Current Research/-Motivation

Proof in Coq

## VERIFICATION OF QUANTUM COMPUTING AND THE BLOCH SPHERE REPRESENTATION

Chris Henson

Drexel University

#### CURRENT RESEARCH/MOTIVATION

CURRENT RESEARCH/-MOTIVATION

SU(2) -SO(3)

Proof I

COQ

- proof assistants are programming languages that allow formal proofs of mathematical theorems
- quantum computing is a great candidate for formal verification: it is both conceptually unintuitive and practically expensive
- algorithms such as Grover, Shor, etc. have been formally verified in languages such as Coq, Isabelle/HOL, etc.
- methods for optimizing general quantum circuits can also be verified
- some formalization frameworks provide the ability to export to formats such as OpenQASM and provide bindings in more accessible languages like Python

#### THE BLOCH SPHERE

 $SU(2) \rightarrow$ SO(3)

- Why do we expect any connection between qubits and a sphere?
- In a sense the relationship between a qubit and a complex number is the same as complex and real numbers




FIGURE: Bloch sphere representation of a Qubit

#### GROUPS

Current Research, Motivatio

### $SU(2) \rightarrow SO(3)$

Proof II Coq

Q&A

#### DEFINITION (GROUPS)

A group is a set G with operation  $\cdot$ , such that:

- $\forall a, b \in G : a \cdot b \in G$
- $\forall a, b, c \in G : (a \cdot b) \cdot c = a \cdot (b \cdot c)$
- $\forall a \in G, \exists 1 \in G : a \cdot 1 = a = 1 \cdot a$
- $\bullet \ \forall a \in G, \exists a^{-1} \in G : a \cdot a^{-1} = \mathbb{1}$

#### DEFINITION (GROUP HOMOMORPHISM)

Given two groups  $(G,\cdot)$  and  $(H,\times)$ , a function  $f:G\to H$  is a homomorphism if  $\forall g_1,g_2\in G: f(g_1\cdot g_2)=f(g_1)\times f(g_2)$ 

#### DEFINITION (GROUP ISOMORPHISM)

A homomorphism  $f: G \to H$  that is bijective (one-to-one and onto) is called an isomorphism

#### GROUPS

Current Research/-Motivation

 $SU(2) \rightarrow SO(3)$ 

Proof in Coq

#### DEFINITION (SPECIAL UNITARY GROUP)

SU(2) is the group of all  $2\times 2$  complex matrices that are unitary  $(A^\dagger A=I_2)$  and have determinant 1. It can be parameterized by

$$A = \begin{pmatrix} \alpha & \beta \\ -\overline{\beta} & \overline{\alpha} \end{pmatrix}, \ |\alpha|^2 + |\beta|^2 = 1$$

#### DEFINITION (SPECIAL ORTHOGONAL GROUP)

SO(3) is the group of all  $3 \times 3$  real matrices that are orthogonal  $(A^{\top}A = I_3)$  and have determinant 1.

#### QUATERNIONS

Current Research/-Motivation

 $SU(2) \rightarrow SO(3)$ 

Proof in Coq

O&A

#### DEFINITION (QUATERNIONS)

A quaternion may be desribed by the expression

$$a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$$
  $a, b, c, d \in \mathbb{R}$ 

where the basis quaternions 1, i, j, k additionally satisfy

$$i^2 = j^2 = k^2 = -1,$$
  
 $ij = -ji = k,$   $jk = -kj = i,$   $ki = -ik = j.$ 

and have 1 as a left and right identity.

#### QUATERNIONS

Current Research/ Motivation

 $SU(2) \rightarrow SO(3)$ 

Proof in Coq

004

- We can also view quaternions as pairs of complex numbers  $(a + bi, c + di) \equiv a + bi + cj + dk$
- This idea of repeatedly taking a product of an algebra with itself is refered to as the *Cayley–Dickson construction*, where at each step we loose some property of the algebra
- Another way of viewing the quaternions that connects with quantum computing is through an isomorphism to the Pauli matrices. One such isomorphism is given by:

$$1 \mapsto I_2$$
,  $\mathbf{i} \mapsto i Z$ ,  $\mathbf{j} \mapsto i Y$ ,  $\mathbf{k} \mapsto i X$ 

- By identifying the *pure imaginary* quaternions (a = 0) with  $\mathbb{R}^3$ , we can model rotations in a numerically stable way
- This is the intuition behind the Bloch sphere, that both qubits and quaternions can be represented by pairs of complex numbers

### $SU(2) \cong \mathbb{S}^3$

Current Research/-Motivation

$$SU(2) \rightarrow$$
  
 $SO(3)$   
PROOF IN

Proof II Coq

More formally, we have that the groups SU(2) and  $\mathbb{S}^3$  of unit quaternions are isomorphic. The isomorphism can be written quite neatly:

$$a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \rightarrow \begin{pmatrix} a + bi & c + di \\ -c + di & a - bi \end{pmatrix}$$

 $SU(2) \rightarrow SO(3)$ 

Proof in Coq

#### DEFINITION (CONJUGATION MAP)

Consider a three dimensional vector represented as a pure imaginary quaternion  $r \in \mathbb{R}^3 \cong \mathbb{R}\mathbf{i} + \mathbb{R}\mathbf{j} + \mathbb{R}\mathbf{k}$  and a unit quaternion  $q \in \mathbb{S}^3$ .

The conjugation of r is given by the map  $\mathbb{R}^3 o \mathbb{R}^3$  given by

$$r o q r q^{-1}$$

and can be shown to be a rotation. Additionally we have that  $qrq^{-1}=(-q)r(-q)^{-1}$ 

 $SU(2) \rightarrow$ SO(3)

For a unit quaternion  $a + b \mathbf{i} + c \mathbf{j} + d \mathbf{k}$ , the homomorphism can also be expressed directly as:

$$\begin{pmatrix} 1-2c^2-2d^2 & 2bc-2da & 2bd+2ca \\ 2bc+2da & 1-2b^2-2d^2 & 2cd-2ba \\ 2bd-2ca & 2cd+2ba & 1-2b^2-2c^2 \end{pmatrix}$$

This was know as early as Euler:

eorum summa

#### PROBLEMATIS INITIO PROPOSITI SOLUTIO GENERALIS IN NUMERIS RATIONALIBUS 33. Coronidis loco solutionem problematis nostri e methodo Diophantea petitam subjungam, quae sequenti modo satis concinne exhiberi potest. Sumantur pro lubitu quatuor numeri p, q, r, s ac posita quadratorum

pp + qq + rr + ss = unovem numeri quaesiti ita determinati reperiuntur')

$$\begin{array}{llll} A = \frac{pp + qq - rr - ss}{u}, & B = -\frac{2qr + 2ps}{u}, & C = -\frac{2qr - 2pr}{u}, \\ D = -\frac{2qr - 2ps}{u}, & E = \frac{pp - qq + rr - ss}{u}, & F = -\frac{2pq + 2rs}{u}, \\ G = -\frac{2qs + 2pr}{u}, & H = -\frac{2rs - 2pq}{u}, & I = \frac{pp - qq - rr + ss}{u}. \end{array}$$

FIGURE: Euler's representation of rotations

# REPRESENTING MATRICES, COMPLEX NUMBERS, AND QUATERNIONS

CURRENT RESEARCH/-MOTIVATION

 $SU(2) \rightarrow SO(3)$ 

Proof in Coq

Coq

- we can represent groups abstractly, but it somewhat interesting to directly work with the underlying types
- complex numbers and quaternions are represented as pairs of real numbers
- a matrix is represented as a function that takes two natural numbers and returns a complex number

 $\texttt{Definition} \ \texttt{C} := (\texttt{R} * \texttt{R}) \% \texttt{type}$ 

Definition Quaternion := (R \* R \* R \* R)%type.

Definition Matrix (m n : nat) := nat  $\rightarrow$  nat  $\rightarrow$  C.

#### Representing Groups

PROOF IN

Coo

- groups are represented as a typeclass
- essentially a typeclass is a dependently-typed record, with some extra type inference capabilities
- a subtle point is that this allows different equivalence relations

```
Class Group (G: Type) (op: G \to G \to G) (rel: relation G) := {
        id · G
       inverse G \to G
      ; rel_equiv: equiv G rel
      ; id_left {x}: rel (op id x) x
      ; id_right {x}: rel (op x id) x
      ; assoc \{x y z\}: rel (op (op x y) z) (op x (op y z))
      ; right_inv {x}: rel (op x (inverse x)) id
```

#### Representing Morphisms

CURRENT RESEARCH/-MOTIVATION

SO(3)

Proof in Coq

Q&A

Likewise, homomorphism and isomorphism can be represented with typeclasses

```
Class GroupHomomorphism G H
  (Gop: G \rightarrow G \rightarrow G)
  (Hop: H \rightarrow H \rightarrow H)
  (Grel: relation G)
  (Hrel: relation H)
  (hom_f: G \rightarrow H)
: Type
    hom_left_group: Group G Gop Grel
  ; hom_right_group: Group H Hop Hrel
  ; hom_mul {a1 a2}: Hrel
                         (hom_f (Gop a1 a2))
                         (Hop (hom_f a1) (hom_f a2))
```

#### Representing Morphisms

PROOF IN Coo

```
Another subtle point about having multiple equivalence
relations is that we must prove they are "compatible"
```

```
Section hom_trans.
Variables A B C : Type.
Variable Arel: relation A
Variable Brel: relation B.
Variable Crel: relation C
Variable A_Eq: Equivalence Arel.
Variable B_Eq: Equivalence Brel.
Variable C Eq: Equivalence Crel.
Variable AtoB: A → B.
Variable BtoC: B → C
Variables Aop: A \rightarrow A \rightarrow A.
Variables Bop: B \rightarrow B \rightarrow B.
Variables Cop: C \rightarrow C \rightarrow C.
Variable BtoC_rw : Proper (Brel == > Crel) BtoC.
Lemma GroupHomomorphism_trans:
 GroupHomomorphism A B Aop Bop Arel Brel AtoB \rightarrow
 GroupHomomorphism B C Bop Cop Brel Crel BtoC →
 GroupHomomorphism A C Aop Cop Arel Crel (fun a ⇒ BtoC (AtoB a)).
Proof.
```

#### Subset Types

Current Research/-Motivation

SU(2) = SO(3)

Proof in Coq

Q&A

- Subset types carry both a value and a proof that the values satisfies some predicate
- any function involving these types also carries a proof of closure

#### SUBSET TYPES

PROOF IN Coo

```
Definition Versor_to_SU2 (v: Versor): SU2.
Proof
 destruct v as [(((x, y), z), w) E].
 unfold SII2
 exists (
   fun row ⇒ fun col ⇒
    match row, col with
    | 0, 0 \Rightarrow (x, y)
    | 0, 1 \Rightarrow (z, w)
    1, 0 \Rightarrow (Ropp z, w)
    | 1, 1 \Rightarrow (x, Ropp y)
    | _, _ ⇒ co
    end
  unfold Onorm in E.
  apply pow_eq with (n := 2%nat) in E.
 rewrite pow2_sqrt in E.
 replace ((1^2)\%R) with 1 in E by 1ra.
 repeat split.
  - show wf.
  — 1 ca
  — 1 ca
  - unfold d2_det, Cmult, Cminus, Cplus, C1.
    simpl. f equal.
    rewrite ← E
    all: lra.
 - repeat apply Rplus_le_le_0_compat.
     all: apply pow2_ge_0.
Defined.
```

#### Subset Types

Current Research/-Motivation

SO(3)

Proof in Coq

- I define equality of these subset types as equality of their values (first projection)
- this is general, as this also takes an equivalence relation

```
Definition sigma_proj1_equiv
{X: Type}
{A: X → Prop}
{rel: relation X}
(e: equiv X rel)
(s1 s2: sig A)
: Prop
:= rel (proj1_sig s1) (proj1_sig s2).
```

#### $SU(2) \rightarrow SO(3)$

CURRENT RESEARCH/-MOTIVATION

 $SU(2) \rightarrow SO(3)$ 

Proof in Coq

```
Q&A
```

```
Theorem SU2 Homomorphism SO3:
  GroupHomomorphism
    SU2
    S03
    SU2 mul
   SO3_mul
   SU2_equiv
    SO3 equiv
    (fun U ⇒ Versor to SO3 (SU2 to Versor U)).
Proof.
  apply GroupHomomorphism trans with versor equiv Vmul.
  - constructor
    + constructor.
    + apply (sigma_proj1_sym eq_equiv).
    + apply (sigma_proj1_trans eq_equiv).

    constructor.

    + constructor.
    + apply (sigma_proj1_sym (mat_equiv_equiv 3 3)).
    + apply (sigma_proj1_trans (mat_equiv_equiv 3 3)).
  - unfold Morphisms.Proper, Morphisms.respectful.
    intros
    unfold versor_equiv, SO3_equiv, sigma_proj1_equiv, proj1_sig in *.
    destruct x as [(((a1, b1), c1), d1) E1].
    destruct y as [(((a2, b2), c2), d2) E2].
    by cell: inversion H: subst: reflexivity.
  - apply SU2_Iso_to_hom.
  - apply Versor_Homomorphism_SO3.
Qed.
```

CURRENT RESEARCH/-MOTIVATION

SU(2) SO(3)

Proof II Coq

Q&A

## Questions?

#### References I

Current Research, Motivatio

SO(3) Proof 1

Coq

Q&A

J.H. Gallier and J. Quaintance, *Linear algebra and optimization with applications to machine learning*, no. v. 1, World Scientific Publishing Company Pte. Limited, 2020.

Kesha Hietala, Robert Rand, Shih-Han Hung, Liyi Li, and Michael Hicks, *Proving Quantum Programs Correct*, 12th International Conference on Interactive Theorem Proving (ITP 2021) (Dagstuhl, Germany) (Liron Cohen and Cezary Kaliszyk, eds.), Leibniz International Proceedings in Informatics (LIPIcs), vol. 193, Schloss Dagstuhl – Leibniz-Zentrum für Informatik, June 2021, pp. 21:1–21:19.

Kesha Hietala, Robert Rand, Shih-Han Hung, Xiaodi Wu, and Michael Hicks, *A verified optimizer for Quantum circuits*, Proceedings of the ACM on Programming Languages **5** (2021), no. POPL, 37, POPL '21.

#### References II

Q&A

Robert Rand, Verified Quantum Computing, https://rand.cs.uchicago.edu/vqc/index.html, 2019.

Alistair Savage, Introduction to Lie Groups, https://alistairsavage.ca/mat4144/notes/

MAT4144-5158-LieGroups.pdf, 2015.

Who discovered the covering homomorphism between su(2) and so(3)?, History of Science and Mathematics Stack Exchange

https://hsm.stackexchange.com/questions/11057/ who-discovered-the-covering-homomorphism-between-su2

2019.

James Wells, Lectures on Standard Model Particle Physics, https://indico.cern.ch/event/243629/, 2013.