

Supplementary Figures: Strain–Doping Coupling in Graphullerene

Xingqiang Chen
(Dated: June 19, 2026)

Companion to the main manuscript on coupled (ϵ, δ) response in qHP C₆₀. Figures follow the observables vocabulary of Capobianco *et al.* [1] (localization, λ ,

J , FCWD). **DATA:** Fig. S5 Marcus λ panels refresh from `exp9_polaron_verification.json` after vertical SP converge; until then literature λ schematics only.

[1] A. Capobianco, A. Breglia, P. Gentile, G. Capano, A. Asinari, G. Drera, L. Sangaletti, M. Cossi, and S. Casassa,

Nano Letters **24**, 7891 (2024), key reference from refer-info.md - polaron transport mechanisms.




FIG. 1. VBM/CBM isosurfaces and π -resolved DOS (Exp. 7). **[pending: VMD]** when PDF absent.




FIG. 2. Marcus configuration-coordinate schematic (Capobianco-style). Verified adiabatic IP/EA from converged Exp. 9 GEO-OPT pairs label the vertical arrows (pristine IP, B EA); λ panels remain **[pending: vertical SP at neutral geometry]**—see `derived.lambda.eV` in audit JSON.




FIG. 3. Nearest-neighbor coupling J from Exp. 4 audit (PBE+D3, two verified tetramer points). [FCWD panel pending MolFC]; legacy figure6_dimer_coupling_fcwd.pdf removed in repo tidy.