Template Document

Academic Support Agent

February 19, 2025

Insert Initials and surnameShort title
Insert title here
First authora, Second authora, and Third authorb
aInsert address here
bInsert address here Corresponding author address. E-mail: xxx@xxxxx.xx.xx

Abstract

Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text. Insert your abstract text.

Insert keywords here
Insert article history

1 Insert A head here

This demo file is intended to serve as a "starter file" for OUP journals produced under LaTeX using oupau.cls.

1.1 Insert B head here

Subsection text here.

1.1.1 Insert C head here

Subsubsection text here.

2 Equations

Sample equations.

$$\begin{aligned}\frac{\partial u(t,x)}{\partial t} &= Au(t,x)\left(1-\frac{u(t,x)}{K}\right) - B\frac{u(t-\tau,x)w(t,x)}{1+Eu(t-\tau,x)},\\ \frac{\partial w(t,x)}{\partial t} &= \delta\frac{\partial^2 w(t,x)}{\partial x^2} - Cw(t,x) + D\frac{u(t-\tau,x)w(t,x)}{1+Eu(t-\tau,x)},\end{aligned} \tag{1}$$

$$\begin{aligned}\frac{dU}{dt} &= \alpha U(t)(\gamma - U(t)) - \frac{U(t-\tau)W(t)}{1+U(t-\tau)},\\ \frac{dW}{dt} &= -W(t) + \beta\frac{U(t-\tau)W(t)}{1+U(t-\tau)}.\end{aligned} \tag{2}$$

$$\frac{\partial(F_1,F_2)}{\partial(c,\omega)}_{(c_0,\omega_0)} = \left|\begin{array}{cc}\frac{\partial F_1}{\partial c} & \frac{\partial F_1}{\partial \omega}\\ \frac{\partial F_2}{\partial c} & \frac{\partial F_2}{\partial \omega}\end{array}\right|_{(c_0,\omega_0)} = -4c_0q\omega_0 - 4c_0\omega_0p^2 = -4c_0\omega_0(q+p^2) > 0.$$

3 Enunciations

Assume that $\alpha > 0, \gamma > 1, \beta > \frac{\gamma+1}{\gamma-1}$. Then there exists a small $\tau_1 > 0$, such that for $\tau \in [0, \tau_1)$, if c crosses $c(\tau)$ from the direction of to a small amplitude periodic traveling wave solution of (2.1), and the period of $(\check{u}^p(s), \check{w}^p(s))$ is

$$\check{T}(c) = c \cdot \left[\frac{2\pi}{\omega(\tau)} + O(c - c(\tau)) \right].$$

From (0.8) and (2.10), it holds $\frac{d\omega}{d\tau} < 0$, $\frac{dc}{d\tau} < 0$ for $\tau \in [0, \tau_1)$. This fact yields that the system (2.1) with delay $\tau > 0$ has the periodic traveling waves for smaller wave speed c than that the system (2.1) with $\tau = 0$ does. That is, the delay perturbation stimulates an early occurrence of the traveling waves.

4 Figures & Tables

The output for figure is:

Figure 1: Insert figure caption here

An example of a double column floating figure using two subfigures. (The subfig.sty package must be loaded for this to work.) The subfigure `\label` commands are set within each subfloat command, the `\label` for the overall figure must come after `\caption`. `\hfil` must be used as a separator to get equal spacing. The subfigure.sty package works much the same way, except `\subfigure` is used instead of `\subfloat`.

The output for table is:

Table 1: An Example of a Table

One	Two
Three	Four

5 Conclusion

The conclusion text goes here.

Acknowledgment

Insert the Acknowledgment text here.

References

[1] Reference details . . .

[2] Reference details