



university of
 groningen

faculty of science
and engineering

bernoulli institute for mathematics,
computer science and artificial intelligence

The differentiation index of DAEs vs. the relative degree of control systems

the 91st GAMM Annual Meeting

Yahao Chen, Stephan Trenn

Jan C. Willems Center for Systems and Control
University of Groningen, Netherlands

A joint work with Witold Respondek

Nonlinear DAEs

Consider a nonlinear differential-algebraic equation (DAE):

$$\Xi : E(x)\dot{x} = F(x), \quad (1)$$

- › where the **generalized state** $x \in X$ and X is an open subset of \mathbb{R}^n ;
- › $E : TX \rightarrow \mathbb{R}^l$ and $F : X \rightarrow \mathbb{R}^l$ are **C^∞ -smooth** maps.
- › The DAE (1) is denoted by $\Xi_{l,n} = (E, F)$.
- › Application: constrained mechanics, electrical circuits, chemical processes, etc.

Solutions of nonlinear DAEs

- › A *solution* of Ξ is a \mathcal{C}^1 -curve $x : I \rightarrow X$ s.t. $E(x(t))\dot{x}(t) = F(x(t))$, $\forall t \in I$.
- › A point x_0 is called *admissible* if \exists a solution $x(\cdot)$ and $t_0 \in I$ s.t. $x(t_0) = x_0$. The *admissible set* (or **consistency set**) $S_a := \{x_0 \mid \exists x(\cdot), t_0 \in I : x(t_0) = x_0\}$.

Definition (geometric reduction method)

Fix $x_p \in X$. Step 0: Set $M_0 = X$, $M_0^c = U_0$. Step k :

$$M_k := \{x \in M_{k-1}^c : F(x) \in E(x)T_x M_{k-1}^c\}. \quad (2)$$

Assume: $x_p \in M_k$ and $M_k^c = M_k \cap U_k$ is a smooth **connected** submanifold.

- › Reich (1991); Rabier and Rheinboldt (2002); Berger(2016,2017); Chen and Trenn (2020); Chen, Trenn and Respondek. (2021).
- › $M^* = M_{k^*+1}^c = M_{k^*}^c$ and $M^* \cap U^* = S_a \cap U^*$ (Chen, Trenn and Respondek. (2021)).

Differentiation index

Definition (differentiation index)

Let $H(x, \dot{x}) = E(x)\dot{x} - F(x)$, define the **k -th order differential array** of $H(x, \dot{x}) = 0$ by

$$H_k(x, x', x^{(2)}, \dots, x^{(k+1)}) = \begin{bmatrix} D_x H x' + D_{x'} H x'' \\ \vdots \\ \frac{d^k}{dt^k} H \end{bmatrix} (x, x', w) = 0, \quad (3)$$

The differentiation index ν_d : the least integer k s.t. (3) **uniquely** determines x' as a function of x , i.e., $x' = v(x)$.

› The vector field v is defined on X or M^* ?

Example

Consider the following DAE $\Xi_{3,3} = (E, F)$ around $x_0 = (x_{10}, x_{20}, x_{30}) = (0, 1, 1) \in X$, where $X = \{x \in \mathbb{R}^3 : x_2 > 0\}$,

$$\begin{bmatrix} x_2 & x_1 & 0 \\ 0 & \sin x_3 & x_2 \cos x_3 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \end{bmatrix} = \begin{bmatrix} \ln x_2 \\ x_2 x_3^2 \cos x_3 \\ x_1 x_2 \end{bmatrix} \quad (4)$$

› We only differentiate the constraints $0 = x_1 x_2$ to have $(x_1 x_2)' = x_2 \dot{x}_1 + x_1 \dot{x}_2 = \ln x_2 = 0$,
 $(x_1 x_2)'' = \frac{1}{x_2} x_2' = 0$.

› The vector field $v(x) = \begin{bmatrix} x_2 & x_1 & 0 \\ 0 & \sin x_3 & x_2 \cos x_3 \\ 0 & \frac{1}{x_2} & 0 \end{bmatrix}^{-1} \begin{bmatrix} \ln x_2 \\ x_2 x_3^2 \cos x_3 \\ 0 \end{bmatrix}$.

› Ξ is of differentiation index-2, i.e. $\nu_d = 2$.

› $M_1^c := \{x \in X : x_1 x_2 = 0\}$; $M_2^c := \{x \in M_1^c : \ln x_2 = 0\}$; $M^* = M_3^c = M_2^c$. Solutions exists on M^* only, $v(x)|_{M^*} = \begin{bmatrix} 0 \\ 0 \\ x_3^2 \end{bmatrix}$.

› The solution of Ξ (passing through x_0) is $(x_1(t), x_2(t), x_3(t)) = (0, 1, \frac{1}{1-t})$.

The explicitation of nonlinear DAEs

- › For a DAE $\Xi_{l,n} = (E, F)$, assume (on a nbh. U of x_p) $\text{rank } E(x) = \text{const.} = q$.
- › $\exists Q : U \rightarrow GL(l, \mathbb{R})$ s.t. $QE(x)\dot{x} = QF(x)$:

$$\begin{cases} E_1(x)\dot{x} = F_1(x), \\ 0 = F_2(x). \end{cases} \quad (5)$$

where $E_1(x)$ of $Q(x)E(x) = \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix}$ is of **full row rank** and $Q(x)F(x) = \begin{bmatrix} F_1(x) \\ F_2(x) \end{bmatrix}$.

- › **(crucial)** The collection of all \dot{x} satisfying $E_1(x)\dot{x} = F_1(x)$ of (5) is given by the differential inclusion:

$$\dot{x} \in f(x) + \ker E_1(x) = f(x) + \ker E(x). \quad (6)$$

where $f(x) = E_1^\dagger F_1(x)$, where E_1^\dagger is a **right inverse** of E_1 .

The explicitation of nonlinear DAEs

- › Let $m = n - q$ and $g_1, \dots, g_m : X \rightarrow \mathbb{R}^n$ s.t. $\ker E(x) = \text{span} \{g_1, \dots, g_m\}(x)$. By introducing **driving variables** v_i , $i = 1, \dots, m$, we get

$$\dot{x} = f(x) + \sum_{i=1}^m g_i(x)v_i. \quad (7)$$

- › All solutions of Ξ are in **one-to-one correspondence** with all solutions (**corresponding to all C^0 -controls $v(t)$**) of

$$\begin{cases} \dot{x} = f(x) + g(x)v, \\ 0 = h(x). \end{cases} \quad (8)$$

where $h(x) = F_2(x)$.

- › To (8), we attach the control system $\Sigma = \Sigma_{n,m,p} = (f, g, h)$, given by

$$\Sigma : \begin{cases} \dot{x} = f(x) + g(x)v, \\ y = h(x), \end{cases} \quad (9)$$

where $n = \dim x$, $m = \dim v = n - q$, $p = \dim y = l - q$.

The explicitation of nonlinear DAEs

Definition (explicitation with driving variables)

Given a DAE $\Xi_{l,n} = (E, F)$, by a (Q, v) -explicitation, we will call a control system $\Sigma = \Sigma_{n,m,p} = (f, g, h)$ given by

$$\Sigma : \begin{cases} \dot{x} = f(x) + g(x)v, \\ y = h(x), \end{cases}$$

with

$$f(x) = E_1^\dagger F_1(x), \quad \text{Im } g(x) = \ker E(x), \quad h(x) = F_2(x),$$

where $QE(x) = \begin{bmatrix} E_1(x) \\ 0 \end{bmatrix}$, $QF(x) = \begin{bmatrix} F_1(x) \\ F_2(x) \end{bmatrix}$.

› Σ is **not uniquely defined** since $Q(x)$, $E_1^\dagger(x)$ and $g(x)$ are **not** unique !

The explicitation of nonlinear DAEs

Proposition

Assume: $\Sigma_{n,m,p} = (f, g, h)$, $\tilde{\Sigma}_{n,m,p} = (\tilde{f}, \tilde{g}, \tilde{h})$ are two (Q, v) -explicitations of a DAE $\Xi_{l,n} = (E, F)$ corresponding to *different choices* of Q , E_1^\dagger and g .

Then $\exists \alpha, \gamma$ and invertible η, β , which map

$$f \mapsto \tilde{f} = f + \gamma h + g\alpha, \quad g \mapsto \tilde{g} = g\beta, \quad h \mapsto \tilde{h} = \eta h.$$

- › Σ is defined up to a **feedback transformation** $v = \alpha + \beta\tilde{v}$, an **output injection** γy and an **output multiplication** ηh :

$$\tilde{\Sigma} : \begin{cases} \dot{x} = f(x) + g(x)(\alpha(x) + \beta(x)\tilde{v}) + \gamma(x)h(x) \\ y = \eta(x)h(x) \end{cases}$$

- › The (Q, v) -explicitation of Ξ is a class, denoted $\mathbf{Expl}(\Xi)$. We write $\Sigma \in \mathbf{Expl}(\Xi)$.

External equivalence and system equivalence

Definition (external equivalence)

Two DAEs $\Xi_{l,n} = (E, F)$ and $\tilde{\Xi}_{l,n} = (\tilde{E}, \tilde{F})$ are called **externally equivalent**, shortly ex-equivalent, if \exists diffeomorphism $\psi : X \rightarrow \tilde{X}$ and $Q : X \rightarrow GL(l, \mathbb{R})$ s.t.

$$\tilde{E}(\psi(x)) = Q(x)E(x) \left(\frac{\partial \psi(x)}{\partial x} \right)^{-1} \quad \text{and} \quad \tilde{F}(\psi(x)) = Q(x)F(x). \quad (10)$$

Definition (system equivalence)

Two control systems $\Sigma_{n,m,p} = (f, g, h)$ and $\tilde{\Sigma}_{n,m,p} = (\tilde{f}, \tilde{g}, \tilde{h})$ are called **system equivalent**, or shortly sys-equivalent if

$$\tilde{f} \circ \psi = \frac{\partial \psi}{\partial x} (f + \gamma h + g\alpha), \quad \tilde{g} \circ \psi = \frac{\partial \psi}{\partial x} g\beta, \quad \tilde{h} \circ \psi = \eta h.$$

External equivalence and system equivalence

Theorem

Assume: $\text{rank } E(x)$ and $\text{rank } \tilde{E}(\tilde{x})$ are **constant** around two points x_p and \tilde{x}_p , respectively, for two DAEs $\Xi_{l,n} = (E, F)$ and $\tilde{\Xi}_{l,n} = (\tilde{E}, \tilde{F})$.

Then for two systems $\Sigma_{n,m,p} = (f, g, h) \in \mathbf{Expl}(\Xi)$ and $\tilde{\Sigma}_{n,m,p} = (\tilde{f}, \tilde{g}, \tilde{h}) \in \mathbf{Expl}(\tilde{\Xi})$, we have that locally

$$\Xi \stackrel{ex}{\sim} \tilde{\Xi} \Leftrightarrow \Sigma \stackrel{sys}{\sim} \tilde{\Sigma}.$$

Example

$$\Xi = (E, F) \stackrel{ex}{\sim} \tilde{\Xi} : \left\{ \begin{array}{l} \dot{x}_1 = F_1(x_1) \\ 0 = x_2^1 \\ \dot{x}_2^1 = x_2^1 \\ \vdots \\ \dot{x}_2^{\rho-1} = x_2^{\rho} \end{array} \right. \Leftrightarrow \Sigma = (f, g, g) \in \mathbf{Expl}(\Xi) \stackrel{sys}{\sim} \tilde{\Sigma} : \left\{ \begin{array}{l} \dot{x}_1 = F_1(x_1) \\ y = x_2^1 \\ \dot{x}_2^1 = x_2^1 \\ \vdots \\ \dot{x}_2^{\rho-1} = x_2^{\rho} \\ \dot{x}_2^{\rho} = v \end{array} \right.$$

Differentiation index vs. relative degree

Definition (relative degree)

$\Sigma_{n,m,m} = (f, g, h)$ has a (vector) relative degree $\rho = (\rho_1, \dots, \rho_m)$ at a point x_0 if (i) $L_g L_f^k h(x) = 0$, for all $1 \leq j \leq m$, for all $k < \rho_i - 1$ and $1 \leq i \leq m$, and for all x around x_0 ;
(ii) $D(x_0) = \left(L_{g_j} L_f^{\rho_i - 1} h_i(x_0) \right)_{i,j=1,\dots,m}$ is invertible.

- › Any control system $\Sigma = (f, g, h)$ with relative degree $\rho = (\rho_1, \dots, \rho_m)$ is **feedback equivalent** to the Byrnes-Isidori form

$$\text{B-I : } \begin{cases} \dot{z} &= f_0(z, \xi_1, \dots, \xi_m) + g_0(z, \xi_1, \dots, \xi_m)v \\ y_i &= \xi_i^1, \quad i = 1, \dots, m, \\ \dot{\xi}_i^1 &= \xi_i^2 \\ &\vdots \\ \dot{\xi}_i^{\rho_i - 1} &= \xi_i^{\rho_i} \\ \dot{\xi}_i^{\rho_i} &= v_i \end{cases}$$

Theorem

Assume: *an* explicitaion $\Sigma \in \mathbf{Expl}(\Xi)$ has *a well-defined relative degree* $\rho = (\rho_1, \dots, \rho_m)$ at x_0 , then

(i) *any* $\tilde{\Sigma} \in \mathbf{Expl}(\Xi)$ has *either the same* relative degree ρ with Σ or *no well-defined relative degree* at x_0 ;

(ii) $\nu_d = \max \{\rho_1, \dots, \rho_m\}$,

(iii) Ξ is ex-equivalent to

$$\begin{cases} \dot{z} - g_0(z, \xi) \dot{\xi}^\rho = f_0(z, \xi) \\ N \dot{\xi} = \xi \end{cases}$$

where $N = \text{diag}(N_1, \dots, N_m)$, and where N_i , $i = 1, \dots, m$ are *nilpotent matrices of index* ρ_i .

Continuation of the Example

$$\Xi : \begin{bmatrix} x_2 & x_1 & 0 \\ 0 & \sin x_3 & x_2 \cos x_3 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \end{bmatrix} = \begin{bmatrix} \ln x_2 \\ x_2 x_3^2 \cos x_3 \\ x_1 x_2 \end{bmatrix},$$

› A control system

$$\Sigma = (f, g, h) \in \mathbf{Expl}(\Xi) : \begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \\ \dot{x}_3 \end{bmatrix} = \begin{bmatrix} \frac{\ln x_2}{x_2} \\ 0 \\ x_2^2 \end{bmatrix} + \begin{bmatrix} x_1 \\ -x_2 \\ \tan x_3 \end{bmatrix} v, \quad y = x_1 x_2$$

› The relative degree of Σ at $x_0 = (0, 1, 1)$ is $\rho = 2$.

$$\Xi \stackrel{ex}{\sim} \begin{cases} \dot{z} - \tan z \dot{\xi}_2 = z^2 \\ 0 = \xi_1 \\ \dot{\xi}_1 = \xi_2 \end{cases} \Leftrightarrow \Sigma \stackrel{sys}{\sim} \begin{cases} \dot{z} = z^2 + \tan z v \\ \dot{\xi}_1 = \xi_2 \\ \dot{\xi}_2 = v \\ y = \xi_1 \end{cases}$$

Summary

- › Geometric reduction method and differential array to solve DAEs.
- › A universal way to connected nonlinear DAEs with nonlinear control systems: **the explicitation with driving variables**.
- › **External equivalence** of DAEs and **system equivalence** of control systems.
- › **The differentiation index** of DAEs and the **relative degree** of control systems.
- › Normal forms (nonlinear generalizations of the Weierstrass form ?).