

### Quiz5

- 1) (2pts) Given four documents A, B, C, and D and their top two TF-IDF words, A: nba, basketball; B: cancer, health; C: vote, democratic; D: basketball, baseball, write the Boolean feature vectors for each document (1pt) and calculate the cosine similarity between A, D (1pt)

$$\text{similarity} = \cos(\theta) = \frac{A \cdot B}{\|A\| \|B\|} = \frac{\sum_{i=1}^n A_i \times B_i}{\sqrt{\sum_{i=1}^n (A_i)^2} \times \sqrt{\sum_{i=1}^n (B_i)^2}}$$

Feature vector (nba, basketball, cancer, health, vote, democratic, baseball)

	nba	basketball	cancer	health	vote	democratic	Baseball
A	1	1	0	0	0	0	0
B	0	0	1	1	0	0	0
C	0	0	0	0	1	1	0
D	0	1	0	0	0	0	1

$$\text{Cosine Similarity}(A,D) = 1/(\sqrt{2} \times \sqrt{2}) = 1/2$$

- 2) (2pts) Briefly explain one advantage and one disadvantage of 1) using the content-based approach for finding recommendations and 2) using the dimensionality reduction techniques in your CF recommendation systems.

1. Advantage: No need for data on other users, Able to recommend to users with unique tastes, Able to recommend new & unpopular items, Able to provide explanations

Disadvantage: Finding the appropriate features is hard, Recommendations for new users, Overspecialization

2. Advantage: Reduce the computation

Disadvantage: May lose useful information

3)

$$1. w_{2,3} = \frac{(4-2.5)(3-2.5) + (1-2.5)(2-2.5)}{\sqrt{(4-2.5)^2 + (1-2.5)^2} \times \sqrt{(3-2.5)^2 + (2-2.5)^2}} = 1$$

$$w_{3,4} = 0$$

$$w_{3,5} = \frac{(3-3)(2-\frac{10}{3}) + (2-3)(3-\frac{10}{3}) + (4-3)(5-\frac{10}{3})}{\sqrt{(3-3)^2 + (2-3)^2 + (4-3)^2} \times \sqrt{(2-\frac{10}{3})^2 + (3-\frac{10}{3})^2 + (5-\frac{10}{3})^2}} = 0.65$$

$$P_{2,3} = \frac{(2-\frac{5}{2}) \times 1 + (4-4) \times 0 + (1-\frac{10}{3}) \times 0.65}{1+0+0.65} + 3 = 1.78 \text{ (1pt, no need to show the result)}$$

4)

(1) average ratings based on all ratings

$$w_{1,3} = \frac{\left(3 - \frac{10}{3}\right)\left(5 - \frac{8}{3}\right) + \left(5 - \frac{10}{3}\right)\left(1 - \frac{8}{3}\right)}{\sqrt{\left(3 - \frac{10}{3}\right)^2 + \left(5 - \frac{10}{3}\right)^2} \times \sqrt{\left(5 - \frac{8}{3}\right)^2 + \left(1 - \frac{8}{3}\right)^2}} = -0.73$$

$$w_{2,3} = \frac{\left(4 - \frac{8}{3}\right)\left(2 - \frac{8}{3}\right) + \left(3 - \frac{8}{3}\right)\left(1 - \frac{8}{3}\right)}{\sqrt{\left(4 - \frac{8}{3}\right)^2 + \left(3 - \frac{8}{3}\right)^2} \times \sqrt{\left(2 - \frac{8}{3}\right)^2 + \left(1 - \frac{8}{3}\right)^2}} = -0.58$$

$$w_{3,4} = \frac{\left(5 - \frac{8}{3}\right)\left(2 - \frac{8}{3}\right) + \left(2 - \frac{8}{3}\right)\left(3 - \frac{8}{3}\right)}{\sqrt{\left(5 - \frac{8}{3}\right)^2 + \left(2 - \frac{8}{3}\right)^2} \times \sqrt{\left(2 - \frac{8}{3}\right)^2 + \left(3 - \frac{8}{3}\right)^2}} = -0.98$$

(0.5pt, no need to show the result to get full credit)

$$P_{1,3} = \frac{2 \times w_{1,3} + 1 \times w_{2,3}}{|w_{1,3}| + |w_{2,3}|} \quad (0.5pt, no need to show the result.)$$

(2) average ratings for co-rated items

$$w_{3,1} = \frac{(3-4)(5-3) + (5-4)(1-3)}{\sqrt{(3-4)^2 + (5-4)^2} \times \sqrt{(5-3)^2 + (1-3)^2}} = -1$$

$$w_{3,2} = \frac{(4-7/2)(2-3/2) + (3-7/2)(1-3/2)}{\sqrt{(4-7/2)^2 + (3-7/2)^2} \times \sqrt{(2-3/2)^2 + (1-3/2)^2}} = 1$$

$$w_{3,4} = \frac{(5-7/2)(2-5/2) + (2-7/2)(3-5/2)}{\sqrt{(5-7/2)^2 + (2-7/2)^2} \times \sqrt{(2-5/2)^2 + (3-5/2)^2}} = -1$$

$$P_{1,3} = \frac{1 \times w_{3,2} + 2 \times w_{3,1}}{|w_{3,2}| + |w_{3,1}|} \quad \text{or} \quad P_{1,3} = \frac{1 \times w_{3,2} + 3 \times w_{3,4}}{|w_{3,2}| + |w_{3,4}|}$$

5) Inverse frequency.  $fi = \log(N/ni)$  (1pt)

e.g) clustering --- get rid of the outsiders. Case Amplification (1pt)

6) Advantage: Simple representation

Disadvantage: Complicated for large dataset