

Linear Regression with Multiple Variables

Quiz, 5 questions

1
point

1. Suppose $m=4$ students have taken some class, and the class had a midterm exam and a final exam. You have collected a dataset of their scores on the two exams, which is as follows:

midterm exam	(midterm exam) ²	final exam
89	7921	96
72	5184	74
94	8836	87
69	4761	78

You'd like to use polynomial regression to predict a student's final exam score from their midterm exam score. Concretely, suppose you want to fit a model of the form $h_{\theta}(x) = \theta_0 + \theta_1 x_1 + \theta_2 x_2$, where x_1 is the midterm score and x_2 is (midterm score)². Further, you plan to use both feature scaling (dividing by the "max-min", or range, of a feature) and mean normalization.

What is the normalized feature $x_2^{(2)}$? (Hint: midterm = 72, final = 74 is training example 2.) Please round off your answer to two decimal places and enter in the text box below.

-0.37

1
point

2. You run gradient descent for 15 iterations with $\alpha = 0.3$ and compute

$J(\theta)$ after each iteration. You find that the

value of $J(\theta)$ **decreases slowly** and is still

decreasing after 15 iterations. Based on this, which of the

following conclusions seems most plausible?

- ☒ Rather than use the current value of α , it'd be more promising to try a larger value of α (say $\alpha = 1.0$).
- ☐ $\alpha = 0.3$ is an effective choice of learning rate.
- ☐ Rather than use the current value of α , it'd be more promising to try a smaller value of α (say $\alpha = 0.1$).

1
point

3. Suppose you have $m = 28$ training examples with $n = 4$ features (excluding the additional all-ones feature for the intercept term, which you should add). The normal equation is $\theta = (X^T X)^{-1} X^T y$. For the given values of m and n , what are the dimensions of θ , X , and y in this equation?

- ☐ X is 28×4 , y is 28×1 , θ is 4×4
- ☒ X is 28×5 , y is 28×1 , θ is 5×1
- ☐ X is 28×5 , y is 28×5 , θ is 5×5
- ☐ X is 28×4 , y is 28×1 , θ is 4×1

1
point

4. Suppose you have a dataset with $m = 50$ examples and $n = 15$ features for each example. You want to use multivariate linear regression to fit the parameters θ to our data. Should you prefer gradient descent or the normal equation?

- ☒ The normal equation, since it provides an efficient way to directly find the solution.

- ☐ The normal equation, since gradient descent might be unable to find the optimal θ .
 - ☐ Gradient descent, since $(X^T X)^{-1}$ will be very slow to compute in the normal equation.
 - ☐ Gradient descent, since it will always converge to the optimal θ .
-

1
point

5. Which of the following are reasons for using feature scaling?

- ☐ It is necessary to prevent the normal equation from getting stuck in local optima.
 - ☐ It speeds up gradient descent by making it require fewer iterations to get to a good solution.
 - ☐ It prevents the matrix $X^T X$ (used in the normal equation) from being non-invertable (singular/degenerate).
 - ☐ It speeds up gradient descent by making each iteration of gradient descent less expensive to compute.
-

☒ I, **Zhaiyu Chen**, understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account.

[Learn more about Coursera's Honor Code](#)

Submit Quiz