


← Support Vector Machines

Quiz, 5 questions

1
point

1. Suppose you have trained an SVM classifier with a Gaussian kernel, and it learned the following decision boundary on the training set:



When you measure the SVM's performance on a cross validation set, it does poorly. Should you try increasing or decreasing C ? Increasing or decreasing σ^2 ?

- ☐ It would be reasonable to try **increasing** C . It would also be reasonable to try **increasing** σ^2 .
- ☐ It would be reasonable to try **decreasing** C . It would also be reasonable to try **decreasing** σ^2 .
- ☒ It would be reasonable to try **decreasing** C . It would also be reasonable to try **increasing** σ^2 .
- ☐ It would be reasonable to try **increasing** C . It would also be reasonable to try **decreasing** σ^2 .

1
point


- 2.

The formula for the Gaussian kernel is given by $\text{similarity}(x, l^{(1)}) = \exp\left(-\frac{\|x - l^{(1)}\|^2}{2\sigma^2}\right)$.

Support Vector Machines

The figure below shows a plot of $f_1 = \text{similarity}(x, l^{(1)})$ when $\sigma^2 = 1$.

Quiz, 5 questions



Which of the following is a plot of f_1 when $\sigma^2 = 0.25$?





Figure 3.





Support Vector Machines

Quiz, 5 questions




Figure 2.




Figure 4.


1
point

3.

The SVM solves

$$\min_{\theta} C \sum_{i=1}^m y^{(i)} \text{cost}_1(\theta^T x^{(i)}) + (1 - y^{(i)}) \text{cost}_0(\theta^T x^{(i)}) + \sum_{j=1}^n \theta_j^2$$

Quiz, 5 questions
where the functions $\text{cost}_0(z)$ and $\text{cost}_1(z)$ look like this:



The first term in the objective is:

$$C \sum_{i=1}^m y^{(i)} \text{cost}_1(\theta^T x^{(i)}) + (1 - y^{(i)}) \text{cost}_0(\theta^T x^{(i)}).$$

This first term will be zero if two of the following four conditions hold true. Which are the two conditions that would guarantee that this term equals zero?

- ☒ For every example with $y^{(i)} = 0$, we have that $\theta^T x^{(i)} \leq -1$.
- ☐ For every example with $y^{(i)} = 0$, we have that $\theta^T x^{(i)} \leq 0$.
- ☒ For every example with $y^{(i)} = 1$, we have that $\theta^T x^{(i)} \geq 1$.
- ☐ For every example with $y^{(i)} = 1$, we have that $\theta^T x^{(i)} \geq 0$.

1
point

4.
Suppose you have a dataset with $n = 10$ features and $m = 5000$ examples.

After training your logistic regression classifier with gradient descent, you find that it has underfit the training set and does not achieve the desired performance on the training or cross validation sets.

Which of the following might be promising steps to take? Check all that apply.

- ☐ Use an SVM with a linear kernel, without introducing new features.
- ☒ Use an SVM with a Gaussian Kernel.
- ☒ Create / add new polynomial features.
- ☐ Increase the regularization parameter λ .

1
point

5.
Which of the following statements are true? Check all that apply.

- ☐ If the data are linearly separable, an SVM using a linear kernel will return the same parameters θ regardless of the chosen value of C (i.e., the resulting value of θ does not depend on C).
- ☐ Suppose you are using SVMs to do multi-class classification and would like to use the one-vs-all approach. If you have K different classes, you will train $K - 1$ different SVMs.
- ☒ It is important to perform feature normalization before using the Gaussian kernel.



The maximum value of the Gaussian kernel (i.e., $\text{sim}(x, l^{(1)})$) is 1.



Support Vector Machines

Quiz, 5 questions



I, **Zhaiyu Chen**, understand that submitting work that isn't my own may result in permanent failure of this course or deactivation of my Coursera account.

[Learn more about Coursera's Honor Code](#)

Submit Quiz

