

Exploring predators as samplers

How can we use diet data to quantify prey distribution and density?

Peri E. Gerson*, Cheryl L. Barnes

Coastal Oregon Marine Experiment Station, Oregon State University

* presenting author

Background

Prey availability is an important factor influencing habitat quality for predator species. Distributions of groundfish predators are also known to respond to localized prey abundances. Currently, standardized surveys in the Gulf of Alaska (GOA) do not adequately sample prey species (e.g., euphausiids, forage fishes, benthic invertebrates) and the capacity to expand survey scope is limited. Predator diet data has been used to quantify spatial and temporal trends of prey in other systems¹⁻⁴. This project aims to utilize readily available diet data from the National Oceanic and Atmospheric Administration (NOAA), Alaska Fisheries Science Center's (AFSC) bottom trawl survey to assess prey distribution and density in the GOA.

Research Objectives

- Use diet data from groundfish predators to quantify spatiotemporal variation in prey densities throughout the GOA.
- Utilize a multivariate statistical framework – joint species distribution models (JSDMs) – to model prey taxa. JSDMs provide a multivariate context to quantify species correlations and improve model performance for more rare taxa.
- Create prey informed species distribution models (SDMs) for commercially important groundfish in the GOA.
- Quantify differences between prey informed and non-prey informed predator groundfish SDMs.

References

1. Gaichas SK, Gartland J, Smith BE, Wood A, Ng E, Celestino M, et al. Assessing small pelagic fish trends in space and time using piscivore stomach contents. *Can J Fish Aquat Sci* [Internet]. 2023 Oct 4 [cited 2023 Dec 15]; Available from: <https://cdnscepub.com/doi/abs/10.1139/cjfas-2023-0093>
2. Ng EL, Deroba JJ, Essington TE, Grüss A, Smith BE, Thorson JT. Predator stomach contents can provide accurate indices of prey biomass. *ICES Journal of Marine Science*. 2021 Jul 1;78(3):1146-59.
3. Gunther KM, Baker MR, Aydin KY. Using predator diets to infer forage fish distribution and assess responses to climate variability in the eastern Bering Sea. *Marine Ecology Progress Series* [Internet]. 2023 Sep 28 [cited 2024 Feb 13];SPF2. Available from: https://www.int-res.com/meeps/SPF2/p_av3/
4. Fahrig L, Lilly GR, Miller DS. Predator Stomachs as Sampling Tools for Prey Distribution: Atlantic Cod (*Gadus morhua*) and Capelin (*Mallotus villosus*). *Can J Fish Aquat Sci*. 1993 Jul;50(7):1541-7.
5. Chipp SR, Garvey JE. 11. Assessment of Food Habits and Feeding Patterns.
6. Thorson, in development

Acknowledgements




Financial and logistical support were provided by discretionary funds from NOAA's Alaska Regional Office (AKRO) and an EFH RFP administered by AKRO and the Alaska Fisheries Science Center (AFSC). Conceptual and analytical support will be provided by co-PI Jim Thorson (AFSC), committee member Jodi Pirtle (AKRO), committee member Jessica Miller (OSU), collaborator Jon Reum (AFSC), and collaborator Lauren Rogers (AFSC). Fish images provided by Larry Allen (California State University Northridge).



Oregon State University
Coastal Oregon Marine
Experiment Station



NOAA
FISHERIES



Predator feeding strategy diagram⁵

Diet data subset to 50% length at maturity for Pacific cod, pollock, and arrowtooth flounder. All samples retained for sablefish and Pacific halibut.

$$\text{Frequency of Occurrence} = O_i = \frac{J_i}{P}$$

$$J_i = \text{number of fish containing prey } i$$

$$P = \text{number of fish with food in their stomachs}$$


$$\text{Prey-specific Abundance} = P_i = \frac{S_{i,t}}{S_{t,t}} \cdot 100$$

$$S_{i,t} = \text{grams of prey } i \text{ in stomachs}$$

$$S_{t,t} = \text{total grams of prey in predators that contain prey } i$$

Workflow Diagram


Grams of prey per grams of predator, weighted for predator size representation, will be used as the sampling unit for each unique predator-haul combination.



Joint species distribution models (JSDMs) will be used to model prey densities as a function of predator species and size, along with a suite of environmental covariates (e.g., depth, occurrence of structure-forming invertebrates). Model development will be done using *tinyVAST*⁶.



Prey distributions and densities will be predicted for the GOA from each predator's sampling perspective.



Predator-specific prey predictions will be combined in a composite index to provide interspecies corroborations of prey distribution and density.

