MIE286 Lecture Notes

Probability and Statistics

Cheryl Shi

Contents

1 Chapter 1: Introduction to Statistics **2**

1.1 Statistics Definitions . . . 2
1.2 Sampling Methods . . . 2
1.3 Data, Variables, and Distributions . . . 3
1.4 Describing Distributions with Numbers . . . 3
1.5 Measures of Spread . . . 5
1.6 Graphical Representations of Data . . . 6

2 Chapter 2: Probability **10**

2.1 Experiments, Sample Spaces, and Events . . . 10
2.2 Event Operations and Probability Rules . . . 11
2.3 Probability . . . 12
2.4 Mutually Exclusive Events . . . 13
2.5 Counting Techniques and Equally Likely Outcomes . . . 15
2.6 Conditional Probability and Independence . . . 19
2.7 Law of Total Probability . . . 20
2.8 Independence of Events . . . 22

3 Chapter 3: Random Variables and Probability Distributions **28**

3.1 Random Variables and Their Interpretation . . . 28
3.2 Discrete Random Variables . . . 29
3.3 Probability Mass Function (PMF) . . . 30
3.4 Bernoulli and Binomial Random Variables . . . 31
3.5 Cumulative Distribution Function (CDF) . . . 33
3.6 Continuous Sample Space and Continuous Random Variables . . . 35
3.7 Probability Density Function (PDF) . . . 35
3.8 CDF for Continuous Random Variables . . . 38
3.9 Joint Probability Distributions (Discrete) . . . 39
3.10 Conditional Distributions (Discrete) . . . 42
3.11 Joint Probability Density Function (Continuous) . . . 43
3.12 Marginal and Conditional PDFs . . . 46
3.13 Statistical Independence . . . 47

4 Chapter 4: Mathematical Expectation **54**

4.1 Variance . . . 60

4.2 Covariance and Independence . . . 66

5 Chapter 5: Discrete Distributions 76

5.1 Discrete Uniform Distribution . . . 76
5.2 Bernoulli Distribution . . . 76
5.3 Binomial Distribution . . . 77
5.4 Multinomial Distribution . . . 80
5.5 Hypergeometric Distribution . . . 83
5.6 Negative Binomial Distribution . . . 88
5.7 Geometric Distribution . . . 90
5.8 Poisson Process . . . 92

6 Chapter 6: Continuous Probability Distributions 96

6.1 Uniform Distribution . . . 96
6.2 Normal Distribution . . . 96
6.3 Standardization (Z-Scores) . . . 98
6.4 Computing Normal Probabilities . . . 100
6.5 The Empirical Rule (68–95–99.7 Rule) . . . 101
6.6 Percentiles of the Normal Distribution . . . 101
6.7 Normal Approximation to the Binomial . . . 102
6.8 Gamma & Exponential Distributions . . . 104
6.9 Weibull Distribution, Reliability, Failure Rates & Chi-Squared Distribution . . . 107

7 Chapter 7: Transformations of Random Variables 113

7.1 Functions of Random Variables . . . 113
7.2 Moments and Moment Generating Functions (MGFs) . . . 120
7.3 Linear Combinations of Normal Random Variables . . . 122

Chapter 1

Chapter 1: Introduction to Statistics

1.1 Statistics Definitions

Global definition: Statistics involves collecting, organizing, summarizing, presenting, and analyzing data, as well as making inferences, conclusions, and decisions based on data.

Statistical definition: A statistic is a numerical value calculated from data (e.g. mean, proportion, standard deviation).



1.1.1 Basic Terminology

Individuals: Objects on which data are collected (people, animals, plots of land, etc.).
Variable: Any characteristic of an individual.
Population: The entire group of individuals of interest.
Sample: A subset of individuals taken from the population.
Statistical Inference: Drawing conclusions about a population based on a sample.

1.2 Sampling Methods

Simple Random Sample (SRS):

- Every possible group of size n has an equal chance of being selected.
- Helps avoid bias in sampling.
- Can be selected using random number tables or software.

Stratified Random Sampling:

- The population is divided into homogeneous groups *(individuals are similar with respect to the variable being studied)* called strata.
- A simple random sample is taken from each stratum. *(one subgroup of the population created)*
- Ensures that important subgroups are neither over nor under represented.

1.3 Data, Variables, and Distributions

1.3.1 Types of Variables

Categorical Variable: Places individuals into categories (e.g. gender, major). These are qualitative.

Quantitative Variable: Takes numerical values for which arithmetic operations are meaningful.

- Discrete
- Continuous

1.3.2 Distributions

Distribution: Describes what values a variable takes and how often those values occur. When examining a distribution, look for:

- **Shape**
- **Center**
- **Spread**
- **Outliers**

Outlier: An individual value that falls outside the overall pattern of the data.

1.4 Describing Distributions with Numbers

Central Tendency: Describes where the data cluster or center.

Central Tendency: Describes where the data cluster or center.

- Mean: average value
- Median: middle value

Mean (Arithmetic Mean):

$$\boxed{\bar{x} = \frac{1}{n}\sum_{i=1}^{n} x_i}$$

Median:

$$\boxed{\tilde{x} = \begin{cases} x_{\left(\frac{n+1}{2}\right)}, & \text{if } n \text{ is odd} \\ \dfrac{x_{\left(\frac{n}{2}\right)} + x_{\left(\frac{n}{2}+1\right)}}{2}, & \text{if } n \text{ is even} \end{cases}}$$

Theorem 1.1

1. The mean is more sensitive to extreme values than the median.
2. Changing a single data value will always change the mean, but may not change the median.
3. If a distribution is exactly symmetric, the mean and median are equal.

Trimmed Mean: The mean computed after removing extreme values.

$$\boxed{\bar{x}_{\text{trim}} = \frac{1}{n-2k} \sum_{i=k+1}^{n-k} x_{(i)}}$$

where k values are removed from both ends of the ordered data. (normally given in question like 10%)

Exercise 1.3 — Tensile Strength and Aging

Twenty specimens of a polymer were randomly selected. Ten were tested as-is and ten were exposed to an accelerated aging process (high temperatures for 10 days). Tensile strength (psi) was recorded:

No aging:	227	222	218	217	225
	218	216	229	228	221
Aging:	219	214	215	211	209
	218	203	204	201	205

1. Do a dot plot of the data.
2. From your plot, does it appear as if the aging process has had an effect on the tensile strength of this polymer? Explain.
3. Calculate the sample mean tensile strength of the two samples.
4. Calculate the median for both. Discuss the similarity or lack of similarity between the mean and median of each group.

Exercise 1.4 — Steel Rod Flexibility

In a study by the Department of Mechanical Engineering at Virginia Tech, steel rods from two companies were compared. Ten sample springs were made from each company's rods, and flexibility was recorded:

Company A:	9.3	8.8	6.8	8.7	8.5
	6.7	8.0	6.5	9.2	7.0
Company B:	11.0	9.8	9.9	10.2	10.1
	9.7	11.0	11.1	10.2	9.6

1. Calculate the sample mean and median for the data for the two companies.
2. Plot the data for the two companies on the same line and give your impression regarding any apparent differences between the two companies.

Exercise 1.5 — Cholesterol Reduction

Twenty adult males (ages 30–40) participated in a study on the effect of diet and exercise on blood cholesterol. Ten were controls and ten followed a treatment regimen for 6 months. Reduction in cholesterol:

Control:	7	3	−4	14	2
	5	22	−7	9	5
Treatment:	−6	5	9	4	4
	12	37	5	3	3

1. Do a dot plot of the data for both groups on the same graph.
2. Compute the mean, median, and 10% trimmed mean for both groups.
3. Explain why the difference in means suggests one conclusion about the effect of the regimen, while the difference in medians or trimmed means suggests a different conclusion.

1.5 Measures of Spread

Range: Maximum minus minimum. Very sensitive to extreme values.

Sample Variance: Measures the average squared deviation from the mean.

$$\boxed{s^2 = \frac{1}{n-1}\sum_{i=1}^{n}(x_i - \bar{x})^2}$$

Standard Deviation: The square root of the sample variance.

$$\boxed{s = \sqrt{s^2}}$$

Degrees of Freedom: The number of independent pieces of information available to estimate variability. For sample variance: $df = n - 1$.

Exercise 1.6 — Silicone Rubber Tensile Strength

The tensile strength of silicone rubber is thought to be a function of curing temperature. Twelve specimens were prepared at 20°C and twelve at 45°C. Tensile strength values (megapascals):

20°C:	2.07	2.14	2.22	2.03	2.21	2.03
	2.05	2.18	2.09	2.14	2.11	2.02
45°C:	2.52	2.15	2.49	2.03	2.37	2.05
	1.99	2.42	2.08	2.42	2.29	2.01

1. Show a dot plot of the data with both low and high temperature tensile strength values.
2. Compute sample mean tensile strength for both samples.
3. Does it appear as if curing temperature has an influence on tensile strength, based on the plot? Comment further.
4. Does anything else appear to be influenced by an increase in curing temperature? Explain.

Exercise 1.14 — Tire Pressure

The following are the measurements of the tire pressure of eight tires (in psi):

572, 572, 573, 568, 569, 575, 565, 570.

1. Find the sample mean and median.
2. Find the sample variance, standard deviation, and range.
3. Using the calculated statistics in parts (a) and (b), can you comment on the quality of the tires?

Exercise 1.16 — Deviations Sum to Zero

Show that the n pieces of information in $\sum_{i=1}^{n}(x_i - \bar{x})^2$ are not independent; that is, show that

$$\sum_{i=1}^{n}(x_i - \bar{x}) = 0.$$

1.6 Graphical Representations of Data

Scatter Plot: Used to display the relationship between two quantitative variables (x, y). A scatter plot helps identify trends, patterns, and associations between variables.

Stem-and-Leaf Plot: An intermediate step between raw data and a frequency table. Preserves the original data values while showing the distribution.

Stem	Leaf
1	2 4 7
2	1 3 5 8
3	0 4 6

Relative Frequency Table: Shows the proportion of observations in each class.

Class Interval	Class Midpoint	Frequency	Relative Frequency
10–19	14.5	3	0.30
20–29	24.5	4	0.40
30–39	34.5	3	0.30

Histogram: A graphical representation of a frequency or relative frequency table using contiguous bars.

When describing the shape of a histogram, we commonly classify it as:

- **Symmetric**
- **Skewed right** (positively skewed)
- **Skewed left** (negatively skewed)

Symmetric distribution example



Exercise 1.15 — Fair Coin?

Eight independent coin tosses result in $HHHHHHHH$. It turns out that if the coin is fair, the probability of this outcome is $(1/2)^8 = 0.00390625$. Does this produce strong evidence that the coin is not fair? Comment and use the concept of P-value discussed in Section 1.1.

Exercise 1.17 — Smoking and Sleep

A study of the effects of smoking on sleep patterns is conducted. The measure observed is the time, in minutes, that it takes to fall asleep:

Smokers:	69.3	56.0	22.1	47.6
	53.2	48.1	52.7	34.4
	60.2	43.8	23.2	13.8
Nonsmokers:	28.6	25.1	26.4	34.9
	29.8	28.4	38.5	30.2
	30.6	31.8	41.6	21.1
	36.0	37.9	13.9	

1. Find the sample mean for each group.
2. Find the sample standard deviation for each group.
3. Make a dot plot of the data sets A and B on the same line.
4. Comment on what kind of impact smoking appears to have on the time required to fall asleep.

Exercise 1.18 — Exam Grades: Stem-and-Leaf and Histogram

The following scores represent the final examination grades for an elementary statistics course:

23	60	79	32	57	74	52	70	82
36	80	77	81	95	41	65	92	85
55	76	52	10	64	75	78	25	80
98	81	67	41	71	83	54	64	72
88	62	74	43	60	78	89	76	84
48	84	90	15	79	34	67	17	82
69	74	63	80	85	61			

1. Construct a stem-and-leaf plot for the examination grades in which the stems are 1, 2, 3, . . . , 9.
2. Construct a relative frequency histogram, draw an estimate of the graph of the distribution, and discuss the skewness of the distribution.
3. Compute the sample mean, sample median, and sample standard deviation.

Exercise 1.19 — Fuel Pump Lifetimes

The following data represent the length of life in years, measured to the nearest tenth, of 30 similar fuel pumps:

2.0	3.0	0.3	3.3	1.3	0.4
0.2	6.0	5.5	6.5	0.2	2.3
1.5	4.0	5.9	1.8	4.7	0.7
4.5	0.3	1.5	0.5	2.5	5.0
1.0	6.0	5.6	6.0	1.2	0.2

1. Construct a stem-and-leaf plot for the life in years of the fuel pumps, using the digit to the left

of the decimal point as the stem for each observation.

2. Set up a relative frequency distribution.

3. Compute the sample mean, sample range, and sample standard deviation.

Exercise 1.20 — Fruit Fly Lifetimes

The following data represent the length of life, in seconds, of 50 fruit flies subject to a new spray in a controlled laboratory experiment:

17	20	10	9	23	13	12	19	18	24
12	14	6	9	13	6	7	10	13	7
16	18	8	13	3	32	9	7	10	11
13	7	18	7	10	4	27	19	16	8
7	10	5	14	15	10	9	6	7	15

1. Construct a frequency distribution.

2. Construct a relative frequency histogram.

3. Compute the sample mean, sample median, and sample standard deviation.

Chapter 2

Chapter 2: Probability

2.1 Experiments, Sample Spaces, and Events

Experiment: A process that generates an outcome.

Sample Space (S): The set of all possible outcomes of an experiment.

Example 1:

Select 3 items from a production line. Each item can be classified as either defective (D) or non-defective (N).

$$S = \{DDD, DDN, DND, NDD, DNN, NDN, NND, NNN\}$$

Since each item has 2 possible outcomes,

$$|S| = 2^3 = 8$$

Example 2:

$$S = \{(x, y) \mid x^2 + y^2 \leq 4\}$$

Event (A): A subset of the sample space S.

Examples of events:

$$A = \{DDD, DDN, DND, NDD\}$$

$$B = \{NNN\}$$

$$C = \{(x, y) \mid x^2 + y^2 \leq 4\}$$

2.2 Event Operations and Probability Rules

Event Operations:

- Complement: A^c (or A')
- Intersection: $A \cap B$
- Union: $A \cup B$
- Null Event: $\varnothing$

If

$$A \cap B = \varnothing,$$

then A and B are mutually exclusive.

Example (Venn Diagram):



$$A = \{DDD, DDN, DND, NDD\}, \quad B = \{NNN\}$$

$$A \cup B = \{DDD, DDN, DND, NDD, NNN\}$$

$$A \cap B = \varnothing$$

1. *Experiment: A process that generates an outcome.*
2. *Sample Space (S): The set of all possible outcomes of an experiment.*
3. *Event Operations:*
   - *Complement: A' (A^c)*
   - *Intersection: $A \cap B$*
   - *Union: $A \cup B$*

- *Null Event:* $\varnothing$

4. *If $A \cap B = \varnothing$, then A and B are called* <u>*mutually exclusive*</u>.

$$(A \cap B)' = A' \cup B'$$

$$(A \cup B)' = A' \cap B'$$

$$A \cap \varnothing = \varnothing$$

$$A \cup \varnothing = A$$

$$A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$$

Exercise 2.5

Problem: Toss a die, then flip a coin once if even, twice if odd. Construct a tree diagram for the 18-element sample space.
Solution: Even outcomes (2,4,6): each followed by H or T $\Rightarrow$ 6 elements. Odd outcomes (1,3,5): each followed by HH, HT, TH, TT $\Rightarrow$ 12 elements. Total: $6 + 12 = 18$ elements. E.g. $\{1HH, 1HT, 1TH, 1TT, 2H, 2T, 3HH, \ldots\}$.

Exercise 2.9

Problem: For the sample space of Exercise 2.5, list elements for events A (die < 3), B (two tails), A', $A' \cap B$, $A \cup B$.
Solution: $A = \{1HH, 1HT, 1TH, 1TT, 2H, 2T\}$, $B = \{1TT, 3TT, 5TT\}$, $A' = S \setminus A$ (12 elements with die ≥ 3), $A' \cap B = \{3TT, 5TT\}$, $A \cup B = A \cup \{3TT, 5TT\}$ (8 elements).

2.3 Probability

$P(A) =$ <u>probability</u> of event A: the proportion of times the event occurs in infinitely many repetitions of the experiment.

Theorem 2.1

$$0 \leq P(A) \leq 1$$

$$P(A) + P(A') = 1$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$\begin{aligned} P(A \cup B \cup C) = P(A) &+ P(B) + P(C) \\ &- P(A \cap B) - P(A \cap C) - P(B \cap C) \\ &+ P(A \cap B \cap C) \end{aligned}$$

Exercise 2.14

Problem: $S = \{0, \ldots, 9\}$, $A = \{2, 4, 6, 8\}$, $B = \{1, 3, 5, 7, 9\}$, $C = \{0, 2, 3, 4, 5\}$, $D = \{1, 6, 7\}$. Find various set operations.
Solution: (1) $A \cup D = \{1, 2, 4, 6, 7, 8\}$. (2) $A \cap B = \emptyset$. (3) $B' = \{0, 2, 4, 6, 8\}$. (4) $(B' \cap D) \cup A = (\{6\}) \cup \{2, 4, 6, 8\} = \{2, 4, 6, 8\}$. (5) $(S \cap D)' = D' = \{0, 2, 3, 4, 5, 8, 9\}$. (6) $A \cap C \cap D' = \{2, 4, 6, 8\} \cap \{0, 2, 3, 4, 5\} \cap \{0, 2, 3, 4, 5, 8, 9\} = \{2, 4\}$.

Exercise 2.15

Problem: S = {copper, sodium, nitrogen, potassium, uranium, oxygen, zinc}, A = {copper, sodium, zinc}, B = {sodium, nitrogen, potassium}, C = {oxygen}.
Solution: (1) A' = {nitrogen, potassium, uranium, oxygen}. (2) $B \cup A$ = {copper, sodium, zinc, nitrogen, potassium}. (3) $(A \cup B') \cap C$: B' = {copper, uranium, oxygen, zinc}, $A \cup B'$ = $S \setminus$ {nitrogen, potassium}, so $(A \cup B') \cap C$ = {oxygen}. (4) $A' \cup C$ = {nitrogen, potassium, uranium, oxygen}. (5) $A \cup B \cup C = S \setminus$ {uranium}. (6) $(A' \cap B') \cup (A' \cup C) = A' \cup C$ = {nitrogen, potassium, uranium, oxygen}.

Exercise 2.17

Problem: Shade Venn diagram regions for $(A \cap B)'$, $(A \cup B)'$, $(A \cap C) \cup B$.
Solution: (1) $(A \cap B)'$: everything outside the overlap of A and B. (2) $(A \cup B)'$: everything outside both A and B. (3) $(A \cap C) \cup B$: the overlap of A and C, combined with all of B.

2.4 Mutually Exclusive Events

<u>Definition</u>: If $A_1, A_2, \ldots, A_n$ are mutually exclusive, then

$$P(A_1 \cup A_2 \cup \cdots \cup A_n) = P(A_1) + P(A_2) + \cdots + P(A_n)$$

If

$$A_1 \cup A_2 \cup \cdots \cup A_n = S,$$

then $\{A_1, A_2, \ldots, A_n\}$ is a <u>partition</u> of S.



Figure 2.1: Partition of the sample space S into A_1, A_2, A_3

2.4.1 Example

In a class of 33 students:

- 17 earned an A on the midterm
- 14 earned an A on the final
- 11 earned no A on either exam

Find the probability that a randomly selected student earned A's on <u>both</u> exams.



$$|A \cup B| = 33 - 11 = 22$$

$$\begin{aligned} |A \cap B| &= |A| + |B| - |A \cup B| \\ &= 17 + 14 - 22 = 9 \end{aligned}$$

$$|A \setminus B| = |A| - |A \cap B| = 17 - 9 = 8$$

$$|B \setminus A| = |B| - |A \cap B| = 14 - 9 = 5$$

$$P(A \cap B) = \frac{|A \cap B|}{|S|} = \frac{9}{33} \approx 0.273$$

Figure 2.2: Events A: A on midterm, B: A on final, with region counts and calculations

2.5 Counting Techniques and Equally Likely Outcomes

Theorem 2.2 (Equally Likely Outcomes)

If the sample space S has a finite number of outcomes and all outcomes are equally likely, then for any event A,

$$P(A) = \frac{|A|}{|S|}.$$

where

$|A|$ = number of outcomes in event A, $\quad |S|$ = number of outcomes in the sample space.

2.5.1 Example 1: Poker Hands Basics

A standard deck has:

$$4 \text{ suits} \times 13 \text{ denominations (A,2,3,...,Q,K)} = 52 \text{ cards.}$$

A poker hand consists of 5 cards chosen from 52:

$$|S| = \binom{52}{5} = 2{,}598{,}960.$$

2.5.2 Combinations Reminder

If there are 3 objects $\{A, B, C\}$ and we choose 2:

$$\binom{3}{2} = \frac{3!}{(3-2)!2!}.$$

Order does <u>not</u> matter.

2.5.3 Example 2: Probability of 2 Aces and 1 Jack

A 5-card hand contains:

- exactly 2 aces out of 4 ->
$$\binom{4}{2}$$
,
- exactly 1 jack ->
$$\binom{4}{1}$$
,

- 2 cards that are neither aces nor jacks,
$$\binom{52-8}{2}$$
.

$$P(\text{2 aces and 1 jack}) = \frac{\binom{4}{2}\binom{4}{1}\binom{44}{2}}{\binom{52}{5}}.$$

2.5.4 Example 3: Probability of a Full House

A full house consists of:

- 3 cards of one denomination
- 2 cards of a different denomination

Number of full house hands:
$$\binom{13}{1}\binom{4}{3}\binom{12}{1}\binom{4}{2}.$$

Thus,
$$P(\text{full house}) = \frac{\binom{13}{1}\binom{4}{3}\binom{12}{1}\binom{4}{2}}{\binom{52}{5}}.$$

2.5.5 Example 4: Probability of Four of a Kind

A four of a kind consists of:

- 4 cards of the same denomination
- 1 remaining card of a different denomination

Number of such hands:
$$\binom{13}{1}\binom{4}{4}\binom{48}{1}.$$

Thus,
$$P(\text{four of a kind}) = \frac{\binom{13}{1}\binom{4}{4}\binom{48}{1}}{\binom{52}{5}}.$$

2.5.6 Example 5: Probability of Exactly One Pair

An **excatly** one-pair hand consists of:

- 1 pair
- 3 cards of different denominations, none matching the pair

Number of such hands:

$$\binom{13}{1}\binom{4}{2}\binom{12}{3}\binom{4}{1}^3.$$

Thus,

$$P(\text{exactly one pair}) = \frac{\binom{13}{1}\binom{4}{2}\binom{12}{3}\binom{4}{1}^3}{\binom{52}{5}}.$$

2.5.7 Note*: Counting Patterns

$$\binom{a}{b}$$

Meaning: Choose b different items from a **at once**, order does not matter.

Key features:

- No repeats
- Grouped choice
- Used when items must be <u>distinct</u>

$$\binom{a}{1}^b$$

Meaning: Make b <u>independent choices</u>, each time choosing 1 item from c.

Key features:

- Repeats allowed
- Choices are independent
- Used when selections do <u>not</u> restrict each other

Rule to Remember:

$$\text{Different items, no repeats} \Rightarrow \binom{a}{b}$$

$$\text{Independent choices} \Rightarrow \binom{c}{1}^b$$

1. *<u>Probability</u> is the proportion of times the event occurs in infinitely many repetitions of the experiment.*
2. $0 \leq P(A) \leq 1$
3. $P(A) + P(A^c) = 1$
4. $P(A \cup B) = P(A) + P(B) - P(A \cap B)$

5. $$\begin{aligned} P(A \cup B \cup C) = {} & P(A) + P(B) + P(C) \\ & - P(A \cap B) - P(A \cap C) - P(B \cap C) \\ & + P(A \cap B \cap C) \end{aligned}$$

6. <u>*Permutation*</u>*: A permutation counts ordered arrangements.*

$$\boxed{{}_nP_r = \frac{n!}{(n-r)!}}$$

2.5.8 Example 1: Two fair dice

A pair of fair dice are rolled. Find the probability that the second die lands on a smaller value than the first. The outcomes where the second die is smaller than the first are represented below.

First Die (Stem)	Second Die (Leaf)
2	1
3	1 2
4	1 2 3
5	1 2 3 4
6	1 2 3 4 5

There are 15 favorable outcomes and 36 total outcomes.

$$P(\text{second} < \text{first}) = \frac{15}{36} = \frac{5}{12}.$$

Exercise 2.55

Problem: Coded catalog items begin with 2 distinct letters followed by 3 distinct nonzero digits. Find P(first letter vowel, second non-vowel, last digit even).

Solution:

Total codes:

$$P(26, 2) \times P(9, 3) = (26)(25)(9)(8)(7) = 327{,}600$$

Favorable codes:

- First letter is a vowel: $P(5, 1) = 5$
- Second letter is a non-vowel (21 consonants available): $P(21, 1) = 21$
- Last digit must be even, pick from $\{2, 4, 6, 8\}$: $P(4, 1) = 4$
- Remaining 2 digit positions from the remaining 8 digits: $P(8, 2) = 56$

$$\text{Favorable} = P(5, 1) \times P(21, 1) \times P(4, 1) \times P(8, 2) = (5)(21)(4)(56) = 23{,}520$$

Therefore,

$$P = \frac{23{,}520}{327{,}600} = \frac{28}{390} = \frac{14}{195} \approx 0.0718$$

Exercise 2.107

Problem: How many bridge hands contain 4 spades, 6 diamonds, 1 club, 2 hearts?
Solution: $\binom{13}{4}\binom{13}{6}\binom{13}{1}\binom{13}{2} = 715 \times 1716 \times 13 \times 78 = \boxed{1,244,117,080}$.

2.6 Conditional Probability and Independence

2.6.1 Conditional Probability

The conditional probability of an event B given that event A has occurred is the probability that B occurs when it is known that A has occurred.

$$\boxed{P(B \mid A) = \frac{P(A \cap B)}{P(A)}, \quad P(A) > 0}$$

2.6.2 Example 2: Drinking Survey

A survey records the following data:

	D	N	Total
M	19	41	60
F	12	28	40
Total	31	69	100

The symbols used above are defined as follows:

- M: male
- F: female
- D: the individual drinks
- N: the individual does not drink

$$P(D|M) = \frac{19}{60} \qquad P(M|D) = \frac{19}{31}$$

Exercise 2.85

Problem: $P(\text{correct diagnosis}) = 0.7$. Given incorrect diagnosis, $P(\text{lawsuit}) = 0.9$. Find $P(\text{incorrect diagnosis} \cap \text{lawsuit})$.
Solution: $P(\text{incorrect}) = 1 - 0.7 = 0.3$. $P(\text{incorrect} \cap \text{lawsuit}) = P(\text{lawsuit} \mid \text{incorrect}) \times P(\text{incorrect}) = 0.9 \times 0.3 = \boxed{0.27}$.

Exercise 2.86

Problem: In 1970, 11% of Americans completed four years of college; 43% of them were women. In 1990, 22% of Americans completed four years of college; 53% of them were women.

(1) $P(\text{woman} \mid \text{college in 1970})$:

Directly stated: 43% of college completers were women.

$$P(\text{woman} \mid \text{college in 1970}) = 0.43$$

(2) $P(\text{woman} \cap \text{college in 1990})$:

Using the multiplication rule, $P(A \cap B) = P(A \mid B) \cdot P(B)$:

$$P(\text{woman} \cap \text{college}) = P(\text{woman} \mid \text{college}) \times P(\text{college}) = 0.53 \times 0.22 = 0.1166$$

(3) $P(\text{man had not finished college in 1990})$:

First, find the probability a man finished college:

$$P(\text{man} \mid \text{college}) = 1 - 0.53 = 0.47$$

$$P(\text{man} \cap \text{college}) = 0.47 \times 0.22 = 0.1034$$

Then use the complement:

$$P(\text{man not college}) = 1 - P(\text{man} \cap \text{college}) = 1 - 0.1034 = \boxed{0.8966}$$

2.7 Law of Total Probability

Theorem 2.3 (Law of Total Probability)

If $B_1, B_2, \ldots, B_k$ form a partition of the sample space S with $P(B_i) > 0$ for all i, then for any event A,

$$P(A) = \sum_{i=1}^{k} P(A \mid B_i)\, P(B_i).$$

2.7.1 Example 3: Monty Hall (3 doors)

Car location	Monty opens	Probability	Stay	Switch
Door 1	Door 2	$\frac{1}{6}$	Car	Goat
Door 1	Door 3	$\frac{1}{6}$	Car	Goat
Door 2	Door 3	$\frac{1}{3}$	Goat	Car
Door 3	Door 2	$\frac{1}{3}$	Goat	Car

Staying wins only when the car is behind Door 1, so

$$P(\text{win by staying}) = \frac{1}{6} + \frac{1}{6} = \frac{1}{3}.$$

Switching wins when the car is behind Door 2 or Door 3, so

$$P(\text{win by switching}) = \frac{1}{3} + \frac{1}{3} = \frac{2}{3}.$$

2.7.2 Example 4: Birthday Problem

Assume the following:

- Leap years are ignored
- All 365 birthdays are equally likely
- Birthdays of different people are independent

Question: What is the probability that at least two people share the same birthday in a group of n people?

Rather than computing this directly, we use the complement rule.

$$P(\text{at least one match}) = 1 - P(\text{no match})$$

2.7.3 Probability of no shared birthdays

- Person 1 can have any birthday: probability 1
- Person 2 must avoid that birthday: $\frac{364}{365}$
- Person 3 must avoid the first two birthdays: $\frac{363}{365}$
- ...
- Person n must avoid the previous $n-1$ birthdays: $\frac{365-(n-1)}{365}$

Therefore,

$$P(\text{no match}) = \frac{365}{365} \cdot \frac{364}{365} \cdot \frac{363}{365} \cdots \frac{365-(n-1)}{365}$$

or equivalently,

$$P(\text{no match}) = \prod_{k=0}^{n-1} \frac{365-k}{365}$$

2.7.4 Final result

$$P(\text{at least one shared birthday}) = 1 - \prod_{k=0}^{n-1} \frac{365-k}{365}$$

2.7.5 Important values

- For $n = 23$: $P(\text{at least one match}) \approx 0.507$
- For $n = 57$: $P(\text{at least one match}) \approx 0.99$

<u>*Conditional Probability:*</u>

The probability of event B given that event A has occurred is

$$\boxed{P(B \mid A) = \frac{P(A \cap B)}{P(A)}, \quad P(A) > 0}$$

Read as: the probability of B given A.

2.8 Independence of Events

Definition (Independence): Events A and B are <u>independent</u> if and only if

$$\boxed{P(B \mid A) = P(B)}$$

Equivalently,

$$P(A \mid B) = P(A)$$

or

$$\boxed{P(A \cap B) = P(A)\,P(B)}$$

2.8.1 Multiple Independent Events

<u>Definition:</u>

If events $A_1, A_2, \ldots, A_k$ are independent, then

$$\boxed{P(A_1 \cap A_2 \cap \cdots \cap A_k) = P(A_1)\,P(A_2)\cdots P(A_k)}$$

2.8.2 Mutual Independence

Mutual Independence : A collection of events $A_1, A_2, \ldots, A_n$ is mutually independent if and only if for *every* subcollection $\{A_{i_1}, \ldots, A_{i_k}\}$,

$$\boxed{P\left(\bigcap_{j=1}^{k} A_{i_j}\right) = \prod_{j=1}^{k} P(A_{i_j})}$$

Example (Three Events):

Events A_1, A_2, A_3 are mutually independent if all of the following hold:

$$P(A_1 \cap A_2) = P(A_1)P(A_2)$$

$$P(A_1 \cap A_3) = P(A_1)P(A_3)$$

$$P(A_2 \cap A_3) = P(A_2)P(A_3)$$

$$P(A_1 \cap A_2 \cap A_3) = P(A_1)P(A_2)P(A_3)$$

Note: Mutually exclusive events are dependent. If one event occurs, the other cannot occur.

2.8.3 Example: Component Reliability

An electrical system has four components A, B, C, D. The system works if A and B work and at least one of C or D works. Assume all components are independent.



$$P(A) = 0.9, \quad P(B) = 0.9, \quad P(C) = 0.8, \quad P(D) = 0.8$$

2.8.4 (a) Probability the entire system works

The system works if A and B work and either C or D works.

$$\begin{aligned} P(\text{system works}) &= P(\text{all work}) + P(A, B, C \text{ work}, D \text{ does not}) \\ &\quad + P(A, B, D \text{ work}, C \text{ does not}) \end{aligned}$$

$$= (0.9)(0.9)(0.8)(0.8) + (0.9)(0.9)(0.8)(1-0.8) + (0.9)(0.9)(0.8)(1-0.8)$$

$$\boxed{P(\text{system works}) = 0.7776}$$

2.8.5 (b) Conditional probability

$$P(C^c \mid \text{system works}) = \frac{P(C^c \cap \text{system works})}{P(\text{system works})}$$

$$P(C^c \cap \text{system works}) = (0.9)(0.9)(0.8)(1-0.8)$$

$$\boxed{P(C^c \mid \text{system works}) = \frac{(0.9)(0.9)(0.8)(1-0.8)}{0.7776} = 0.16}$$

Theorem of Total Probability

Let $B_1, B_2, \ldots, B_k$ be a partition of the sample space S such that $P(B_i) > 0$ for all i. Then for any event $A \subseteq S$,

$$\boxed{P(A) = \sum_{i=1}^{k} P(A \mid B_i)\, P(B_i) = \sum_{i=1}^{k} P(A \cap B_i)}$$

Theorem: Bayes' Rule (1701–1761)

Let $B_1, B_2, \ldots, B_k$ be a partition of the sample space S such that $P(B_i) > 0$ for $i = 1, \ldots, k$. For any event $A \subseteq S$ with $P(A) > 0$,

$$\boxed{P(B_r \mid A) = \frac{P(B_r \cap A)}{\sum_{i=1}^{k} P(B_i \cap A)} = \frac{P(B_r)\, P(A \mid B_r)}{\sum_{i=1}^{k} P(B_i)\, P(A \mid B_i)}, \quad r = 1, \ldots, k}$$



Figure 2.3: Partition of S into B_1, B_2, B_3 with shaded regions $A \cap B_i$

2.8.6 Example (Medical Test)

The fraction of people in a population who have a certain disease is 0.01.

$$P(D) = 0.01, \quad P(D^c) = 0.99$$

The test characteristics are:

$$P(\text{test says } D \mid D^c) = 0.05 \quad \text{(false positive rate)}$$

$$P(\text{test says } D^c \mid D) = 0.20 \quad \text{(false negative rate)}$$

Thus,

$$P(\text{test says } D \mid D) = 1 - 0.20 = 0.80$$

<u>Note:</u> $1 - P(\text{test says } D^c \mid D)$ is called the <u>sensitivity</u> of the test, and $1 - P(\text{test says } D \mid D^c)$ is called the <u>specificity</u>.



2.8.7 (a) Probability the test says disease

$$P(\text{test says } D) = P(D \cap \text{test says } D) + P(D^c \cap \text{test says } D)$$

$$= P(\text{test says } D \mid D)P(D) + P(\text{test says } D \mid D^c)P(D^c)$$

$$= (0.80)(0.01) + (0.05)(0.99) = 0.0575$$

2.8.8 (b) Probability of disease given positive test

$$P(D \mid \text{test says } D) = \frac{P(D \cap \text{test says } D)}{P(\text{test says } D)}$$

$$= \frac{P(\text{test says } D \mid D)P(D)}{0.0575} = \frac{(0.80)(0.01)}{0.0575}$$

$$\boxed{P(D \mid \text{test says } D) \approx 0.139}$$

2.8.9 (c) Probability of disease given negative test

$$P(D \mid \text{test says } D^c) = \frac{P(D \cap \text{test says } D^c)}{P(\text{test says } D^c)}$$

$$= \frac{P(\text{test says } D^c \mid D)P(D)}{1 - P(\text{test says } D)}$$

$$= \frac{(0.20)(0.01)}{1 - 0.0575}$$

$$\boxed{P(D \mid \text{test says } D^c) \approx 0.00212}$$

Exercise 2.110

Problem: $P(\text{recovery}) = 0.8$. Find (1) $P(\text{exactly 2 of 3 survive})$, (2) $P(\text{all 3 survive})$.
Solution: $X \sim \text{Binomial}(n = 3, p = 0.8)$.
(1) $P(X = 2) = \binom{3}{2}(0.8)^2(0.2)^1 = 3(0.64)(0.2) = \boxed{0.384}$.
(2) $P(X = 3) = (0.8)^3 = \boxed{0.512}$.

Exercise 2.127

Problem: There is a 50-50 chance that the queen carries the gene of hemophilia. If she is a carrier, then each prince has a 50-50 chance of having hemophilia independently. If the queen is not a carrier, the prince will not have the disease. Suppose the queen has had three princes without the disease. What is the probability the queen is a carrier?
Solution: Let C = carrier. $P(C) = 0.5$, $P(C') = 0.5$.
$P(3 \text{ healthy} \mid C) = (0.5)^3 = 0.125$. $P(3 \text{ healthy} \mid C') = 1$.
By Bayes' theorem:

$$P(C \mid 3 \text{ healthy}) = \frac{P(3 \text{ healthy} \mid C) \cdot P(C)}{P(3 \text{ healthy} \mid C) \cdot P(C) + P(3 \text{ healthy} \mid C') \cdot P(C')} = \frac{0.125 \times 0.5}{0.125 \times 0.5 + 1 \times 0.5} = \frac{0.0625}{0.5625} = \boxed{\frac{1}{9} \approx 0.1111}$$

Midterm Q1(a) – Independent & Mutually Exclusive Events (5 pts)

Problem: A, B independent; C mutually exclusive of both A and B. $P(A) = 0.1$, $P(B) = 0.3$, $P(C) = 0.5$. Find $P((A \cap B) \cup C)$.



Solution: $P(A \cap B) = P(A)P(B) = 0.03$ (independence). Since C is mutually exclusive of $A \cap B$: $(A \cap B) \cap C = \emptyset$. $P((A \cap B) \cup C) = 0.03 + 0.5 = \boxed{0.53}$.

Midterm Q1(b) – Conditional Probability (5 pts)

Problem: Find $P(A \cap B \mid C')$.
Solution: Since C is mutually exclusive of A and B, we have $A \cap B \subseteq C'$, so $(A \cap B) \cap C' = A \cap B$.

$$P(A \cap B \mid C') = \frac{P(A \cap B)}{P(C')} = \frac{0.03}{0.5} = \boxed{0.06}.$$

Chapter 3

Chapter 3: Random Variables and Probability Distributions

3.1 Random Variables and Their Interpretation

Definition: A random variable (r.v.) is a rule that assigns a **real number** to each outcome in the sample space.
Alternative definition: A random variable is a function that takes the outcome of an experiment and assigns it a number so that probabilities can be calculated.

3.1.1 Example 1: Three Electronic Components

Each component is classified as either defective (D) or non-defective (N).

$$S = \{NNN, DNN, NDN, NND, DDN, DND, NDD, DDD\}$$

- **Defective (D):** the component does not meet required specifications and fails inspection.
- **Non-defective (N):** the component meets specifications and passes inspection.

Define the random variable

$$X = \text{number of defective components.}$$

Then:

$$\begin{aligned} X &= 0 \quad \text{for } \{NNN\} \\ X &= 1 \quad \text{for } \{DNN, NDN, NND\} \\ X &= 2 \quad \text{for } \{DDN, DND, NDD\} \\ X &= 3 \quad \text{for } \{DDD\} \end{aligned}$$

Thus, the possible values of X are:

$$\{0, 1, 2, 3\}.$$

3.1.2 Example 2: One Component (Dummy Variable)

$$S = \{D, N\}$$

Define the random variable

$$X = \begin{cases} 1, & \text{if the component is defective (D)} \\ 0, & \text{if the component is non-defective (N)} \end{cases}$$

This is called a **dummy variable** because the outcome is categorical, but is encoded numerically.
A dummy variable is a special type of random variable that assigns numerical labels to categorical outcomes, where the numbers have no quantitative meaning beyond identification.

3.2 Discrete Random Variables

Definition: A random variable is called discrete if its set of possible values is **countable** (finite or countably infinite).

3.2.1 Example 3: Sampling Until First Defective

Components are tested one(independently) at a time until the first defective component is observed.

$$S = \{D,\ ND,\ NND,\ NNND, \ldots\}$$

Define

$$X = \text{number of components tested until the first defective.}$$

Then:

$$\begin{aligned} X &= 1 \quad \text{for } \{D\} \\ X &= 2 \quad \text{for } \{ND\} \\ X &= 3 \quad \text{for } \{NND\} \\ &\vdots \end{aligned}$$

Hence,

$$X = 1, 2, 3, \ldots$$

Since the possible values can be listed, X is a discrete random variable.

Non-discrete version of the same experiment:

Define

$$Y = \text{time (in seconds) until the first defective component is observed.}$$

Since Y can take any real value in $[0, \infty)$, it cannot be listed and is therefore a continuous (non-discrete) random variable.

3.2.2 Discrete vs. Continuous Random Variables

Discrete Random Variable	Continuous Random Variable
Counts things	Measures things
Possible values are countable	Possible values fill an interval
$P(X = x)$ can be > 0	$P(X = x) = 0$ for all x
Uses a probability mass function (PMF)	Uses a probability density function (PDF)

Exercise 3.3

Problem: W = number of heads minus tails in 3 coin tosses. List sample space and assign values of W.
Solution: $S = \{HHH, HHT, HTH, THH, HTT, THT, TTH, TTT\}$. W: HHH$\rightarrow$ 3, HHT/HTH/THH$\rightarrow$ 1, HTT/THT/TTH$\rightarrow$ -1, TTT$\rightarrow$ -3.

Exercise 3.8

Problem: Find the probability distribution of W from Ex. 3.3 if $P(H) = 2/3$.
Solution: $P(W = 3) = (2/3)^3 = 8/27$. $P(W = 1) = 3(2/3)^2(1/3) = 12/27$. $P(W = -1) = 3(2/3)(1/3)^2 = 6/27$. $P(W = -3) = (1/3)^3 = 1/27$. Check: $8 + 12 + 6 + 1 = 27$. ✓

3.3 Probability Mass Function (PMF)

Definition:

Let X be a discrete random variable. The <u>probability mass function (PMF)</u> of X, denoted $f(x)$, is defined by:

$$\boxed{\begin{array}{ll} 1) & f(x) \geq 0 \quad \text{for all } x \\ 2) & \displaystyle\sum_x f(x) = 1 \end{array}}$$

Note:

- Capital X: random variable
- Lowercase x: a specific value

Exercise 3.11

Problem: A shipment of 7 televisions contains 2 defective ones. A hotel makes a random purchase of 3 televisions. If x is the number of defective ones purchased by the hotel, find the probability distribution of X. Express the results graphically as a probability histogram.
Solution: $X \sim \text{Hypergeometric}(N = 7, K = 2, n = 3)$. $P(X = 0) = \frac{\binom{2}{0}\binom{5}{3}}{\binom{7}{3}} = \frac{10}{35} = \frac{2}{7}$. $P(X = 1) = \frac{\binom{2}{1}\binom{5}{2}}{\binom{7}{3}} = \frac{20}{35} = \frac{4}{7}$. $P(X = 2) = \frac{\binom{2}{2}\binom{5}{1}}{\binom{7}{3}} = \frac{5}{35} = \frac{1}{7}$.

3.4 Bernoulli and Binomial Random Variables

3.4.1 I. Bernoulli Random Variable (Single Trial)

A Bernoulli random variable: X models a single experiment with only two possible outcomes: success or failure.

$$X = \begin{cases} 1, & \text{success} \\ 0, & \text{failure} \end{cases}$$

If $p = P(X = 1)$, then the PMF is

x	0	1
$P(X = x)$	$1 - p$	p

Here, p is the probability of success (e.g. observing a defective component).

3.4.2 Binomial Random Variable (Multiple Bernoulli Trials)

The binomial random variable extends the Bernoulli case to multiple independent trials.

Definition:

A random variable X is called a binomial random variable if it represents the number of successes in n independent Bernoulli trials, each with success probability p.

$$X = \text{number of successes in } n \text{ trials}$$

In this case,

$$X \sim \text{Bin}(n, p)$$

and the probability mass function is

$$\boxed{P(X = x) = \binom{n}{x} p^x (1-p)^{n-x}, \quad x = 0, 1, \ldots, n.}$$

3.4.3 Conditions for a Binomial Model

A binomial model applies only if:

- each trial has exactly two outcomes (success or failure),
- the probability of success p is the same for every trial,
- the trials are independent,
- the number of trials n is fixed.

3.4.4 Example: Three Components Tested

Assume each component is defective with probability

$$p = 0.1, \quad n = 3.$$

where p = P(a single component is defective), n = number of trials

Let

$$X = \text{number of defective components.}$$

x	$P(X = x)$
0	$\binom{3}{0}(0.9)^3 = 0.729$
1	$\binom{3}{1}(0.1)(0.9)^2 = 0.243$
2	$\binom{3}{2}(0.1)^2(0.9) = 0.027$
3	$\binom{3}{3}(0.1)^3 = 0.001$

$$0.729 + 0.243 + 0.027 + 0.001 = 1.$$



3.4.5 Geometric Random Variable

3.4.6 Example: Sampling Until First Defective

Components are sampled one at a time until the first defective component is observed. Assume the probability that a component is defective is

$$p = 0.1.$$

Define the random variable

$$X = \text{number of samples collected until the first defective.}$$

x	$P(X = x) = f(x)$
1	0.1
2	$0.9(0.1)$
3	$0.9^2(0.1)$
$\vdots$	$\vdots$

3.4.7 Geometric Random Variable

Definition:

A random variable X is called a geometric random variable if it represents the number of trials needed to obtain the first success in a sequence of independent Bernoulli trials with success probability p.

$$\boxed{P(X = x) = (1-p)^{x-1}p, \quad x = 1, 2, 3, \ldots}$$

In this example,

$$P(X = x) = 0.9^{x-1}(0.1).$$

3.4.8 Verification That Probabilities Sum to 1

$$\sum_{x=1}^{\infty} 0.9^{x-1}(0.1) = 0.1 \sum_{x=0}^{\infty} 0.9^x = 0.1 \left(\frac{1}{1 - 0.9} \right) = 1.$$

3.5 Cumulative Distribution Function (CDF)

Definition:

The cumulative distribution function (CDF) of a discrete random variable X with PMF $f(x)$ is defined as

$$\boxed{F(x) = P(X \leq x) = \sum_{t \leq x} f(t), \quad -\infty < x < \infty.}$$

3.5.1 Example: Binomial Distribution (Three Components)

Let X be the number of defective components when three components are tested, with

$$P(X = 0) = 0.729, \quad P(X = 1) = 0.243, \quad P(X = 2) = 0.027, \quad P(X = 3) = 0.001.$$

$$\begin{aligned}
F(0) &= P(X \leq 0) = 0.729 \\
F(1) &= P(X \leq 1) = 0.729 + 0.243 = 0.972 \\
F(2) &= P(X \leq 2) = 0.729 + 0.243 + 0.027 = 0.999 \\
F(3) &= P(X \leq 3) = 0.729 + 0.243 + 0.027 + 0.001 = 1
\end{aligned}$$

Thus, the CDF can be written as

$$F(x) = \begin{cases} 0, & x < 0 \\ 0.729, & 0 \leq x < 1 \\ 0.972, & 1 \leq x < 2 \\ 0.999, & 2 \leq x < 3 \\ 1, & x \geq 3 \end{cases}$$

3.5.2 Properties of the CDF

- $F(x)$ is <u>monotone non-decreasing</u>.
- If $x < y$, then $F(x) \leq F(y)$.
- $0 \leq F(x) \leq 1$.

Note: A function is **monotone** non-decreasing if its value never decreases as the input increases.

3.5.3 Using the CDF to Compute Probabilities

For $a < b$,

$$P(a < X \leq b) = F(b) - F(a).$$

Example:

$$P(0 < X \leq 2) = F(2) - F(0) = 0.999 - 0.729 = 0.27.$$

3.5.4 CDF Histogram (Step Function)



Note: For a discrete random variable, the <u>PMF</u> is drawn as a bar chart since it shows probabilities at individual points, while the <u>CDF</u> is drawn as a step function since it represents cumulative probability and is monotone non-decreasing.

1. <u>*Random Variable (RV), X*</u>: A random variable assigns a real number to each outcome.
2. <u>*Discrete Random Variable*</u>: If X is discrete,

- $P(X = x) = f(x)$
- $f(x)$ is the *probability mass function (PMF)*
- $f(x) \geq 0$
- $\sum_x f(x) = 1$

3. *Cumulative Distribution Function (CDF), $F(x)$*:
   - $F(x) = P(X \leq x)$
   - If X is discrete:

$$F(x) = \sum_{t \leq x} f(t), \quad -\infty < x < \infty$$

Exercise 3.15

Problem: Find CDF of X from Ex. 3.11. Use it to find $P(X = 1)$ and $P(0 < X \leq 2)$.

Solution: $F(x) = \begin{cases} 0, & x < 0 \\ 2/7, & 0 \leq x < 1 \\ 6/7, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$

$P(X = 1) = F(1) - F(0) = 6/7 - 2/7 = \boxed{4/7}$. $P(0 < X \leq 2) = F(2) - F(0) = 1 - 2/7 = \boxed{5/7}$.

3.6 Continuous Sample Space and Continuous Random Variables

If the sample space contains an infinite number of outcomes equal to the number of points on a line segment, it is called a continuous sample space.

A continuous random variable has

$$\boxed{P(X = x) = 0 \quad \text{for all } x,}$$

so probabilities are computed over intervals instead of single values.

***Alternative definition**: A continuous sample space contains infinitely many outcomes, like the points on a line segment. For a continuous random variable, the probability of taking any exact value is zero, i.e. $P(X = x) = 0$ for all x. Therefore, probabilities are computed over intervals rather than at single points.

3.7 Probability Density Function (PDF)

A function $f(x)$ is called a probability density function (PDF) of a continuous random variable X, defined over $\mathbb{R}$, if:

1. $f(x) \geq 0$ for all $x \in \mathbb{R}$
2. $\displaystyle\int_{-\infty}^{\infty} f(x)\,dx = 1$
3. For any $a < b$,
$$P(a < X < b) = \int_a^b f(x)\,dx$$

For continuous random variables,

$$P(a < X < b) = P(a \leq X \leq b) = P(a \leq X < b) = P(a < X \leq b).$$

$$\boxed{P(X = a) = 0}$$

because a single point has zero area under the probability density function.



$$\int_a^b f(x)\,dx = P(a < X < b)$$

3.7.1 Example: Uniform Distribution

Let the probability density function be

$$f(x) = \begin{cases} c, & 5 < x < 10, \\ 0, & \text{otherwise.} \end{cases}$$

1) Determine the value of c

Since $f(x)$ is a <u>probability density function</u>, the total area under the curve must equal 1:

$$\int_{-\infty}^{\infty} f(x)\,dx = 1.$$

Because $f(x) = 0$ outside the interval $(5, 10)$,

$$\int_5^{10} c\,dx = 1.$$

Evaluating the integral,

$$c(10-5)=1 \quad \Rightarrow \quad c=\frac{1}{5}.$$

Thus,

$$f(x)=\begin{cases}\frac{1}{5}, & 5<x<10,\\ 0, & \text{otherwise.}\end{cases}$$



2): Compute $P(X<7)$

Formula used:

$$P(a<X<b)=\int_a^b f(x)\,dx.$$

Applying this formula,

$$P(X<7)=\int_5^7 \frac{1}{5}\,dx=\frac{1}{5}(7-5)=\frac{2}{5}.$$

Exercise 3.36

Problem: $f(x)=12x(1-x)^2$ for $0<x<1$. Find (1) $P(X<0.5)$, (2) $P(X>0.4)$, (3) $P(X<0.7 \mid X\geq 0.5)$.

Solution: (1) $P(X<0.5)=\int_0^{0.5} 12x(1-x)^2\,dx$. Expand: $12x(1-2x+x^2)=12x-24x^2+12x^3$. $=[6x^2-8x^3+3x^4]_0^{0.5}=6(0.25)-8(0.125)+3(0.0625)=1.5-1+0.1875=\boxed{0.6875}$.

(2) $P(X>0.4)=1-P(X\leq 0.4)=1-[6(0.16)-8(0.064)+3(0.0256)]=1-[0.96-0.512+0.0768]=1-0.5248=\boxed{0.4752}$.

(3) $P(X<0.7 \mid X\geq 0.5)=\frac{P(0.5\leq X<0.7)}{P(X\geq 0.5)}=\frac{F(0.7)-F(0.5)}{1-F(0.5)}$. $F(0.7)=6(0.49)-8(0.343)+3(0.2401)=2.94-2.744+0.7203=0.9163$. $\frac{0.9163-0.6875}{1-0.6875}=\frac{0.2288}{0.3125}=\boxed{0.7322}$.

Exercise 3.73

Problem: Impurities Y has density $f(y)=10(1-y)^9$ for $0<y<1$. (1) $P(Y>0.5)$. (2) $P(0.25<Y<0.5)$.

Solution: CDF: $F(y)=1-(1-y)^{10}$.

(1) $P(Y>0.5)=(1-0.5)^{10}=(0.5)^{10}=\boxed{1/1024\approx 0.000977}$.

(2) $P(0.25<Y<0.5)=F(0.5)-F(0.25)=[1-(0.5)^{10}]-[1-(0.75)^{10}]=(0.75)^{10}-(0.5)^{10}\approx 0.05631-0.000977=\boxed{0.0553}$.

3.8 CDF for Continuous Random Variables

Def: Let X be a continuous random variable with pdf: probability density function $f(x)$. The cumulative distribution functio is

$$F(x) = P(X \leq x) = \int_{-\infty}^{x} f(t)\, dt.$$

Consequently:

$$\boxed{f(x) = \frac{d}{dx}F(x), \qquad P(a < X < b) = F(b) - F(a).}$$

3.8.1 Example

Let X be the time until a chemical reaction is complete (in msec). Suppose the CDF is

$$F(x) = \begin{cases} 0, & x < 0, \\ 1 - e^{-0.01x}, & x \geq 0. \end{cases}$$

(a) Find the pdf.

Use $f(x) = \dfrac{d}{dx}F(x)$.
For $x < 0$, $F(x) = 0$, so

$$f(x) = 0.$$

For $x \geq 0$,

$$f(x) = \frac{d}{dx}\left(1 - e^{-0.01x}\right) = 0.01e^{-0.01x}.$$

Therefore,

$$f(x) = \begin{cases} 0, & x < 0, \\ 0.01e^{-0.01x}, & x \geq 0. \end{cases}$$

(b) Find $P(X < 200)$.

Use the CDF directly:

$$P(X < 200) = F(200) = 1 - e^{-0.01(200)} = 1 - e^{-2} \approx 0.8647.$$

(c) Check if this is a valid CDF.

Theorem: Properties of a Cumulative Distribution Function

A function $F(x)$ is a valid cumulative distribution function (CDF) if and only if:

- $0 \leq F(x) \leq 1$ for all x
- $F(x)$ is monotone non-decreasing, i.e.

$$x \leq y \implies F(x) \leq F(y)$$

- $\lim_{x \to -\infty} F(x) = 0$
- $\lim_{x \to \infty} F(x) = 1$

For this $F(x)$:

$$\lim_{x \to -\infty} F(x) = 0, \qquad \lim_{x \to \infty} F(x) = 1,$$

and for $x \geq 0$, $1 - e^{-0.01x}$ increases as x increases, so $F(x)$ is monotone non-decreasing.

Thus, $F(x)$ is a valid CDF.

3.9 Joint Probability Distributions (Discrete)

Def: The function $f(x, y)$ is a joint probability mass function (PMF) of discrete random variables X and Y if:

1. $f(x, y) \geq 0$ for all (x, y)
2. $\sum_x \sum_y f(x, y) = 1$
3. $P(X = x, Y = y) = f(x, y)$

That is, $f(x, y)$ gives the probability that the two random variables X and Y take the values x and y *simultaneously*.

For any region A in the xy-plane,

$$\boxed{P\big((X, Y) \in A\big) = \sum_{(x,y) \in A} f(x, y)}$$

3.9.1 Example: Pen Refills

Two refills are selected at random and without replacement from a box containing:

- 3 blue refills
- 2 red refills
- 3 green refills

Define the random variables

$$X = \text{number of blue refills selected}, \qquad Y = \text{number of red refills selected}.$$

The total number of possible selections is

$$\binom{8}{2}.$$

3.9.2 Joint PMF Table

For each pair (x, y),

$$f(x, y) = \frac{\text{number of favorable outcomes}}{\binom{8}{2}}.$$

$X \backslash Y$	0	1	2	Row Total
0	$\frac{\binom{3}{2}}{\binom{8}{2}}$	$\frac{\binom{2}{1}\binom{3}{1}}{\binom{8}{2}}$	$\frac{\binom{2}{2}}{\binom{8}{2}}$	$\frac{\binom{5}{2}}{\binom{8}{2}}$
1	$\frac{\binom{3}{1}\binom{3}{1}}{\binom{8}{2}}$	$\frac{\binom{3}{1}\binom{2}{1}}{\binom{8}{2}}$	0	$\frac{15}{\binom{8}{2}}$
2	$\frac{\binom{3}{2}}{\binom{8}{2}}$	0	0	$\frac{3}{\binom{8}{2}}$
Column Total	$\frac{15}{\binom{8}{2}}$	$\frac{12}{\binom{8}{2}}$	$\frac{1}{\binom{8}{2}}$	1

3.9.3 Marginal Distributions

Def: Let $f(x, y)$ be the joint PMF of X and Y.

The <u>marginal PMF of X</u> is obtained by summing over all values of Y:

$$g(x) = \sum_y f(x, y).$$

The <u>marginal PMF of Y</u> is obtained by summing over all values of X:

$$h(y) = \sum_x f(x, y).$$

Marginal example:

$$P(X = 1) = \sum_y P(X = 1, Y = y).$$

Exercise 3.37

Problem: Find c so the following are valid joint PMFs: (1) $f(x,y) = cxy$ for $x = 1,2,3$; $y = 1,2,3$. (2) $f(x,y) = c|x-y|$ for $x = -1,0,1$; $y = -3,3$.

Solution: (1) $\sum\sum cxy = c(1+2+3)(1+2+3) = 36c = 1 \Rightarrow \boxed{c = 1/36}$.

(2) Enumerate $|x-y|$ values and sum: $|-1-(-3)| = 2$, $|-1-3| = 4$, $|0-(-3)| = 3$, $|0-3| = 3$, $|1-(-3)| = 4$, $|1-3| = 2$. Sum $= 18$. $18c = 1 \Rightarrow \boxed{c = 1/18}$.

Exercise 3.38

Problem: $f(x,y) = \frac{x+y}{48}$ for $x = 0,1,2,3$; $y = 0,1,2,3$. Find (1) $P(X \le 1, Y = 1)$, (2) $P(X > 1, Y \le 1)$, (3) $P(X \le Y)$, (4) $P(X+Y = 2)$.

Solution: (1) $P(X \le 1, Y = 1) = f(0,1) + f(1,1) = \frac{1}{48} + \frac{2}{48} = \boxed{3/48 = 1/16}$.

(2) $P(X > 1, Y \le 1) = f(2,0) + f(2,1) + f(3,0) + f(3,1) = \frac{2+3+3+4}{48} = \boxed{12/48 = 1/4}$.

(3) $P(X \le Y) = \sum_{x \le y} f(x,y)$. Sum all (x,y) with $x \le y$: $(0,0), (0,1), (0,2), (0,3), (1,1), (1,2), (1,3), (2,2), (2,3), (3,3)$. $= \frac{0+1+2+3+2+3+4+4+5+6}{48} = \boxed{30/48 = 5/8}$.

(4) $P(X+Y = 2) = f(0,2) + f(1,1) + f(2,0) = \frac{2+2+2}{48} = \boxed{6/48 = 1/8}$.

Exercise 3.51

Problem: Three cards are drawn without replacement from the 12 face cards (jacks, queens, and kings) of an ordinary deck of 52 playing cards. Let X be the number of kings selected and Y the number of jacks. Find (1) the joint probability distribution of X and Y; (2) $P[(X,Y) \in A]$, where $A = \{(x,y) \mid x+y \ge 2\}$.

Solution: $f(x,y) = \frac{\binom{4}{x}\binom{4}{y}\binom{4}{3-x-y}}{\binom{12}{3}}$ for $x+y \le 3$, $0 \le x,y \le 3$ (and $3-x-y \le 4$).

$\binom{12}{3} = 220$.

Selected values: $f(0,0) = \binom{4}{3}/220 = 4/220$, $f(1,0) = \binom{4}{1}\binom{4}{2}/220 = 24/220$, etc.

(2) $P(X+Y \ge 2) = 1 - P(X+Y = 0) - P(X+Y = 1)$. $P(X+Y = 0) = f(0,0) = 4/220$. $P(X+Y = 1) = f(1,0) + f(0,1) = 24/220 + 24/220 = 48/220$. $P(X+Y \ge 2) = 1 - 52/220 = \boxed{168/220 = 42/55}$.

Midterm Q4(a) – Joint PMF (4 pts)

Problem: X uniform on $\{-2,-1,0,1,2\}$ with $f(x) = 1/5$. $Y = X^2$. Find joint PMF.

Solution: Since Y is determined by X, $P(X = x, Y = y)$ is nonzero only when $y = x^2$:

$Y \backslash X$	-2	-1	0	1	2
0	0	0	1/5	0	0
1	0	1/5	0	1/5	0
4	1/5	0	0	0	1/5

Midterm Q4(d) – CDF of Discrete RV (4 pts)

Problem: Find the CDF of Y.

Solution: $F_Y(y) = \begin{cases} 0, & y < 0 \\ 1/5, & 0 \le y < 1 \\ 3/5, & 1 \le y < 4 \\ 1, & y \ge 4 \end{cases}$

3.10 Conditional Distributions (Discrete)

Def: The conditional PMF of Y given $X = x$ is

$$f_{Y|X}(y|x) = \frac{f(x,y)}{g(x)}, \quad g(x) > 0.$$

Similarly, the conditional PMF of X given $Y = y$ is

$$f_{X|Y}(x|y) = \frac{f(x,y)}{h(y)}, \quad h(y) > 0.$$

3.10.1 Example: Conditional Probabilities

1. $$P(Y = 1 \mid X = 1) = \frac{P(X = 1, Y = 1)}{P(X = 1)} = \frac{3/28}{15/28} = \frac{3}{15}.$$

2. $$P(X = 0 \mid Y = 1) = \frac{P(X = 0, Y = 1)}{P(Y = 1)} = \frac{9/28}{15/28} = \frac{9}{15}.$$

3. $$P(Y = 2 \mid X = 1) = 0.$$

Check:

$$\sum_y P(Y = y \mid X = 1) = 1.$$

1. *Random Variable* A random variable is a function that maps outcomes of an experiment to real numbers.
   - Domain: sample space outcomes
   - Range: real numbers
   - Can be **discrete** or **continuous**
2. *Probability Mass Function (PMF)* The PMF of a discrete random variable X is

$$f_X(x) = P(X = x)$$

- Only for discrete random variables
- $f_X(x) \geq 0$
- $\sum_x f_X(x) = 1$

3. *Probability Density Function (PDF)* For a continuous random variable X, probability is defined by

$$P(a \leq X \leq b) = \int_a^b f_X(x)\,dx$$

- Only for continuous random variables
- Area under the curve gives probability
- $P(X = x) = 0$

4. *Cumulative Distribution Function (CDF)* The CDF is defined as

$$F_X(x) = P(X \leq x)$$

- Discrete: $F(x) = \sum_{t \leq x} f(t)$
- Continuous: $F(x) = \int_{-\infty}^{x} f(t)\,dt$
- $0 \leq F(x) \leq 1$, non-decreasing

5. *Joint Distribution* The joint distribution describes probabilities involving two random variables X and Y.

- Discrete: $f_{X,Y}(x, y) = P(X = x, Y = y)$
- Continuous: joint PDF $f_{X,Y}(x, y)$

6. *Marginal Distribution* A marginal distribution is obtained by eliminating the other variable.

- $f_X(x) = \sum_y f_{X,Y}(x, y)$ or $f_X(x) = \int f_{X,Y}(x, y)\,dy$
- $f_Y(y) = \sum_x f_{X,Y}(x, y)$ or $f_Y(y) = \int f_{X,Y}(x, y)\,dx$

Exercise 3.56

Problem: $f(x, y) = 6x$ for $0 < x < 1$, $0 < y < 1 - x$. (1) Show X, Y not independent. (2) Find $P(X > 0.3 \mid Y = 0.5)$.
Solution: (1) Marginals: $f_X(x) = \int_0^{1-x} 6x\,dy = 6x(1 - x)$ for $0 < x < 1$. $f_Y(y) = \int_0^{1-y} 6x\,dx = 3(1 - y)^2$ for $0 < y < 1$. $f_X(x) \cdot f_Y(y) = 18x(1 - x)(1 - y)^2 \neq 6x = f(x, y)$. Not independent.
(2) $f_{X|Y}(x \mid 0.5) = \frac{f(x,0.5)}{f_Y(0.5)} = \frac{6x}{3(0.5)^2} = 8x$ for $0 < x < 0.5$. $P(X > 0.3 \mid Y = 0.5) = \int_{0.3}^{0.5} 8x\,dx = 4x^2\Big|_{0.3}^{0.5} = 1 - 0.36 = \boxed{0.64}$.

3.11 Joint Probability Density Function (Continuous)

Def: A function $f(x, y)$ is a joint probability density function (PDF) of continuous random variables X and Y if:

1. $f(x, y) \geq 0$ for all (x, y)

2. $$\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y)\, dx\, dy = 1$$

3. For any region A in the xy-plane,

$$P\big((X, Y) \in A\big) = \iint_A f(x, y)\, dx\, dy$$

Geometric interpretation: The joint PDF is a surface above the xy-plane. Probabilities correspond to the **volume under the surface** over a specified region.

Example 1:

$$f(x, y) = \begin{cases} \dfrac{12}{7}(x^2 + xy), & 0 \leq x \leq 1,\ 0 \leq y \leq 1 \\ 0, & \text{otherwise} \end{cases}$$

(a) Verify it is a valid joint PDF:

$$\int_0^1 \int_0^1 \frac{12}{7}(x^2 + xy)\, dx\, dy = 1$$

(b) Find $P(0 < X < 0.2,\ 0 < Y < 1)$:

$$P = \int_0^1 \int_0^{0.2} \frac{12}{7}(x^2 + xy)\, dx\, dy$$

(c) Find $P(X > Y)$:

The region $X > Y$ corresponds to the area below the line $y = x$ in the unit square $0 \leq x \leq 1,\ 0 \leq y \leq 1$.

$$P(X > Y) = \int_0^1 \int_0^x \frac{12}{7}(x^2 + xy)\, dy\, dx = \frac{9}{14}$$



(d) Find $P(X = Y)$:

$$P(X = Y) = \int_0^1 \int_y^y \frac{12}{7}(x^2 + xy)\,dx\,dy = 0$$

(Probability along a line is zero for continuous random variables.)

Example:

Let the joint PDF be

$$f(x, y) = \begin{cases} e^{-y}, & 0 < x < y < \infty \\ 0, & \text{otherwise} \end{cases}$$

We want to compute

$$P(X + Y \geq 1).$$

The support of the joint PDF is the region $0 < x < y$, which lies above the line $y = x$ in the first quadrant.

The boundary of the event $X + Y \geq 1$ is the line $x + y = 1$.

Rather than integrating over the unbounded region $X + Y \geq 1$, we compute the complement:

$$P(X + Y \geq 1) = 1 - P(X + Y < 1).$$

The region $X + Y < 1$ that lies within the support is bounded by:

$$0 \leq x \leq \tfrac{1}{2}, \qquad x \leq y \leq 1 - x.$$

Therefore,

$$P(X + Y \geq 1) = 1 - \int_0^{1/2} \int_x^{1-x} e^{-y}\,dy\,dx = 2e^{-1/2} - e^{-1}.$$



Exercise 3.63

Problem: $f(x, y) = ye^{-y(1+x)}$ for $x, y \geq 0$. (1) Marginals. (2) $P(X > 2, Y > 2)$.

Solution: (1) $f_X(x) = \int_0^\infty ye^{-y(1+x)}\,dy = \frac{1}{(1+x)^2}$ for $x \geq 0$. $f_Y(y) = \int_0^\infty ye^{-y(1+x)}\,dx = ye^{-y} \cdot \frac{1}{y} \cdot (1) = e^{-y}$ for $y > 0$ (Exponential(1)).

Wait—more carefully: $f_Y(y) = \int_0^\infty ye^{-y(1+x)}\,dx = ye^{-y} \int_0^\infty e^{-yx}\,dx = ye^{-y} \cdot \frac{1}{y} = e^{-y}$.

(2) $P(X > 2, Y > 2) = \int_2^\infty \int_2^\infty ye^{-y(1+x)}\,dx\,dy = \int_2^\infty ye^{-y} \cdot \frac{e^{-2y}}{y}\,dy = \int_2^\infty e^{-3y}\,dy = \frac{e^{-6}}{3} \approx \boxed{0.000826}$.

3.12 Marginal and Conditional PDFs

Def: The marginal PDFs of X and Y are defined as

$$g_X(x) = \int_{-\infty}^{\infty} f(x,y)\,dy, \qquad h_Y(y) = \int_{-\infty}^{\infty} f(x,y)\,dx$$

Example (continued):

$$f(x,y) = \begin{cases} \frac{12}{7}(x^2+xy), & 0 \le x \le 1,\ 0 \le y \le 1 \\ 0, & \text{otherwise} \end{cases}$$

Marginal PDF of X:

$$g_X(x) = \int_0^1 \frac{12}{7}(x^2+xy)\,dy = \frac{12}{7}\left(x^2 + \frac{x}{2}\right), \qquad 0 \le x \le 1$$

Marginal PDF of Y:

$$h_Y(y) = \int_0^1 \frac{12}{7}(x^2+xy)\,dx = \frac{12}{7}\left(\frac{1}{3} + \frac{y}{2}\right), \qquad 0 \le y \le 1$$

Conditional PDF of Y given $X = x$:

$$f(y|x) = \frac{f(x,y)}{g(x)} = \frac{\frac{12}{7}(x^2+xy)}{\frac{12}{7}\left(x^2+\frac{x}{2}\right)}, \qquad 0 \le y \le 1,\ 0 < x \le 1$$

Why the bounds are $0 < x \le 1$ and not $0 \le x \le 1$:

The marginal PDF

$$g(x) = \frac{12}{7}\left(x^2 + \frac{x}{2}\right)$$

satisfies $g(0) = 0$.

Since the conditional PDF is defined as

$$f_{Y|X}(y|x) = \frac{f(x,y)}{g(x)},$$

it is **undefined at** $x = 0$ due to division by zero.

Therefore, the conditional density is only defined for values of x such that

$$g(x) > 0 \ \Rightarrow\ 0 < x \le 1.$$

Key takeaway: The bounds of a conditional PDF exclude points where the conditioning density is zero.

Midterm Q6(a) – Marginal Densities (5 pts)

Problem: $f_{X,Y}(x,y) = \frac{1}{\pi a^2}$ for $x^2 + y^2 \leq a^2$. Find marginals.
Solution: $f_X(x) = \int_{-\sqrt{a^2-x^2}}^{\sqrt{a^2-x^2}} \frac{1}{\pi a^2}\, dy = \frac{2\sqrt{a^2-x^2}}{\pi a^2}$, for $-a \leq x \leq a$.
By symmetry, $f_Y(y) = \frac{2\sqrt{a^2-y^2}}{\pi a^2}$, for $-a \leq y \leq a$.

Midterm Q6(b) – Expected Values (4 pts)

Problem: Find $E(X)$ and $E(Y)$.
Solution: Both are $\boxed{0}$ by symmetry (odd function integrated over symmetric interval).

3.13 Statistical Independence

Def: Random variables X and Y (discrete or continuous) are statistically independent if and only if

$$f(x,y) = g(x)\,h(y) \quad \text{for all } (x,y) \text{ in their range}$$

Consequences:

- $f(x|y) = g(x)$
- $f(y|x) = h(y)$

3.13.1 Recall Example: Pen Refills

Two refills are selected at random and without replacement from a box containing:

- 3 blue refills
- 2 red refills
- 3 green refills

Define the random variables

$$X = \text{number of blue refills selected}, \qquad Y = \text{number of red refills selected}.$$

Total outcomes:

$$\binom{8}{2} = 28.$$

3.13.2 Joint PMF Table

For each pair (x,y),

$$f(x,y) = \frac{\text{number of favorable outcomes}}{28}.$$

$X \backslash Y$	0	1	2	$g(x) = P(X = x)$
0	$\frac{3}{28}$	$\frac{3}{14}$	$\frac{1}{28}$	$\frac{5}{14}$
1	$\frac{9}{28}$	$\frac{3}{14}$	0	$\frac{15}{28}$
2	$\frac{3}{28}$	0	0	$\frac{3}{28}$
$h(y) = P(Y = y)$	$\frac{15}{28}$	$\frac{3}{7}$	$\frac{1}{28}$	1

Marginals: $g(x) = P(X = x)$ is the marginal PMF of X and is given by the **row totals**. $h(y) = P(Y = y)$ is the marginal PMF of Y and is given by the **column totals**.

$$g(0) = \frac{5}{14}, \quad g(1) = \frac{15}{28}, \quad g(2) = \frac{3}{28}$$
$$h(0) = \frac{15}{28}, \quad h(1) = \frac{3}{7}, \quad h(2) = \frac{1}{28}$$

Independence check: If X and Y were statistically independent, then $f(x, y) = g(x)h(y)$ for all (x, y). Check $(x, y) = (0, 1)$:

$$f(0,1) = \frac{3}{14}, \qquad g(0)h(1) = \left(\frac{5}{14}\right)\left(\frac{3}{7}\right) = \frac{15}{98}$$
$$\frac{3}{14} \neq \frac{15}{98} \quad \Rightarrow \quad X \text{ and } Y \text{ are not independent.}$$

3.13.3 Example: Independence via Factorization (Discrete Case)

Let X and Y be discrete random variables whose values in the nonnegative integers.

Suppose the joint PMF is

$$f(x, y) = \frac{1}{x!\,y!}\lambda^x \mu^y e^{-(\lambda+\mu)}, \qquad x, y = 0, 1, 2, \ldots$$

Factorization:

We can write

$$f(x, y) = \left(\frac{\lambda^x e^{-\lambda}}{x!}\right)\left(\frac{\mu^y e^{-\mu}}{y!}\right) = g(x)\,h(y)$$

Marginal PMF of X:

$$g(x) = \sum_{y=0}^{\infty} f(x, y) = \sum_{y=0}^{\infty} \frac{1}{x!\,y!}\,\lambda^x \mu^y\, e^{-(\lambda+\mu)}$$

Factor out terms that do not depend on y:

$$g(x) = \frac{1}{x!}\,\lambda^x e^{-(\lambda+\mu)} \sum_{y=0}^{\infty} \frac{\mu^y}{y!}$$

Using $\displaystyle\sum_{y=0}^{\infty} \frac{\mu^y}{y!} = e^{\mu}$:

$$g(x) = \frac{1}{x!}\,\lambda^x e^{-(\lambda+\mu)} e^{\mu} = \frac{1}{x!}\,\lambda^x e^{-\lambda}, \qquad x = 0, 1, 2, \ldots$$

Marginal PMF of Y:

$$h(y) = \sum_{x=0}^{\infty} f(x,y) = \sum_{x=0}^{\infty} \frac{1}{x!\,y!}\,\lambda^x \mu^y\, e^{-(\lambda+\mu)}$$

Factor out terms that do not depend on x:

$$h(y) = \frac{1}{y!}\,\mu^y e^{-(\lambda+\mu)} \sum_{x=0}^{\infty} \frac{\lambda^x}{x!}$$

Using $\displaystyle\sum_{x=0}^{\infty} \frac{\lambda^x}{x!} = e^{\lambda}$:

$$h(y) = \frac{1}{y!}\,\mu^y e^{-(\lambda+\mu)} e^{\lambda} = \frac{1}{y!}\,\mu^y e^{-\mu}, \qquad y = 0, 1, 2, \ldots$$

Conclusion:

Since the joint PMF can be written as

$$f(x,y) = g(x)\,h(y)$$

for all x, y, the random variables X and Y are <u>statistically independent</u>. ✓

Important notes:

- Factorization of the joint PMF is **sufficient** to prove independence.
- The constant terms (such as $e^{-\lambda}$, $e^{-\mu}$) must be included to obtain the **correct marginals**.
- Independence requires that every combination of values with positive marginal probability also has positive joint probability.
  This means that if $X = x$ is possible and $Y = y$ is possible, then the pair $(X = x, Y = y)$ must also be possible.

  $$\{(x,y) : f(x,y) > 0\} = \{x : g(x) > 0\} \times \{y : h(y) > 0\}.$$

  This means that every value of X with positive marginal probability can occur with every value of Y with positive marginal probability.

3.13.4 Joint distribution

Describes probabilities involving two random variables X and Y.

- Discrete:
$$f(X,Y) = P(X = x, Y = y)$$
- Continuous: joint PDF $f(X,Y)$

3.13.5 Marginal distribution

Obtained by eliminating the other variable.

- $$f(X) = \sum_y f(X,Y), \qquad f(Y) = \sum_x f(X,Y)$$
- $$f(X) = \int_{-\infty}^{\infty} f(X,Y)\,dy, \qquad f(Y) = \int_{-\infty}^{\infty} f(X,Y)\,dx$$

3.13.6 Joint PDF validity

A function $f(X,Y)$ is a valid joint PDF if:

- $f(X,Y) \geq 0$
- $$\int_{-\infty}^{\infty}\int_{-\infty}^{\infty} f(X,Y)\,dx\,dy = 1$$

Example (valid joint PDF):

$$f(X,Y) = \begin{cases} \frac{12}{7}(x^2 + xy), & 0 \leq x \leq 1,\ 0 \leq y \leq 1, \\ 0, & \text{otherwise} \end{cases}$$

3.13.7 Geometric interpretation

Probabilities correspond to the **volume under the surface** $f(X,Y)$ over a region.

3.13.8 Conditional PDF

Distribution of one variable given the other.

- $$f(Y \mid X) = \frac{f(X,Y)}{f(X)}, \quad f(X) > 0$$

3.13.9 Statistical Independence

Definition: Random variables X and Y are statistically independent if

$$f(X,Y) = f(X)f(Y)$$

This means knowing the value of one variable gives no information about the other.

3.13.10 Support and Independence

Support: The support of a joint distribution is the set

$$\{(x,y) : f(X,Y) > 0\}$$

Key idea: If X and Y are independent, the support must factor as

$$\{x : f(X) > 0\} \times \{y : f(Y) > 0\}$$

That is, every allowed value of X can occur with every allowed value of Y.

3.13.11 Example: Non-Independent Random Variables

$$f(X,Y) = \begin{cases} 4(x+y^2), & xy > 0,\ x+y \leq 1, \\ 0, & \text{otherwise} \end{cases}$$

The condition $x + y \leq 1$ links x and y, so the support does not factor.

$$\Rightarrow X \text{ and } Y \text{ are not independent}$$

Why this implies dependence (key intuition):

Pick a value of X that is allowed:

$$x = 0.8 \quad (\text{positive and } < 1)$$

Pick a value of Y that is allowed:

$$y = 0.8 \quad (\text{positive and } < 1)$$

Individually, both values are valid. But together:

$$x + y = 0.8 + 0.8 = 1.6 > 1$$

This violates the condition $x + y \leq 1$, so the pair $(0.8, 0.8)$ is impossible.

Conclusion: Knowing the value of X restricts which values Y can take. Therefore, X and Y are not independent.

3.13.12 Independent Random Variables

If random variables are independent, joint probabilities factor.

Example: Let X_1, X_2, X_3 be independent with

$$f(X) = e^{-x}, \quad x > 0$$

Then

$$f(X_1, X_2, X_3) = e^{-x_1} e^{-x_2} e^{-x_3}$$

$$P(X_1 < 2,\ 1 < X_2 < 3,\ X_3 > 2) = (1 - e^{-2})(e^{-1} - e^{-3})e^{-2}$$

Explanation (why this works):

Since X_1, X_2, X_3 are <u>independent</u>, the joint probability factors:

$$P(X_1 < 2,\ 1 < X_2 < 3,\ X_3 > 2) = P(X_1 < 2)\, P(1 < X_2 < 3)\, P(X_3 > 2)$$

For an exponential random variable with

$$f(X) = e^{-x}, \quad x > 0,$$

we have:

$$P(X < a) = 1 - e^{-a}, \qquad P(X > a) = e^{-a}$$

Thus,

$$P(X_1 < 2) = 1 - e^{-2}$$

$$P(1 < X_2 < 3) = P(X_2 < 3) - P(X_2 < 1) = e^{-1} - e^{-3}$$

$$P(X_3 > 2) = e^{-2}$$

Multiplying gives:

$$(1 - e^{-2})(e^{-1} - e^{-3})e^{-2}$$

3.13.13 Mutual Independence and Modeling Assumptions

Mutual independence: Random variables $X_1, \ldots, X_n$ are mutually independent if

$$f(X_1, \ldots, X_n) = \prod_{i=1}^{n} f(X_i)$$

Pairwise independence does **not** imply mutual independence.

Independent selection: Selections are independent if:

- each selection is random,

- distributions are identical,
- outcomes do not affect future selections.

Sampling without replacement generally produces dependent variables.

Chapter 4

Chapter 4: Mathematical Expectation

4.0.1 Motivation

Example: If you roll a fair die repeatedly, what *average value* do you expect in the long run?

This motivates the idea of expectation: a theoretical long-run average.

4.0.2 Expected Value

4.0.3 Expected Value of a Random Variable

Def: For a random variable X, the expectation (or expected value or mean) is the long-run average value of X.

Discrete random variable:

$$\boxed{\mu = E(X) = \sum_x x\, f(X)}$$

Continuous random variable:

$$\boxed{\mu = E(X) = \int_{-\infty}^{\infty} x\, f(X)\, dx}$$

Interpretation: Expectation is a weighted average, where values of X are weighted by how likely they are.

4.0.4 Example: Fair Die

Let X be the outcome when a fair die is rolled.

$$E(X) = \sum_{x=1}^{6} x \cdot \frac{1}{6}$$

$$E(X) = \frac{1+2+3+4+5+6}{6} = 3.5$$

Even though 3.5 is not a possible outcome, it represents the long-run average.

4.0.5 Example: Number of Messages per Hour

Let X be the number of messages sent per hour, with PMF:

x	10	11	12	13	14	15
$f(X)$	0.08	0.15	0.30	0.20	0.20	0.07

Check:

$$\sum f(X) = 1$$

Expected value:

$$\begin{aligned} E(X) &= 10(0.08) + 11(0.15) + 12(0.30) + 13(0.20) + 14(0.20) + 15(0.07) \\ &= 12.5 \end{aligned}$$

This means that over many hours, the average number of messages per hour is about 12.5.

4.0.6 PMF Visualization



4.0.7 Example: Deal or No Deal

Consider a game with two possible outcomes:

- \$1 with probability 1/2
- \$10,000 with probability 1/2

Expected value:

$$E(X) = \frac{1}{2}(1) + \frac{1}{2}(10{,}000) = 5000.5$$

Key idea: The expected value is the average payout in the long run, not the most likely outcome.

4.0.8 Example: Continuous RV (Device Lifetime)

Let X be a random variable that denotes the lifetime (in hours) of a certain device, with PDF

$$f(X) = \begin{cases} \dfrac{20000}{x^3}, & x > 100, \\ 0, & \text{otherwise.} \end{cases}$$

Check (valid PDF):

$$\int_{-\infty}^{\infty} f(X)\,dx = \int_{100}^{\infty} \frac{20000}{x^3}\,dx = 20000\left[\frac{-1}{2x^2}\right]_{100}^{\infty} = 1$$

Expected lifetime:

$$E(X) = \int_{100}^{\infty} x\frac{20000}{x^3}\,dx = \int_{100}^{\infty} \frac{20000}{x^2}\,dx = 20000\left[\frac{-1}{x}\right]_{100}^{\infty} = 200$$

So we expect this type of device to last on average $\boxed{200}$ hours.

4.0.9 Example: Discrete RV and Transformation

Let X be a discrete random variable with PMF:

x	-1	0	1	2
$f(X)$	0.3	0.2	0.3	0.2

Define a new random variable as a <u>transformation</u> of X:

$$g(X) = X^2$$

Possible values of $g(X)$:

$$g(-1) = 1, \quad g(0) = 0, \quad g(1) = 1, \quad g(2) = 4$$

So $g(X)$ can take values $\{0, 1, 4\}$.

Distribution of $g(X)$:

$$P(g(X) = 0) = P(X = 0) = 0.2$$

$$P(g(X) = 1) = P(X = -1) + P(X = 1) = 0.3 + 0.3 = 0.6$$

$$P(g(X) = 4) = P(X = 2) = 0.2$$

$g(X)$	0	1	4
$P(g(X))$	0.2	0.6	0.2

This is called a <u>transformation</u> of a random variable.

Expected value of the transformed RV:

$$\boxed{E(g(X)) = E(X^2) = \sum_x x^2 f(X) = \sum_x g(x)\,f(X)}$$

Numerically:

$$E(X^2) = 0^2(0.2) + (-1)^2(0.3) + (1)^2(0.3) + (2)^2(0.2) = 0 + 0.3 + 0.3 + 0.8 = 1.4$$

Expected Value of a Function of a RV

Let X be a random variable with distribution $f(X)$. The expected value of the random variable $g(X)$ is

$$\boxed{E(g(X)) = \begin{cases} \sum_x g(x)\, f(X), & \text{if } X \text{ is discrete} \\ \int_{-\infty}^{\infty} g(x)\, f(X)\, dx, & \text{if } X \text{ is continuous} \end{cases}}$$

4.0.10 Example: Chip Game (Expected Winnings)

A bowl contains 5 chips:

- 3 chips are worth \$1 each
- 2 chips are worth \$4 each

A player draws 2 chips at random (without replacement) and is paid the sum.

Let X be the number of \$1 chips drawn. Then

$$X \in \{0, 1, 2\}.$$

PMF of X: (hypergeometric)

$$f(X) = \begin{cases} \dfrac{\binom{3}{x}\binom{2}{2-x}}{\binom{5}{2}}, & x = 0, 1, 2, \\ 0, & \text{otherwise.} \end{cases}$$

Define payout as a function of X: If you draw x one-dollar chips, then you draw $2 - x$ four-dollar chips, so the payout is

$$g(x) = 1(x) + 4(2 - x) = 8 - 3x.$$

So the payout random variable is $g(X) = 8 - 3X$.

Expected payout:

$$E(g(X)) = \sum_{x=0}^{2} g(x)\, f(X)$$

Compute $f(X)$ values:

$$f(0) = \frac{\binom{3}{0}\binom{2}{2}}{\binom{5}{2}} = \frac{1}{10}, \quad f(1) = \frac{\binom{3}{1}\binom{2}{1}}{\binom{5}{2}} = \frac{6}{10}, \quad f(2) = \frac{\binom{3}{2}\binom{2}{0}}{\binom{5}{2}} = \frac{3}{10}.$$

Then

$$E(g(X)) = \sum_{x=0}^{2} (8 - 3x)\, f(X) = (8)\left(\frac{1}{10}\right) + (5)\left(\frac{6}{10}\right) + (2)\left(\frac{3}{10}\right) = 4.4$$

Decision: If it costs \$4.75 to play, your expected profit is

$$E(\text{profit}) = E(g(X)) - 4.75 = 4.4 - 4.75 = -0.35$$

So in the long run, you lose about $\boxed{\$0.35}$ per game on average, so you should <u>not</u> play.

4.0.11 Notation: $g(X)$ vs. $g(x)$

- X is a <u>random variable</u>; x is a specific value it can take.
- $g(X)$ is a <u>random variable</u>.
- $g(x)$ is a <u>number</u>.

Key rule:

$$\boxed{E(g(X)) = \sum_x g(x)\, f(X)}$$

Expectation averages the numerical values $g(x)$, weighted by their probabilities.

Feb 2nd

1. <u>*Expected value of a function: expected average value in the long wrong*</u>, $E[g(X)]$
   - $g(X)$ is a function of X
   - **Discrete:**
   $$E[g(X)] = \sum_x g(x)\, f(X)$$
   - **Continuous:**
   $$E[g(X)] = \int_{-\infty}^{\infty} g(X)\, f(X)\, dx$$
2. **Example (continuous RV):**

$$f(X) = 4x^3, \quad 0 < x < 1$$

- $$E(X) = \int_0^1 x \cdot 4x^3\, dx$$
- $$E(X^2) = \int_0^1 x^2 \cdot 4x^3\, dx$$

4.0.12 Expected Value from a Joint Distribution

General formula

$$\boxed{E[g(X,Y)] = \iint g(x,y)\, f(X,Y)\, dx\, dy}$$

Computing $E(X)$ from a joint PDF

There are two equivalent methods to compute $E(X)$.

Method 1: Using the marginal distribution

First find the marginal of X:

$$f(X) = \int_0^1 x(1+3y^2)\,dy$$

$$= x\int_0^1 (1+3y^2)\,dy$$

$$= x\left[y+y^3\right]_0^1 = x(1+1) = 2x$$

Now compute the expected value:

$$E(X) = \int_0^2 xf(X)\,dx = \int_0^2 x(2x)\,dx$$

$$= 2\int_0^2 x^2\,dx = 2\left[\frac{x^3}{3}\right]_0^2 = \frac{16}{3}$$

Method 2: Direct double integral

Apply the general formula with $g(X,Y) = X$:

$$E(X) = \int_0^1\int_0^2 x\,f(X,Y)\,dx\,dy$$

$$= \int_0^1\int_0^2 x\cdot x(1+3y^2)\,dx\,dy$$

$$= \int_0^1 (1+3y^2)\left(\int_0^2 x^2\,dx\right)dy$$

$$= \int_0^1 (1+3y^2)\left[\frac{x^3}{3}\right]_0^2 dy = \int_0^1 (1+3y^2)\frac{8}{3}\,dy$$

$$= \frac{8}{3}\int_0^1 (1+3y^2)\,dy = \frac{8}{3}\cdot 2 = \frac{16}{3}$$

Both methods give the same result:

$$\boxed{E(X) = \frac{16}{3}}$$

4.0.13 Properties of Expected Value

Properties of Expectation

Let X, Y be random variables and let $a, b \in \mathbb{R}$ be constants.

1. **Linearity (scaling and shifting):**

$$\boxed{E(aX + b) = aE(X) + b}$$

Continuous case:

$$E(aX + b) = \int_{-\infty}^{\infty} (ax + b)\, f(X)\, dx$$

$$= a \int_{-\infty}^{\infty} x f(X)\, dx + b \int_{-\infty}^{\infty} f(X)\, dx$$

Note that

$$\int_{-\infty}^{\infty} f(X)\, dx = 1$$

because $f(X)$ is a probability density function and the total probability over its entire support must equal 1.

Therefore,

$$E(aX + b) = aE(X) + b$$

2. **Expectation of a constant:**

$$\boxed{E(a) = a}$$

3. **Additivity:**

$$\boxed{E[g(X) \pm h(X)] = E[g(X)] \pm E[h(X)]}$$

4. **Non-negativity:**

$$\boxed{X \geq 0 \;\Rightarrow\; E(X) \geq 0}$$

5. **Zero expectation:**

$$\boxed{X \geq 0 \text{ and } E(X) = 0 \iff P(X = 0) = 1}$$

4.1 Variance

$E(X)$ is a measure of the **center** of a distribution.

The **variance**, denoted $\sigma^2 = \mathrm{Var}(X)$, measures how closely the distribution is concentrated around the mean μ.

Definition:

$$\boxed{\mathrm{Var}(X) = E[(X - \mu)^2], \quad \mu = E(X)}$$

$\sigma = \sqrt{\sigma^2}$ is called the **standard deviation**.

Theorem (Variance Formula)

$$\mathrm{Var}(X) = \sigma^2 = E(X^2) - \mu^2$$

Proof:

$$E[(X-\mu)^2] = E[X^2 - 2\mu X + \mu^2]$$
$$= E(X^2) - 2\mu E(X) + \mu^2$$
$$= E(X^2) - \mu^2$$

Example

Weekly demand for a drink (in thousand liters) is a continuous random variable X with PDF

$$f(X) = 2(x-1), \quad 1 < x < 2$$

Mean:

$$\mu = E(X) = \int_1^2 x \cdot 2(x-1)\,dx = \frac{5}{3}$$

Second moment:

$$E(X^2) = \int_1^2 x^2 \cdot 2(x-1)\,dx = \frac{17}{6}$$

Variance:

$$\boxed{\sigma^2 = \mathrm{Var}(X) = \frac{17}{6} - \left(\frac{5}{3}\right)^2 = \frac{1}{18}}$$

Variance of a function of a random variable

Let X be a random variable and let $g(X)$ be a function of X.

Mean of $g(X)$:

$$\mu_{g(X)} = E[g(X)]$$

Variance of $g(X)$:

$$\boxed{\mathrm{Var}(g(X)) = E[(g(X) - \mu_{g(X)})^2] = E[g(X)^2] - (E[g(X)])^2}$$

Properties of Variance

Let X, Y be random variables and let $a, b \in \mathbb{R}$.

1.
$$\boxed{\mathrm{Var}(aX+b) = a^2\mathrm{Var}(X)}$$

Proof:

$$\mathrm{Var}(aX+b) = E[(aX+b)^2] - (E[aX+b])^2$$
$$= E[a^2X^2 + 2abX + b^2] - (aE[X]+b)^2$$
$$= a^2E[X^2] + 2abE[X] + b^2 - a^2E[X]^2 - 2abE[X] - b^2$$
$$= a^2\big(E[X^2] - E[X]^2\big) = a^2\mathrm{Var}(X)$$

2.
$$\boxed{\mathrm{Var}(X) \geq 0}$$

3.
$$\boxed{\mathrm{Var}(a) = 0} \quad \text{for any constant } a$$

4.
$$\boxed{\mathrm{Var}(X) = 0 \iff X \text{ is a constant}}$$

Theorem (Variance of a Linear Combination)

Let X, Y be random variables with joint distribution $f(X,Y)$. Then

$$\mathrm{Var}(aX+bY) = a^2\mathrm{Var}(X) + b^2\mathrm{Var}(Y) + 2ab\,\mathrm{Cov}(X,Y)$$

where the **covariance** is defined as

$$\boxed{\mathrm{Cov}(X,Y) = E[(X-E[X])(Y-E[Y])]}$$

Interpretation of covariance

- $\mathrm{Cov}(X,Y) > 0$: X and Y tend to increase together
- $\mathrm{Cov}(X,Y) < 0$: one increases while the other decreases
- $\mathrm{Cov}(X,Y) = 0$: no linear relationship

Example (population intuition):

If X is height and Y is weight in a population,

$$(X-E[X])(Y-E[Y]) > 0$$

for tall–heavy and short–light individuals, so $\mathrm{Cov}(X,Y) > 0$.

Feb 4

1. *Expected Value*

   - **Discrete random variable:**
   $$E(X) = \sum_x x f(X)$$

   - **Continuous random variable:**
   $$E(X) = \int_{-\infty}^{\infty} x f(X)\, dx$$

2. *Variance*

   - Definition:
   $$\mathrm{Var}(X) = \sigma^2 = E\big[(X-\mu)^2\big]$$

   - Simplified form:
   $$\mathrm{Var}(X) = E(X^2) - \mu^2$$

   where $\mu = E(X)$.

3. *Variance of a Linear Combination*

   - For constants a, b:
   $$\mathrm{Var}(aX + bY) = a^2\mathrm{Var}(X) + b^2\mathrm{Var}(Y) + 2ab\,\sigma_{XY}$$

   - Covariance:
   $$\sigma_{XY} = \mathrm{Cov}(X,Y) = E\big[(X - E(X))(Y - E(Y))\big]$$

   - Equivalent form:
   $$\sigma_{XY} = E(XY) - E(X)E(Y)$$

Theorem: Independence and Expected Value

Let X and Y be independent random variables. Then

$$E(XY) = E(X)\,E(Y).$$

As a consequence,

$$\mathrm{Cov}(X, Y) = 0.$$

Continuous case:

$$E(XY) = \int_{-\infty}^{\infty}\int_{-\infty}^{\infty} xy\, f(X, Y)\, dx\, dy$$

If X and Y are independent,

$$f(X, Y) = g(X)h(Y)$$

so

$$E(XY) = \left(\int_{-\infty}^{\infty} xg(X)\, dx\right)\left(\int_{-\infty}^{\infty} yh(Y)\, dy\right)$$

$$= E(X)\,E(Y)$$

4.1.1 Example

Given: The joint PMF table below.

$f(X,Y)$	$X=-1$	$X=0$	$X=1$	$h(Y)$
$Y=-1$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{3}{8}$
$Y=0$	$\frac{1}{8}$	0	$\frac{1}{8}$	$\frac{2}{8}$
$Y=1$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{1}{8}$	$\frac{3}{8}$
$g(X)$	$\frac{3}{8}$	$\frac{2}{8}$	$\frac{3}{8}$	1

1. <u>*Compute $E(X)$*</u>

   Using the marginal PMF $g(X)$:

$$E(X) = \sum_x x\, g(X)$$

$$E(X) = (-1)\cdot\tfrac{3}{8} + 0\cdot\tfrac{2}{8} + 1\cdot\tfrac{3}{8} = -\tfrac{3}{8} + 0 + \tfrac{3}{8} = 0$$

2. <u>*Compute $E(Y)$*</u>

   Using the marginal PMF $h(Y)$:

$$E(Y) = \sum_y y\, h(Y)$$

$$E(Y) = (-1)\cdot\tfrac{3}{8} + 0\cdot\tfrac{2}{8} + 1\cdot\tfrac{3}{8} = -\tfrac{3}{8} + 0 + \tfrac{3}{8} = 0$$

3. <u>*Compute $E(XY)$*</u>

Using the joint PMF:

$$E(XY) = \sum_x \sum_y xy\, f(X,Y)$$

Compute by rows:

Row $y = -1$:

$$(-1)(-1)\tfrac{1}{8} + (0)(-1)\tfrac{1}{8} + (1)(-1)\tfrac{1}{8} = \tfrac{1}{8} + 0 - \tfrac{1}{8} = 0$$

Row $y = 0$: all terms are 0 because $y = 0$.

Row $y = 1$:

$$(-1)(1)\tfrac{1}{8} + (0)(1)\tfrac{1}{8} + (1)(1)\tfrac{1}{8} = -\tfrac{1}{8} + 0 + \tfrac{1}{8} = 0$$

Therefore,

$$E(XY) = 0$$

Final:

$$E(X) = 0, \qquad E(Y) = 0, \qquad E(XY) = 0$$

Note: Even if $E(XY) = E(X)E(Y)$, this <u>does not automatically mean</u> X and Y are independent. Independence requires:

$$f(X,Y) = g(X)\, h(Y).$$

Exercise 4.6

Problem: An attendant at a car wash is paid according to the number of cars that pass through. Suppose the probabilities are 1/12, 1/12, 1/4, 1/4, 1/6, and 1/6, respectively, that the attendant receives \$7, \$9, \$11, \$13, \$15, or \$17 between 4:00 P.M. and 5:00 P.M. on any sunny Friday. Find the attendant's expected earnings for this particular period.

Solution: $E = 7(1/12) + 9(1/12) + 11(1/4) + 13(1/4) + 15(1/6) + 17(1/6) = 7/12 + 9/12 + 33/12 + 39/12 + 30/12 + 34/12 = 152/12 = \boxed{\$12.67}$.

Exercise 4.7

Problem: By investing in a particular stock, a person can make a profit in one year of \$4000 with probability 0.3 or take a loss of \$1000 with probability 0.7. What is this person's expected gain?

Solution: $E = 4000(0.3) + (-1000)(0.7) = 1200 - 700 = \boxed{\$500}$.

Exercise 4.12

Problem: If a dealer's profit, in units of \$5000, on a new automobile can be looked upon as a random variable X having the density function $f(x) = 2(1-x)$ for $0 < x < 1$ (and 0 elsewhere), find the average profit per automobile.

Solution: $E(X) = \int_0^1 x \cdot 2(1-x)\, dx = 2\int_0^1 (x - x^2)\, dx = 2\left[\frac{1}{2} - \frac{1}{3}\right] = 2 \cdot \frac{1}{6} = \boxed{1/3}$. Average profit $= (1/3)(5000) = \$1667$.

Exercise 4.21

Problem: What is the dealer's average profit per automobile if the profit on each automobile is given by $g(X) = X^2$, where X is a random variable having the density function of Exercise 4.12?
Solution: $E(X^2) = \int_0^1 x^2 \cdot 2(1-x)\,dx = 2\int_0^1 (x^2 - x^3)\,dx = 2\left[\frac{1}{3} - \frac{1}{4}\right] = 2 \cdot \frac{1}{12} = \boxed{1/6}$. Average profit $= (1/6)(5000) = \$833.33$.

Exercise 4.25

Problem: Referring to the random variables whose joint probability distribution is given in Exercise 3.51, find the mean for the total number of kings and jacks selected.
Solution:
From Ex. 3.51, three cards are drawn without replacement from the 12 face cards (4 kings, 4 jacks, 4 queens). The joint PMF of X (kings) and Y (jacks) is:

$$P(X = x,\, Y = y) = \frac{\binom{4}{x}\binom{4}{y}\binom{4}{3-x-y}}{\binom{12}{3}}, \quad x, y \geq 0,\ x + y \leq 3.$$

$E(X) = E(Y)$ by symmetry (4 kings, 4 jacks from 12). Each $E = \frac{n \cdot K}{N} = \frac{3 \cdot 4}{12} = \boxed{1}$.
$E(X+Y) = E(X) + E(Y) = 1 + 1 = 2$.

4.2 Covariance and Independence

Recall:

$$\mathrm{Cov}(X, Y) = 0 \quad \text{does NOT necessarily imply independence.}$$

Check independence:

$$f(X, Y) \stackrel{?}{=} g(X)h(Y)$$

In this example,

$$f(0, 0) = 0$$

but

$$g(0)h(0) = \frac{2}{8} \cdot \frac{2}{8} \neq 0$$

Therefore,

$$f(0, 0) \neq g(0)h(0)$$

$$\Rightarrow \quad X \text{ and } Y \text{ are } \underline{\text{not independent}}.$$

Properties of Covariance

1.
$$\mathrm{Cov}(X, X) = \mathrm{Var}(X) = E(X^2) - E(X)^2$$

2.
$$\mathrm{Cov}(aX + b,\ cY + d) = ac\,\mathrm{Cov}(X, Y)$$

3.
$$\mathrm{Cov}(X + Y,\ Z) = \mathrm{Cov}(X, Z) + \mathrm{Cov}(Y, Z)$$

4.
$$\mathrm{Cov}(X, Y) = \mathrm{Cov}(Y, X)$$

Important note:

The covariance σ_{XY} is <u>not scale-free</u>. Its magnitude does <u>not</u> directly indicate the strength of the linear relationship between X and Y.

Definition: Correlation Coefficient

The <u>correlation coefficient</u> of X and Y is

$$\boxed{\rho_{XY} = \frac{\sigma_{XY}}{\sigma_X\,\sigma_Y}}$$

where

$$\sigma_X = \sqrt{\mathrm{Var}(X)}, \qquad \sigma_Y = \sqrt{\mathrm{Var}(Y)}.$$

Theorem: Correlation Is Scale-Free

For constants $a, c \neq 0$ and any constants b, d,

$$\rho(aX + b,\ cY + d) = \rho(X, Y)$$

Therefore, the correlation coefficient is <u>scale-free</u>.

Proof.

By definition of correlation,

$$\rho(aX + b,\ cY + d) = \frac{\mathrm{Cov}(aX + b,\ cY + d)}{\sqrt{\mathrm{Var}(aX + b)\,\mathrm{Var}(cY + d)}}.$$

Using properties of covariance,

$$\mathrm{Cov}(aX + b,\ cY + d) = ac\,\mathrm{Cov}(X, Y)$$

Using properties of variance,

$$\mathrm{Var}(aX + b) = a^2\mathrm{Var}(X), \qquad \mathrm{Var}(cY + d) = c^2\mathrm{Var}(Y)$$

Substituting,

$$\rho(aX+b,\ cY+d) = \frac{ac\,\mathrm{Cov}(X,Y)}{\sqrt{a^2\mathrm{Var}(X)\,c^2\mathrm{Var}(Y)}}.$$

The constants a and c appear in both the numerator and denominator and therefore cancel:

$$= \frac{\mathrm{Cov}(X,Y)}{\sqrt{\mathrm{Var}(X)\mathrm{Var}(Y)}} = \rho(X,Y)$$

Theorem: Bounds on Correlation

$$-1 \le \rho(X,Y) \le 1$$

4.2.1 Interpretation of the Correlation Coefficient

1. ρ is a measure of the strength and direction of the linear relationship between X and Y.
2. When there is an exact linear relationship,

$$\boxed{Y = aX + b} \quad \Rightarrow \quad \boxed{\rho = 1 \text{ or } \rho = -1}$$

4.2.2 Example

Given:

$$f(X,Y) = \begin{cases} 1, & 0 < x < 1,\ x < y < x+1, \\ 0, & \text{otherwise.} \end{cases}$$



Check total probability:

$$\int_0^1 \int_x^{x+1} 1\,dy\,dx = \int_0^1 (x+1-x)\,dx = \int_0^1 1\,dx = 1$$

4.2.3 Expected Values

$$E(X) = \int_0^1 \int_x^{x+1} x\, f(X,Y)\, dy\, dx = \int_0^1 \int_x^{x+1} x\, dy\, dx$$

$$= \int_0^1 x(1)\, dx = \int_0^1 x\, dx = \frac{1}{2}$$

$$\boxed{E(X) = \frac{1}{2}}$$

$$E(Y) = \int_0^1 \int_x^{x+1} y\, f(X,Y)\, dy\, dx = \int_0^1 \int_x^{x+1} y\, dy\, dx$$

$$= \int_0^1 \left[\frac{y^2}{2}\right]_x^{x+1} dx = \int_0^1 \frac{(x+1)^2 - x^2}{2} dx$$

$$= \int_0^1 \frac{2x+1}{2}\, dx = \left[\frac{x^2}{2} + \frac{x}{2}\right]_0^1 = 1$$

$$\boxed{E(Y) = 1}$$

4.2.4 Compute $E(XY)$

$$E(XY) = \int_0^1 \int_x^{x+1} xy\, f(X,Y)\, dy\, dx = \int_0^1 \int_x^{x+1} xy\, dy\, dx$$

$$= \int_0^1 x \left[\frac{y^2}{2}\right]_x^{x+1} dx = \int_0^1 x \cdot \frac{(x+1)^2 - x^2}{2} dx$$

$$= \int_0^1 x \cdot \frac{2x+1}{2} dx = \int_0^1 \left(x^2 + \frac{x}{2}\right) dx$$

$$= \left[\frac{x^3}{3} + \frac{x^2}{4}\right]_0^1 = \frac{7}{12}$$

$$\boxed{E(XY) = \frac{7}{12}}$$

4.2.5 Variances

$$\mathrm{Var}(X) = E(X^2) - E(X)^2$$

$$E(X^2) = \int_0^1 \int_x^{x+1} x^2 \, dy \, dx = \int_0^1 x^2 \, dx = \frac{1}{3}$$

$$\text{Var}(X) = \frac{1}{3} - \left(\frac{1}{2}\right)^2 = \frac{1}{12}$$

$$\boxed{\text{Var}(X) = \frac{1}{12}}$$

$$\text{Var}(Y) = E(Y^2) - E(Y)^2$$

$$E(Y^2) = \int_0^1 \int_x^{x+1} y^2 \, dy \, dx = \int_0^1 \left[\frac{y^3}{3}\right]_x^{x+1} dx$$

$$= \int_0^1 \frac{(x+1)^3 - x^3}{3} dx = \frac{7}{6}$$

$$\text{Var}(Y) = \frac{7}{6} - 1 = \frac{1}{6}$$

$$\boxed{\text{Var}(Y) = \frac{1}{6}}$$

4.2.6 Variances (again but using $E[(X-\mu)^2]$

Variance of X:
Since $E(X) = \dfrac{1}{2}$,

$$\text{Var}(X) = E\big[(X - \tfrac{1}{2})^2\big] = \int_0^1 \int_x^{x+1} \left(x - \tfrac{1}{2}\right)^2 f(X,Y) \, dy \, dx$$

Because $f(X,Y) = 1$,

$$\text{Var}(X) = \int_0^1 \int_x^{x+1} \left(x - \tfrac{1}{2}\right)^2 \, dy \, dx$$

$$= \int_0^1 \left(x - \tfrac{1}{2}\right)^2 (1) \, dx = \int_0^1 \left(x - \tfrac{1}{2}\right)^2 dx$$

$$= \int_0^1 \left(x^2 - x + \tfrac{1}{4}\right) dx = \left[\frac{x^3}{3} - \frac{x^2}{2} + \frac{x}{4}\right]_0^1 = \frac{1}{12}$$

$$\boxed{\text{Var}(X) = \frac{1}{12}}$$

Variance of Y:

Since $E(Y) = 1$,

$$\mathrm{Var}(Y) = E\big[(Y-1)^2\big] = \int_0^1 \int_x^{x+1} (y-1)^2 f(X,Y)\, dy\, dx$$

$$= \int_0^1 \int_x^{x+1} (y-1)^2\, dy\, dx$$

$$= \int_0^1 \left[\frac{(y-1)^3}{3}\right]_{y=x}^{y=x+1} dx = \int_0^1 \frac{(x)^3 - (x-1)^3}{3} dx$$

$$= \int_0^1 \left(x^2 - x + \tfrac{1}{3}\right) dx = \left[\frac{x^3}{3} - \frac{x^2}{2} + \frac{x}{3}\right]_0^1 = \frac{1}{6}$$

$$\boxed{\mathrm{Var}(Y) = \frac{1}{6}}$$

4.2.7 Correlation

$$\rho_{XY} = \frac{E(XY) - E(X)E(Y)}{\sqrt{\mathrm{Var}(X)\mathrm{Var}(Y)}} = \frac{\frac{7}{12} - \frac{1}{2}}{\sqrt{\frac{1}{12} \cdot \frac{1}{6}}} = \frac{1}{\sqrt{2}}$$

$$\boxed{\rho_{XY} = \frac{1}{\sqrt{2}}}$$

4.2.8 Example

Question:

Let the joint PDF be

$$f(X,Y) = \begin{cases} 10, & 0 < x < 1,\ x < y < x + \dfrac{1}{10}, \\ 0, & \text{otherwise.} \end{cases}$$



Idea:

From the support $x < y < x + \frac{1}{10}$, subtract x:

$$0 < y - x < \tfrac{1}{10}.$$

Define

$$U = Y - X.$$

Then the support becomes

$$0 < X < 1, \qquad 0 < U < \tfrac{1}{10}.$$

So X and U are independent

Calculations:
Variance of a Uniform random variable:

$$\boxed{\mathrm{Var}(Z) = \frac{(b-a)^2}{12} \quad \text{for } Z \sim \mathrm{Uniform}(a, b)}$$

This follows from

$$E(Z) = \frac{a+b}{2}, \qquad E(Z^2) = \frac{a^2 + ab + b^2}{3},$$

and

$$\mathrm{Var}(Z) = E(Z^2) - [E(Z)]^2.$$

$$\mathrm{Var}(X) = \frac{(1-0)^2}{12} = \frac{1}{12}.$$

$$\mathrm{Var}(U) = \frac{(1/10 - 0)^2}{12} = \frac{1}{1200}.$$

Because $Y = X + U$ and X and U are independent,

$$\mathrm{Var}(Y) = \mathrm{Var}(X) + \mathrm{Var}(U) = \frac{1}{12} + \frac{1}{1200} = \frac{101}{1200}.$$

Covariance:

$$\mathrm{Cov}(X, Y) = \mathrm{Cov}(X, X + U) = \mathrm{Cov}(X, X) + \mathrm{Cov}(X, U).$$

Since $\mathrm{Cov}(X, U) = 0$,

$$\mathrm{Cov}(X, Y) = \mathrm{Var}(X) = \frac{1}{12}.$$

Correlation coefficient:

$$\rho_{X,Y} = \frac{\frac{1}{12}}{\sqrt{\left(\frac{1}{12}\right)\left(\frac{101}{1200}\right)}} = \sqrt{\frac{100}{101}}.$$

Comparison and interpretation:

In both examples, the support is a diagonal band of the form

$$x < y < x + w,$$

where w is the band width.

- A **smaller band width** means Y stays closer to the line $y = x$.
- This implies less variation in $Y - X$.
- Hence, Y is more tightly determined by X.

In the second example,

$$w = \frac{1}{10},$$

so $Y = X + U$ with $U \sim \mathrm{Unif}(0, \frac{1}{10})$ very small.

As a result,

$$\rho_{X,Y} = \sqrt{\frac{100}{101}} \approx 1,$$

indicating a very strong positive correlation.

Key idea: A thinner diagonal support band implies stronger linear dependence and higher correlation between X and Y.

Exercise 4.42

Problem: Using the results of Exercise 4.21, find the variance of $g(X) = X^2$, where X is a random variable having the density function given in Exercise 4.12.

Solution: From 4.21: $E(X^2) = 1/6$. Need $E(X^4) = \int_0^1 x^4 \cdot 2(1-x)\,dx = 2[1/5 - 1/6] = 2/30 = 1/15$. $\mathrm{Var}(X^2) = E(X^4) - [E(X^2)]^2 = 1/15 - 1/36 = (12-5)/180 = \boxed{7/180}$.

Exercise 4.58

Problem: question makes no sense because its suppose to be all positive for a PDF, but 4(1-x) while $1 \leq x < 2 is not possible, so we change the question to$:f(x) $= \begin{cases} 4x, & 0 < x < 1 \\ 4(2-x), & 1 \leq x < 2 \\ 0, & \text{else} \end{cases}$. $Find$ E(Y) $where$ Y $= 60X^2 + 48X$.

Solution: By linearity: $E(Y) = 60E(X^2) + 48E(X)$.

The density is triangular and symmetric about $x = 1$, so $E(X) = 1$.

$E(X^2) = \int_0^1 4x^3\,dx + \int_1^2 4x^2(2-x)\,dx$.

$\int_0^1 4x^3\,dx = \left[x^4\right]_0^1 = 1$.

$\int_1^2 4x^2(2-x)\,dx = 4\left[\frac{2x^3}{3} - \frac{x^4}{4}\right]_1^2 = 4\left[\left(\frac{16}{3} - 4\right) - \left(\frac{2}{3} - \frac{1}{4}\right)\right] = 4 \cdot \frac{11}{12} = \frac{11}{3}$.

$E(X^2) = 1 + \frac{11}{3} = \frac{14}{3}$.

$E(Y) = 60 \cdot \frac{14}{3} + 48(1) = 280 + 48 = \boxed{328}$.

Exercise 4.65

Problem: Let X represent the number that occurs when a red die is tossed and Y the number that occurs when a green die is tossed. Find (1) $E(X+Y)$; (2) $E(X-Y)$; (3) $E(4XY)$.

Solution: $E(X) = E(Y) = 7/2$.

(1) $E(X+Y) = 7/2 + 7/2 = \boxed{7}$.

(2) $E(X-Y) = 7/2 - 7/2 = \boxed{0}$.

(3) X, Y independent $\Rightarrow E(XY) = E(X)E(Y) = (7/2)^2 = 49/4$. $E(4XY) = 4 \times 49/4 = \boxed{49}$.

Midterm Q4(b) – Covariance (4 pts)

Problem: Find $\mathrm{Cov}(X, Y)$.

Solution: $E(X) = 0$ (symmetric). $E(Y) = 0(1/5) + 1(2/5) + 4(2/5) = 2$. $E(XY) = (-2)(4)(1/5) + (-1)(1)(1/5) + 0 + (1)(1)(1/5) + (2)(4)(1/5) = 0$. $\mathrm{Cov}(X, Y) = E(XY) - E(X)E(Y) = 0 - 0 = \boxed{0}$.

Midterm Q4(c) – Independence (2 pts)

Problem: Are X and Y independent?

Solution: **No.** $P(X = -2, Y = 0) = 0 \neq P(X = -2)P(Y = 0) = \frac{1}{5} \cdot \frac{1}{5} = \frac{1}{25}$. Zero covariance does not imply independence!

Midterm Q6(c) – Covariance (4 pts)

Problem: Find $\mathrm{Cov}(X, Y)$.

Solution: $E(XY) = \iint \frac{xy}{\pi a^2}\,dx\,dy = 0$ (inner integral of x over symmetric limits gives 0). $\mathrm{Cov}(X, Y) = E(XY) - E(X)E(Y) = 0 - 0 = \boxed{0}$.

Midterm Q6(d) – Independence (5 pts)

Problem: Are X and Y independent?

Solution: **No.** The joint support is a disk $x^2 + y^2 \leq a^2$, which is *not* a Cartesian product. Also $f_{X,Y}(x, y) \neq f_X(x)f_Y(y)$. Again, zero covariance $\not\Rightarrow$ independence.

Chapter 5

Chapter 5: Discrete Distributions

5.1 Discrete Uniform Distribution

A random variable X has a discrete uniform distribution if it assumes the values

$$x_1, x_2, \ldots, x_k$$

with equal probability.

$$f(X) = \begin{cases} \frac{1}{k}, & X = x_1, x_2, \ldots, x_k \\ 0, & \text{otherwise} \end{cases}$$

In general, k will be given.

Mean:

$$E(X) = \mu = \frac{1}{k} \sum_{i=1}^{k} x_i$$

Variance:

$$\boxed{\mathrm{Var}(X) = \sigma^2 = \frac{1}{k} \sum_{i=1}^{k} (x_i - \mu)^2}$$

Exercise 5.1

Problem: X is discrete uniform on $x_1, \ldots, x_k$ with $f(x) = 1/k$. Find mean and variance.
Solution: $\mu = \frac{1}{k} \sum_{i=1}^{k} x_i$. $\sigma^2 = \frac{1}{k} \sum_{i=1}^{k} x_i^2 - \mu^2$. For $x_i = 1, 2, \ldots, k$: $\mu = (k+1)/2$, $\sigma^2 = (k^2-1)/12$.

5.2 Bernoulli Distribution

A Bernoulli random variable has only two outcomes:

- success

- failure

(The definition of success is arbitrary. For example, in a cancer experiment, success could be defined as death.)

$$X = \begin{cases} 1, & \text{success} \\ 0, & \text{failure} \end{cases}$$

Let

$$p = P(X = 1), \qquad q = P(X = 0) = 1 - p$$

PMF table:

x	0	1
$f(x)$	q	p

Mean:

$$\mu = E(X) = 1 \cdot p + 0 \cdot q = p$$

Variance:

$$\sigma^2 = E(X^2) - \mu^2$$

$$= 0^2 q + 1^2 p - p^2 = p - p^2 = p(1-p) = pq$$

5.2.1 Example: Independent Bernoulli Trials

Four components are tested independently. Let X be the number of defectives.

$$x = 0, 1, 2, 3, 4$$

x	0	1	2	3	4
$P(X = x)$	$\binom{4}{0}p^0q^4$	$\binom{4}{1}p^1q^3$	$\binom{4}{2}p^2q^2$	$\binom{4}{3}p^3q$	$\binom{4}{4}p^4$

5.3 Binomial Distribution

The <u>binomial distribution</u> gives the number of successes in n independent Bernoulli trials.

$$X \sim \text{Binomial}(n, p)$$

$$\boxed{b(x; n, p) = f(X) = \binom{n}{x} p^x q^{n-x}, \qquad x = 0, 1, 2, \ldots, n}$$

5.3.1 Binomial Expansion Check

$$\binom{n}{0}p^0q^n + \binom{n}{1}p^1q^{n-1} + \cdots + \binom{n}{n}p^nq^0 = (p+q)^n$$

Since $p + q = 1$,

$$(p+q)^n = 1$$

5.3.2 Mean of a Binomial Random Variable

Reminder:

$$E(X_1 + X_2 + \cdots + X_n) = E(X_1) + E(X_2) + \cdots + E(X_n)$$

Let I_i be the indicator random variable for success on trial i:

$$I_i = \begin{cases} 1, & \text{success on trial } i \\ 0, & \text{failure on trial } i \end{cases}$$

Then

$$X = I_1 + I_2 + \cdots + I_n$$

$$\boxed{\mu = E(X) = nE(I_i) = np}$$

5.3.3 Variance of a Binomial Random Variable

Reminder:

$$\mathrm{Var}(aX + bY + c) = a^2\,\mathrm{Var}(X) + b^2\,\mathrm{Var}(Y) + 2ab\,\mathrm{Cov}(X, Y)$$

Since $I_1, \ldots, I_n$ are independent,

$$\mathrm{Cov}(I_i, I_j) = 0 \quad (i \neq j)$$

$$\mathrm{Var}(X) = \mathrm{Var}(I_1 + \cdots + I_n)$$

$$= \mathrm{Var}(I_1) + \cdots + \mathrm{Var}(I_n)$$

Each I_i is Bernoulli(p), so

$$\mathrm{Var}(I_i) = pq$$

$$\boxed{\mathrm{Var}(X) = npq}$$

5.3.4 Example: Testing a Manufacturer's Claim

A manufacturer claims that the defective rate is $p = 0.1$. We test $n = 4$ components independently and observe 3 defectives.

Let

$$X = \text{number of defectives in 4 trials}$$

Assume the claim is true, so $p = 0.1$.

$$X \sim \text{Binomial}(4, 0.1)$$

$$P(X = x) = \binom{4}{x}(0.1)^x(0.9)^{4-x}, \qquad x = 0, 1, 2, 3, 4$$

We compute the probability of observing at least 3 defectives:

$$P(X \geq 3) = P(X = 3) + P(X = 4)$$

$$= \binom{4}{3}(0.1)^3(0.9) + \binom{4}{4}(0.1)^4$$

$$= 0.0037$$

Since this probability is very small, observing 3 or more defectives would be very unlikely if the claim were true.

Therefore, we reject the claim.

Important note: The claim is either true or false; there is no probability attached to the claim itself. The probability calculation assumes the claim is true and measures how surprising the observed data would be under that assumption.

(We will study this idea formally in Chapter 9.)

Exercise 5.6

Problem: According to a survey by the Administrative Management Society, one-half of U.S. companies give employees 4 weeks of vacation after they have been with the company for 15 years. Find the probability that among 8 companies surveyed at random, the number that give employees 4 weeks of vacation after 15 years of employment is (1) anywhere from 3 to 5; (2) fewer than 2.

Solution: n=8, p = 0.5(one-half of the companies) $X \sim \text{Bin}(8, 0.5)$.

(1) $P(3 \leq X \leq 5) = \sum_{x=3}^{5} \binom{8}{x}(0.5)^8 = (56 + 70 + 56)/256 = 182/256 = \boxed{0.7109}$.

(2) $P(X < 2) = P(X = 0) + P(X = 1) = (1 + 8)/256 = 9/256 = \boxed{0.0352}$.

Exercise 5.9

Problem: In testing a certain kind of truck tire over rugged terrain, it is found that 25% of the trucks fail to complete the test run without a blowout. Of the next 15 trucks tested, find the probability that (1) from 3 to 6 have blowouts; (2) fewer than 4 have blowouts; (3) more than 5 have blowouts.
Solution: $X \sim \text{Bin}(15, 0.25)$.
(1) $P(3 \leq X \leq 6) = \sum_{x=3}^{6} \binom{15}{x}(0.25)^x(0.75)^{15-x} \approx 0.2252 + 0.2252 + 0.1651 + 0.0917 = \boxed{0.7072}$.
(2) $P(X < 4) = \sum_{x=0}^{3} \binom{15}{x}(0.25)^x(0.75)^{15-x} \approx 0.0134 + 0.0668 + 0.1559 + 0.2252 = \boxed{0.4613}$.
(3) $P(X > 5) = 1 - P(X \leq 5) \approx 1 - (0.4613 + 0.1651 + 0.0917) = 1 - 0.8516 = \boxed{0.1484}$.

Midterm Q2(b) – Expected Value of Binomial (6 pts)

Problem: Expected number of acceptable specimens in a sample of 10.
Solution: $Y \sim \text{Binomial}(n = 10, p = 0.6826)$. $E(Y) = np = 10 \times 0.6826 = \boxed{6.826}$.

Midterm Q2(c) – Binomial Probability (5 pts)

Problem: $P(\text{exactly 2 of 5 acceptable})$.
Solution: $Y \sim \text{Bin}(5, 0.6826)$. $P(Y = 2) = \binom{5}{2}(0.6826)^2(0.3174)^3 = 10 \times 0.4659 \times 0.03198 \approx \boxed{0.1490}$.

5.4 Multinomial Distribution

The <u>multinomial distribution</u> applies when each trial has more than two possible outcomes.

Definition: Suppose a single trial can result in k outcomes

$$E_1, E_2, \ldots, E_k$$

with probabilities

$$p_1, p_2, \ldots, p_k, \qquad \sum_{i=1}^{k} p_i = 1$$

Let

$$X_i = \text{number of occurrences of outcome } E_i$$

in n independent trials, for $i = 1, 2, \ldots, k$.

Then the joint probability distribution of

$$(X_1, X_2, \ldots, X_k)$$

is given by

$$\boxed{f(x_1, x_2, \ldots, x_k; p_1, \ldots, p_k, n) = \binom{n}{x_1, x_2, \ldots, x_k} p_1^{x_1} p_2^{x_2} \cdots p_k^{x_k}}$$

subject to the constraint

$$\sum_{i=1}^{k} x_i = n$$

Interpretation: This is like Bernoulli trials, but with more than two possible outcomes per trial.

5.4.1 Example: Travel Choice (Multinomial)

Suppose a person travels to school each day using one of three methods:

$$E_1 = \text{bus}, \quad E_2 = \text{train}, \quad E_3 = \text{car}$$

Assume the probabilities are

$$p_1 = P(\text{bus}), \quad p_2 = P(\text{train}), \quad p_3 = P(\text{car}), \qquad p_1 + p_2 + p_3 = 1$$

The person travels for n days independently.

Let

$$X_1 = \text{number of days taking the bus}$$

$$X_2 = \text{number of days taking the train}$$

$$X_3 = \text{number of days taking the car}$$

Then

$$(X_1, X_2, X_3) \sim \text{Multinomial}(n; p_1, p_2, p_3)$$

The joint probability is

$$P(X_1 = x_1, X_2 = x_2, X_3 = x_3) = \binom{n}{x_1, x_2, x_3} p_1^{x_1} p_2^{x_2} p_3^{x_3}$$

subject to

$$x_1 + x_2 + x_3 = n$$

This is similar to a binomial distribution, but instead of two outcomes per trial, there are three or more possible outcomes.

5.4.2 Example: Traffic Light (Multinomial)

As a student drives to school, he encounters a traffic light that is:

- green for 35 s
- yellow for 5 s
- red for 60 s

Assume each encounter is independent.

Define the random variables:

$$X_1 = \text{number of times the light is green}$$

$$X_2 = \text{number of times the light is yellow}$$

$$X_3 = \text{number of times the light is red}$$

Then

$$X_1 + X_2 + X_3 = n$$

The probabilities are:

$$p_1 = P(\text{green}) = 0.35$$

$$p_2 = P(\text{yellow}) = 0.05$$

$$p_3 = P(\text{red}) = 0.60$$

$$p_1 + p_2 + p_3 = 1$$

Suppose the student encounters the traffic light $n = 100$ times.

Then

$$(X_1, X_2, X_3) \sim \text{Multinomial}(100; 0.35, 0.05, 0.60)$$

We compute:

$$P(X_1 = 30, X_2 = 10, X_3 = 60)$$

$$= \binom{100}{30, 10, 60}(0.35)^{30}(0.05)^{10}(0.60)^{60}$$

$$= 0.00117$$

5.4.3 Multinomial Coefficient

The multinomial coefficient is defined as

$$\boxed{\binom{n}{x_1, x_2, \ldots, x_k} = \frac{n!}{x_1! x_2! \cdots x_k!}}$$

It can also be written as a product of binomial coefficients:

$$\boxed{\binom{n}{x_1, x_2, \ldots, x_k} = \binom{n}{x_1}\binom{n - x_1}{x_2} \cdots \binom{n - x_1 - \cdots - x_{k-1}}{x_k}}$$

1. <u>*Discrete uniform distribution*</u> A discrete random variable that takes k possible values with equal probability.

2. *Bernoulli distribution* A discrete random variable with two outcomes: success or failure.

3. *Binomial distribution* The number of successes in n independent Bernoulli trials.

4. *Multinomial distribution* A generalization of the binomial distribution with more than two outcomes per trial.

Exercise 5.22

Problem: According to a genetics theory, a certain cross of guinea pigs will result in red, black, and white offspring in the ratio 8:4:4. Find the probability that among 8 offspring, 5 will be red, 2 black, and 1 white.

Solution: Multinomial: $P = \frac{8!}{5!\,2!\,1!}(1/2)^5(1/4)^2(1/4)^1 = \frac{8!}{5!2!1!} \cdot \frac{1}{32} \cdot \frac{1}{16} \cdot \frac{1}{4} = 168 \times \frac{1}{2048} = \boxed{0.0820}$.

5.5 Hypergeometric Distribution

The hypergeometric distribution models **sampling without replacement**. Therefore, the trials are **dependent**.

Let:

- N = total number of items
- k = number of successes
- $N - k$ = number of failures
- n = sample size
- X = number of successes in the sample

$$\boxed{P(X = x) = \frac{\binom{k}{x}\binom{N-k}{n-x}}{\binom{N}{n}}, \qquad \max\{0, n - (N - k)\} \leq x \leq \min(n, k)}$$

5.5.1 Example

A hat contains 12 tickets:

- 7 black
- 5 white

Three tickets are drawn at random without replacement.

$$N = 12, \quad n = 3, \quad k = 7$$

$$P(X=2) = \frac{\binom{7}{2}\binom{5}{1}}{\binom{12}{3}}$$

$$P(X=x) = \frac{\binom{7}{x}\binom{5}{3-x}}{\binom{12}{3}}$$

5.5.2 Acceptance Sampling

Example:

A company inspects batches of compressors purchased from a vendor.

- Each batch contains $N = 15$ compressors.
- A batch is <u>unacceptable</u> if it contains **two or more faulty compressors**.

To inspect a batch:

- A random sample of $n = 5$ compressors is selected.
- All sampled compressors are tested.
- If **none are defective**, the batch is accepted.

We want to determine whether this is a good inspection plan.

Assume a bad batch:

Suppose the batch actually contains exactly $k = 2$ faulty compressors.

Random variable:

$$X = \text{number of faulty compressors in the sample of 5}$$

Since sampling is done <u>without replacement</u>, X follows a hypergeometric distribution:

$$X \sim \text{Hypergeometric}(N = 15,\ k = 2,\ n = 5)$$

- $N = 15$: total compressors in the batch
- $k = 2$: faulty compressors in the batch
- $n = 5$: compressors tested

Probability of accepting a bad batch:

The batch is accepted if no faulty compressors are found in the sample, i.e. $X = 0$.

$$P(X=0) = \frac{\binom{2}{0}\binom{13}{5}}{\binom{15}{5}} = 0.428$$

Interpretation:

Even when the batch is unacceptable (contains 2 faulty compressors), this inspection plan accepts the batch with probability 0.428 (42.8%).

Conclusion:

This inspection plan has a high chance of accepting bad batches and is therefore not a good plan.

5.5.3 Mean and Variance of the Hypergeometric Distribution

Let

- N = total number of items
- k = number of successes in the population
- n = sample size
- X = number of successes in the sample

Then the mean and variance of a hypergeometric random variable are:

$$\boxed{\mu = E(X) = n\frac{k}{N}}$$

$$\boxed{\sigma^2 = \mathrm{Var}(X) = n\frac{k}{N}\left(1 - \frac{k}{N}\right)\frac{N-n}{N-1}}$$

The extra factor

$$\frac{N-n}{N-1}$$

is called the finite population correction factor and appears because sampling is done without replacement.

5.5.4 Relation to the Binomial Distribution

If the sample size n is much smaller than the population size N, then removing one item changes the population very little.

In this case, a binomial distribution can be used to approximate the hypergeometric distribution with

$$\boxed{p = \frac{k}{N}}$$

Rule of thumb:

$$\frac{n}{N} \ll 0.05$$

5.5.5 Example: Binomial Approximation to Hypergeometric

A tire manufacturer reports that among a shipment of $N = 5000$ tires sent to a distributor,

$$k = 1000$$

are slightly blemished.

A customer purchases $n = 10$ tires at random.

Let

$$X = \text{number of blemished tires in the sample}$$

Exact (hypergeometric) probability:

$$P(X = 3) = \frac{\binom{1000}{3}\binom{4000}{7}}{\binom{5000}{10}} = 0.2014$$

Binomial approximation:

Since

$$\frac{n}{N} = \frac{10}{5000} = 0.002 \ll 0.05$$

the binomial approximation is appropriate.
reminder: p is the probability that a randomly selected item from the population is a success

$$p = \frac{1000}{5000} = 0.2$$

$$P(X = 3) = \binom{10}{3}(0.2)^3(0.8)^7 = 0.2013$$

Conclusion:

The binomial approximation gives nearly the same result as the exact hypergeometric probability and is much easier to compute.

Exercise 5.32

Problem: From a lot of 9 missiles, 3 are selected at random and fired. If the lot contains 4 defective missiles that will not fire, what is the probability that (1) all 3 will fire? (2) at most 2 will not fire?
Solution: $X = \#$ defective, $X \sim \text{Hyper}(N = 9, K = 4, n = 3)$.
(1) $P(X = 0) = \frac{\binom{4}{0}\binom{5}{3}}{\binom{9}{3}} = \frac{10}{84} = \boxed{5/42}$.
(2) $P(X \leq 2) = 1 - P(X = 3) = 1 - \frac{\binom{4}{3}\binom{5}{0}}{\binom{9}{3}} = 1 - \frac{4}{84} = \boxed{80/84 = 20/21}$.

Exercise 5.34

Problem: What is the probability that a waitress will refuse to serve alcoholic beverages to only 2 minors if she randomly checks the IDs of 5 among 9 students, 4 of whom are minors? **Solution:**

$X \sim \text{Hyper}(N = 9, K = 4, n = 5)$. $P(X = 2) = \frac{\binom{4}{2}\binom{5}{3}}{\binom{9}{5}} = \frac{6\times 10}{126} = \boxed{60/126 = 10/21}$.

Exercise 5.36

Problem: A manufacturing company uses an acceptance scheme on items from a production line before they are shipped. The plan is a two-stage one. Boxes of 25 items are readied for shipment, and a sample of 3 items is tested for defectives. If any defectives are found, the entire box is sent back for 100% screening. If no defectives are found, the box is shipped. 1) What is the probability that a box containing 3 defectives will be shipped? 2) What is the probability that a box containing only 1 defective will be sent back for screening?

Solution: (1) $P(X = 0) = \frac{\binom{3}{0}\binom{22}{3}}{\binom{25}{3}} = \frac{1540}{2300} = \boxed{0.6696}$.

(2) $P(\text{back}) = P(X \geq 1) = 1 - P(X = 0) = 1 - \frac{\binom{1}{0}\binom{24}{3}}{\binom{25}{3}} = 1 - \frac{2024}{2300} = 1 - 0.88 = \boxed{0.12}$.

Exercise 5.37

Problem: Suppose that the manufacturing company of Exercise 5.36 decides to change its acceptance scheme. Under the new scheme, an inspector takes 1 item at random, inspects it, and then replaces it in the box; a second inspector does likewise. Finally, a third inspector goes through the same procedure. The box is not shipped if any of the three inspectors find a defective. Answer the questions in Exercise 5.36 for this new plan

Solution: With replacement: $p = K/25$.

(1) $p = 3/25 = 0.12$. $P(\text{no def found}) = (1 - 0.12)^3 = (0.88)^3 = \boxed{0.6815}$.

(2) $p = 1/25 = 0.04$. $P(\text{back}) = 1 - (0.96)^3 = 1 - 0.8847 = \boxed{0.1153}$.

Exercise 5.38

Problem: Among 150 IRS employees in a large city, only 30 are women. If 20 of the employees are chosen at random to provide free tax assistance for the residents of this city, use the binomial approximation to the hypergeometric distribution to find the probability that at least 2 women are selected. **Solution:** $p = 30/150 = 0.2$, $n = 20$. Binomial approx: $X \sim \text{Bin}(20, 0.2)$. $P(X \geq 2) = 1 - P(X = 0) - P(X = 1) = 1 - (0.8)^{20} - 20(0.2)(0.8)^{19}$. $= 1 - 0.01153 - 0.05765 = \boxed{0.9308}$.

Exercise 5.95

Problem: A production process outputs items in lots of 50. The current inspection plan is to periodically sample randomly 10 out of the 50 items in a lot and, if none are defective, to perform no intervention. (1) Suppose in a lot chosen at random, 2 out of 50 are defective. What is the probability that at least 1 in the sample of 10 from the lot is defective? (2) Comment on the quality of this sampling plan. (3) What is the mean number of defects found out of 10 items sampled? from the lot is defective? 2 def in lot).(2)$Comment$.(3)$Meandefectsinsample$.

Solution: (1) $X \sim \text{Hyper}(N = 50, K = 2, n = 10)$. $P(X \geq 1) = 1 - P(X = 0) = 1 - \frac{\binom{2}{0}\binom{48}{10}}{\binom{50}{10}} = 1 - \frac{48!\cdot 40!}{38!\cdot 50!} \cdot \frac{50!}{10!40!}$. $P(X = 0) = \frac{\binom{48}{10}}{\binom{50}{10}} = \frac{48!\,40!}{38!\,50!} = \frac{39\times 40}{49\times 50} = \boxed{0.6367}$. $P(X \geq 1) = 1 - 0.6367 = \boxed{0.3633}$.

(2) The plan is poor: only about 36% chance of detecting a defective lot with 2 defectives.
(3) $E(X) = n \cdot K/N = 10(2)/50 = \boxed{0.4}$.

5.6 Negative Binomial Distribution

The negative binomial distribution models the number of trials required for the k**th success** to occur in a sequence of independent Bernoulli trials.

Each trial has:

$$P(\text{success}) = p, \qquad P(\text{failure}) = q = 1 - p$$

5.6.1 Definition

Let

$$X = \text{trial on which the } k\text{th success occurs}$$

Then the PMF of X is

$$\boxed{P(X = x) = \binom{x-1}{k-1} p^k q^{x-k}, \qquad x = k, k+1, k+2, \ldots}$$

Explanation of the formula:

- The kth success must occur on trial x
- In the first $x - 1$ trials, there must be exactly $k - 1$ successes
- Trial x must be a success

5.6.2 Mean and Variance

$$\boxed{E(X) = \mu = \frac{k}{p}}$$

$$\boxed{\text{Var}(X) = \frac{k(1-p)}{p^2}}$$

5.6.3 Interpretation (intuition)

Application (disease screening):

Suppose people are tested one by one for a disease. Each person independently tests positive with probability p.

Testing continues until k infected individuals have been identified. Let

$$X = \text{number of people tested until the } k\text{th infected person is found}$$

Then X follows a negative binomial distribution.

This framework explains why the negative binomial distribution is useful in applications such as disease screening during the pandemic.

5.6.4 Example

Electronic components are tested independently. Each component is defective with probability

$$p = 0.1$$

Let

$$X = \text{trial on which the 3rd defective component is found}$$

Probability that the 3rd defective occurs on the 4th trial:

$$P(X = 4) = \binom{3}{2}(0.1)^3(0.9)^1$$

Probability that the 3rd defective occurs on the 5th trial:

$$P(X = 5) = \binom{4}{2}(0.1)^3(0.9)^2$$

5.6.5 PMF of the Negative Binomial Distribution

Let X be the trial on which the kth success occurs in a sequence of independent Bernoulli trials. Each trial has success probability p and failure probability $q = 1 - p$.

Then the probability mass function of X is

$$\boxed{b^*(x; k, p) = f(X) = \binom{x-1}{k-1} p^k q^{x-k}, \qquad x = k, k+1, k+2, \ldots}$$

5.6.6 Mean and Variance

$$\boxed{\mu = E(X) = \frac{k}{p}}$$

$$\boxed{\mathrm{Var}(X) = \frac{k(1-p)}{p^2}}$$

5.6.7 Interpretation of the Mean

The mean $\mu = k/p$ represents the average number of trials required to obtain k successes.

For example, in a disease-screening context:

- k is the number of infected individuals you want to identify
- p is the probability that a randomly tested person is infected

On average, $\frac{k}{p}$ people must be tested to find k infected individuals.

Exercise 5.49

Problem: The probability that a person living in a certain city owns a dog is estimated to be 0.4. Find the probability that the seventh person randomly interviewed in that city is the fifth one to own a dog.

Solution: Negative Binomial: $X \sim \text{NegBin}(k = 5, p = 0.4)$. $P(X = 7) = \binom{6}{4}(0.4)^5(0.6)^2 = 15 \times 0.01024 \times 0.36 = \boxed{0.0553}$.

Midterm Q7(b) – Negative Binomial (5 pts)

Problem: $P(\text{3rd activation on 12th person})$.

Solution: $Y \sim \text{NegBin}(k = 3, p = 0.05)$. $P(Y = 12) = \binom{11}{2}(0.05)^3(0.95)^9 = 55 \times 0.000125 \times 0.6302 \approx \boxed{0.0043}$.

Note: The official solution states ≈ 0.0299 but this appears to use a slightly different calculation.

5.7 Geometric Distribution

The geometric distribution models the number of trials required for the **first success** to occur in a sequence of independent Bernoulli trials.

Let

$$X = \text{trial on which the first success occurs}$$

Each trial has

$$P(\text{success}) = p, \qquad P(\text{failure}) = q = 1 - p$$

5.7.1 PMF

$$f(X) = g(x; p) = p\,q^{\,x-1}, \qquad x = 1, 2, 3, \ldots$$

5.7.2 Mean and Variance

$$\boxed{E(X) = \mu = \frac{1}{p}}$$

$$\boxed{\mathrm{Var}(X) = \frac{1-p}{p^2}}$$

5.7.3 Relation to the Negative Binomial

The geometric distribution is a special case of the negative binomial distribution with

$$k = 1$$

That is, it counts the waiting time until the **first** success.

Memoryless Property

Let X follow a geometric distribution. For integers $s > t$,

$$P(X > s \mid X > t) = P(X > s - t)$$

This property is called the memoryless property.

5.7.4 Interpretation

In gambling, the probability of losing or winning the next game does not depend on the fact that you have already lost five games in a row.

Each game is independent, which is why the geometric distribution is memoryless.

5.7.5 Proof

For a geometric random variable,

$$P(X > n) = (1-p)^n$$

Then

$$P(X > s \mid X > t) = \frac{P(X > s \cap X > t)}{P(X > t)} = \frac{P(X > s)}{P(X > t)}$$

$$= \frac{(1-p)^s}{(1-p)^t} = (1-p)^{s-t} = P(X > s - t)$$

Review

1. *Discrete uniform distribution* A discrete random variable that takes k possible values with equal probability.

2. *Bernoulli distribution* A discrete random variable with two outcomes: success or failure.

3. *Binomial distribution* The number of successes in n independent Bernoulli trials.

4. *Multinomial distribution* A generalization of the binomial distribution with more than two outcomes per trial.

5. *Hypergeometric distribution* The number of successes in a sample of size n drawn without replacement from a population of size N containing k successes.

6. *Negative binomial distribution* The number of trials required to obtain a fixed number of successes.

7. *Geometric distribution* The number of trials required to obtain the first success.

Midterm Q7(a) – Geometric Distribution (5 pts)

Problem: 5% activate metal detector. Find $P(10 \text{ pass}, 11\text{th activates})$.
Solution: $X \sim \text{Geometric}(p = 0.05)$. $P(X = 11) = (0.95)^{10}(0.05) \approx \boxed{0.0299}$.

5.8 Poisson Process

Poisson process The basic form is continuous-time counting process $\{N(t),\, t \geq 0\}$ that counts the number of events occurring up to time t.

Basic properties

1. $N(0) = 0$
   Zero arrivals at time 0.

2. *Independent increments*
   The number of events occurring in one time interval is independent of the number occurring in any disjoint time interval.

3. *Stationary increments*
   The probability distribution of the number of events in a time interval depends only on the length of the interval, not on its location.

4. *No simultaneous occurrences*
   The probability that two or more events occur in a very small interval is negligible.

Poisson Approximation to the Binomial

If

$$X \sim \text{Binomial}(n, p),$$

and

$$n \to \infty, \qquad p \to 0, \qquad np \to \mu \text{ (constant)},$$

then

$$\text{Binomial}(n, p) \longrightarrow \text{Poisson}(\mu).$$

Equivalently,

$$b(x; n, p) \longrightarrow P(x; \mu) = \frac{e^{-\mu}\mu^x}{x!}.$$

We can use the *Poisson distribution* to approximate a *Binomial distribution* when modeling **rare occurrences**, that is, when n is large and p is small while np remains

Rule of thumb

The approximation is good if

$$n \geq 20 \text{ and } p \leq 0.05,$$

or

$$n \geq 100 \text{ and } np \leq 10.$$

Example

On average, 1 in every 1000 glass productions are blemished. Find the probability that a random sample of 8000 items yields fewer than 7 defective items.

Let

$$X = \text{number of defective items in the sample.}$$

Then

$$X \sim b(x; 8000, 0.001),$$

where

$$b(x; n, p) = \binom{n}{x} p^x (1-p)^{n-x}$$

denotes the *Binomial pmf*.

We want

$$P(X < 7) = P(X \leq 6) = \sum_{x=0}^{6} \binom{8000}{x} (0.001)^x (0.999)^{8000-x}.$$

$$= 0.3132.$$

R command (exact binomial):

$$\texttt{sum(dbinom(0:6, 8000, 0.001))} = 0.3132.$$

Poisson approximation

Since

$$np = 8000(0.001) = 8,$$

we approximate

$$X \sim P(x; 8),$$

where

$$P(x; \mu) = \frac{e^{-\mu}\mu^x}{x!}$$

denotes the *Poisson pmf*.

$$P(X \leq 6) = \sum_{x=0}^{6} \frac{e^{-8}8^x}{x!} = 0.3134.$$

R command (Poisson approximation):

$$\texttt{sum(dpois(0:6, 8))} = 0.3134.$$

Notation Notes

- $X \sim b(x; n, p)$ means X follows a Binomial distribution with n trials and success probability p.
- $b(x; n, p)$ represents the Binomial pmf evaluated at x.
- $X \sim P(x; \mu)$ means X follows a Poisson distribution with mean μ.
- $P(x; \mu)$ represents the Poisson pmf evaluated at x.
- `dbinom(x, n, p)` computes $P(X = x)$ for a Binomial distribution in R.
- `dpois(x,` μ`)` computes $P(X = x)$ for a Poisson distribution in R.

Exercise 5.56

Problem: On average, 3 traffic accidents per month occur at a certain intersection. What is the probability that in any given month at this intersection (1) $P(X = 5)$. (2) $P(X < 3)$. (3) $P(X \geq 2)$.
Solution: $X \sim \text{Poisson}(\lambda = 3)$.
(1) $P(X = 5) = \frac{e^{-3}3^5}{5!} = \frac{0.0498 \times 243}{120} = \boxed{0.1008}$.
(2) $P(X < 3) = P(0) + P(1) + P(2) = e^{-3}(1 + 3 + 4.5) = 0.0498(8.5) = \boxed{0.4232}$.
(3) $P(X \geq 2) = 1 - P(0) - P(1) = 1 - 0.0498 - 0.1494 = \boxed{0.8009}$.

Exercise 5.63

Problem: Find the mean and variance of the random variable X in Exercise 5.58, representing the number of hurricanes per year to hit a certain area of the eastern United States (which is, on average, hit by 6 hurricanes a year, i.e. $\lambda = 6$).
Solution: $X \sim \text{Poisson}(\lambda = 6)$. $E(X) = \boxed{6}$, $\text{Var}(X) = \boxed{6}$.

Exercise 5.66

Problem: Small aircraft arrive Poisson at rate 6/hour. (1) $P(X = 4)$ in 1 hour. (2) $P(X \geq 4)$ in 1 hour. (3) $P(X \geq 75)$ in 12 hours.
Solution: (1) $\lambda = 6$. $P(X = 4) = \frac{e^{-6}6^4}{4!} = \frac{0.002479 \times 1296}{24} = \boxed{0.1339}$.
(2) $P(X \geq 4) = 1 - P(X \leq 3) = 1 - e^{-6}(1 + 6 + 18 + 36) = 1 - 0.002479(61) = 1 - 0.1512 = \boxed{0.8488}$.
(3) $\lambda = 72$ for 12 hours. Use normal approximation: $X \approx N(72, 72)$. $P(X \geq 75) = P(Z \geq \frac{74.5-72}{\sqrt{72}}) = P(Z \geq 0.295) \approx \boxed{0.384}$.

Midterm Q3(a) – Poisson & Total Probability (8 pts)

Problem: Factory I: defects $\sim$ Poisson(0.25). Factory II: $\sim$ Poisson(0.5). Equal production. Find $P(0 \text{ defects})$.
Solution: $P(0 \mid I) = e^{-0.25} \approx 0.7788$. $P(0 \mid II) = e^{-0.5} \approx 0.6065$. $P(0) = 0.7788 \times 0.5 + 0.6065 \times 0.5 = \boxed{0.6927}$.

Midterm Q3(b) – Bayes' Theorem (8 pts)

Problem: Given 0 defects, find $P(\text{Factory II})$.
Solution: $P(II \mid 0) = \dfrac{0.6065 \times 0.5}{0.6927} = \boxed{0.4378}$.

Midterm Q5 – Bayes with Binomial (10 pts)

Problem: Machine: $P(\text{adjusted}) = 0.9$, $P(\text{unadjusted}) = 0.1$. Adjusted: $P(H) = 0.5$. Unadjusted: $P(H) = 0.25$. Of 5 items, 4 high quality, 1 medium. Find $P(\text{adjusted} \mid 4H, 1M)$.
Solution: $P(4H, 1M \mid \text{adj}) = \binom{5}{4}(0.5)^4(0.5) = 5/32$. $P(4H, 1M \mid \text{unadj}) = \binom{5}{4}(0.25)^4(0.75) = 5 \times 0.003906 \times 0.75 = 0.01465$.
$P(\text{adj} \mid 4H, 1M) = \dfrac{(5/32)(0.9)}{(5/32)(0.9) + (0.01465)(0.1)} = \dfrac{0.1406}{0.1406 + 0.001465} = \boxed{0.9897}$.

Chapter 6

Chapter 6: Continuous Probability Distributions

6.1 Uniform Distribution

Uniform random variable

A continuous random variable X is said to be uniformly distributed on the interval $[A, B]$ if its probability density function is

$$f(x; A, B) = \begin{cases} \dfrac{1}{B-A}, & A \leq x \leq B, \\ 0, & \text{otherwise.} \end{cases}$$



6.2 Normal Distribution

Normal random variable

A continuous random variable X is said to be normally distributed with mean μ and variance σ^2 if its probability density function is

$$n(x;\mu,\sigma) = f(x) = \frac{1}{\sqrt{2\pi}\sigma}\exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right), \qquad -\infty < x < \infty.$$

Graph 1: The Empirical Rule (68-95-99.7 Rule)



Explanation: This graph shows the **Empirical Rule** for normal distributions:

- **68%** of data falls within $\mu \pm 1\sigma$ (one standard deviation from the mean)
- **95%** of data falls within $\mu \pm 2\sigma$ (two standard deviations from the mean)
- **99.7%** of data falls within $\mu \pm 3\sigma$ (three standard deviations from the mean)

This is also called the 68-95-99.7 rule and helps us understand how data is distributed around the mean.

Graph 2: Comparing Different Standard Deviations



Explanation: This graph compares three normal distributions with the same mean ($\mu = 0$) but different standard deviations:

- **Red curve** ($\sigma = 0.5$): Small standard deviation $\rightarrow$ data is tightly concentrated around the mean $\rightarrow$ tall, narrow peak
- **Blue curve** ($\sigma = 1$): Medium standard deviation $\rightarrow$ moderate spread
- **Green curve** ($\sigma = 2$): Large standard deviation $\rightarrow$ data is more spread out $\rightarrow$ short, wide curve

Key insight: A smaller σ means less variability (more consistent data), while a larger σ means more variability (data is more spread out). All three curves are centered at the same mean, but differ in their spread.

Midterm Q2(a) – Normal Distribution (6 pts)

Problem: Rockwell hardness $X \sim N(70, 9)$. Acceptable if $67 < X < 73$. Find $P(67 < X < 73)$.
Solution: $Z = (X - 70)/3$. $P(67 < X < 73) = P(-1 < Z < 1) = 1 - 2(0.1587) = \boxed{0.6826}$.

Midterm Q2(d) – Inverse Normal (5 pts)

Problem: Find c such that $P(70 - c < X < 70 + c) = 0.95$.
Solution: $P(X < 70 - c) = 0.025 \Rightarrow P(Z < -c/3) = 0.025 \Rightarrow -c/3 = -1.96 \Rightarrow \boxed{c = 5.88}$.

6.3 Standardization (Z-Scores)

Notation first: When we write $X \sim n(x;\, \mu, \sigma)$, this means:

- X is a **random variable** — a variable whose value is the outcome of a random process (e.g. a randomly selected person's height).
- $\sim$ means "is distributed as" — i.e. X follows the distribution on the right.
- $n(\,\cdot\,;\, \mu, \sigma)$ is the **normal distribution** function. The semicolon separates the variable from the parameters.
- μ is the **mean** (centre) of the distribution.
- σ is the **standard deviation** — a measure of how spread out the distribution is. σ^2 is the **variance**.
- x is just the placeholder variable (the horizontal axis value).

So $X \sim n(x;\, \mu, \sigma)$ is read: *"X follows a normal distribution with mean μ and standard deviation σ."*

Standardization

If $X \sim N(\mu,\, \sigma^2)$, we transform X into a new random variable Z by subtracting the mean and dividing by the standard deviation:

$$Z = \frac{X - \mu}{\sigma}.$$

- Subtracting μ **shifts** the distribution so it is centred at 0.

- Dividing by σ **rescales** it so the spread becomes exactly 1.

The result is $Z \sim n(z;\, 0,\, 1)$, called the **standard normal distribution** — mean 0, standard deviation 1. Rearranging gives the inverse: $X = \mu + Z\sigma$ (useful for finding x from a known z).

Why do we standardize? Every normal distribution $N(\mu, \sigma^2)$ has a different mean and spread, so we would need a separate probability table for each one. By converting to $Z \sim N(0, 1)$, we only need *one* table (the standard normal table) for all normal distributions.

Visualizing Standardization — Equal Areas

The key insight is that probabilities (areas) are preserved under standardization. Points x_1, x_2 on the X-axis correspond to z_1, z_2 on the Z-axis, and the shaded area is identical in both curves.



The transformation $Z = \dfrac{X - \mu}{\sigma}$ maps $x_1 \mapsto z_1$ and $x_2 \mapsto z_2$, and the shaded areas (probabilities) are exactly the same in both plots. This is why we can compute any normal probability using just the standard normal table.

<u>Def:</u> The **z-score** tells us *how many standard deviations* the original observation x is from the mean μ:

$$z = \frac{x - \mu}{\sigma} \qquad \Longleftrightarrow \qquad x = \mu + z\sigma.$$

A positive z means x is *above* the mean; a negative z means x is *below* the mean.

Exercise 6.5

Problem: Given a standard normal distribution, find the area under the curve that lies: (1) to the left of $z = -1.10$; (2) to the right of $z = 1.645$; (3) between $z = -2.43$ and -0.45; (4) to the left of $z = 0.45$; (5) to the right of $z = -0.45$; (6) between $z = -0.45$ and $z = 0.45$.

Solution: (1) $\Phi(-1.10) = \boxed{0.1357}$. (2) $1 - \Phi(1.645) \approx \boxed{0.05}$. (3) $\Phi(-0.45) - \Phi(-2.43) = 0.3264 - 0.0075 = \boxed{0.3189}$. (4) $\Phi(0.45) = \boxed{0.6736}$. (5) $1 - \Phi(-0.45) = \boxed{0.6736}$. (6) $\Phi(0.45) - \Phi(-0.45) = 0.6736 - 0.3264 = \boxed{0.3472}$.

Exercise 6.9

Problem: Given the normally distributed variable X with mean $\mu = 20$ and standard deviation $\sigma = 2$ (i.e. $X \sim N(20, \sigma^2 = 4)$). Find: (1) $P(X < 16)$. (2) k: $P(X < k) = 0.4090$. (3) k: $P(X > k) = 0.8599$. (4) $P(17 < X < 22)$.

Solution: (1) $Z = (16 - 20)/2 = -2$. $P(Z < -2) = \boxed{0.0228}$. (2) $P(Z < (k - 20)/2) = 0.4090 \Rightarrow (k - 20)/2 = -0.23 \Rightarrow k = \boxed{19.54}$. (3) $P(X > k) = 0.8599 \Rightarrow P(X < k) = 0.1401 \Rightarrow (k - 20)/2 = -1.08 \Rightarrow k = \boxed{17.84}$. (4) $P(-1.5 < Z < 1) = \Phi(1) - \Phi(-1.5) = 0.8413 - 0.0668 = \boxed{0.7745}$.

Exercise 6.22

Problem: If a set of observations is normally distributed, what percent of these differ from the mean by: (1) more than 1.50σ? (2) less than 0.75σ?

Solution: (1) $P(|Z| > 1.5) = 2(1 - \Phi(1.5)) = 2(0.0668) = \boxed{13.36\%}$. (2) $P(|Z| < 0.75) = 2\Phi(0.75) - 1 = 2(0.7734) - 1 = \boxed{54.68\%}$.

6.4 Computing Normal Probabilities

Example: Finding x Given a Probability (Inverse Normal)

Suppose $X \sim N(\mu = 40, \sigma = 6)$. Find x such that $P(X < x) = 0.45$.

Step 1: Standardize.

$$P(X < x) = P\left(Z < \frac{x - 40}{6}\right) = 0.45.$$

Step 2: Look up z such that $P(Z < z) = 0.45$ from the standard normal table (Table A.3).

From the table: $P(Z < -0.13) = 0.4483$ and $P(Z < -0.12) = 0.4522$. Since 0.45 lies between these two values, we interpolate:

$$z \approx -0.13 + \frac{0.45 - 0.4483}{0.4522 - 0.4483} \times 0.01 = -0.13 + \frac{0.0017}{0.0039} \times 0.01 \approx -0.13 + 0.0044 \approx -0.126.$$

The board approximated this as $z \approx -0.13$.

Step 3: Solve for x using $x = \mu + z\sigma$:

$$x = 40 + (-0.13)(6) = 40 - 0.78 = \boxed{39.22}.$$

Interpretation: 45% of the distribution lies below $x = 39.22$, which is slightly below the mean of 40 — this makes sense since 0.45 is slightly less than 0.5.

Using the standard normal table, we look up $P(Z < z)$ — the area to the *left* of z.

- **Left tail:** $P(Z < z)$ read directly from table.
- **Right tail:** $P(Z > z) = 1 - P(Z < z)$ (complement rule)
- **Symmetry:** $P(Z > z) = P(Z < -z)$ (curve is symmetric about 0)

- **Between two values:** $P(a < Z < b) = P(Z < b) - P(Z < a)$

Examples:

$$P(Z < -2.81) = 0.0025$$

$$P(Z < 1.4) = 0.9192$$

$$P(Z > 1.4) = 1 - P(Z < 1.4) = 1 - 0.9192 = 0.0808$$

$$P(Z < -1.4) = P(Z > 1.4) = 0.0808 \qquad \text{(by symmetry)}$$

$$P(0.43 < Z < 2.15) = P(Z < 2.15) - P(Z < 0.43) = 0.9798 - 0.6664 = 0.3134$$

6.5 The Empirical Rule (68–95–99.7 Rule)

$$P(-1 < Z < 1) = 0.6826 \qquad \Longleftrightarrow \qquad P(\mu - \sigma < X < \mu + \sigma) = 0.6826$$

$$P(-2 < Z < 2) = 0.9544 \qquad \Longleftrightarrow \qquad P(\mu - 2\sigma < X < \mu + 2\sigma) = 0.9544$$

$$P(-3 < Z < 3) = 0.9974 \qquad \Longleftrightarrow \qquad P(\mu - 3\sigma < X < \mu + 3\sigma) = 0.9974$$

6.6 Percentiles of the Normal Distribution

<u>Def:</u> If $P(Z < z) = p$, then z is the $p^{\mathbf{th}}$ **percentile** of the standard normal distribution.

Similarly, if $P(X < x) = p$, then x is the $p^{\mathbf{th}}$ **percentile** of the random variable X.

i.e. the pth percentile is the value below which a proportion p of the distribution lies.

Example: Oil Filter Service Time

"An oil filter and oil change in __ minutes or the next one's free." Service time: $X \sim N(\mu = 15, \sigma = 2.5)$ min.

Question: What time x should be advertised if at most 5 free services per 100 customers is acceptable?

We want x such that 95% of services finish within x minutes:

$$P(X < x) = 1 - \frac{5}{100} = 0.95.$$

Step 1: Find the z-value for $p = 0.95$:

$$P(Z < z) = 0.95 \quad \Rightarrow \quad z \approx 1.65.$$

Step 2: Convert back using $x = \mu + z\sigma$:

$$x = 15 + 1.65 \times 2.5 = 19.125 \approx \boxed{20 \text{ min}}.$$

6.7 Normal Approximation to the Binomial

Normal Approximation to the Binomial

If X is a binomial r.v. with parameters n and p (mean $\mu = np$, variance $\sigma^2 = npq$, $q = 1 - p$), then as $n \to \infty$ the distribution of

$$Z = \frac{X - np}{\sqrt{npq}}$$

approaches the standard normal $N(0, 1)$. For large n:

$$X \sim \text{Binom}(n, p) \approx Y \sim N(np,\ npq)\,.$$

- Very accurate when n is large and p is not very small or very large.
- Fairly good even when n is small, provided p is close to 0.5.



The bars show the exact binomial probabilities; the smooth red curve shows the normal approximation $N(np,\, npq)$. Notice how the continuous curve closely approximates the discrete bars — this is why the normal approximation works.

6.7.1 Continuity Correction

Continuity Correction

Because the binomial is **discrete** and the normal is **continuous**, each integer k is approximated by the interval $[k-0.5,\ k+0.5]$:

$$P(X=k) \approx P(k-0.5 < Y < k+0.5).$$

For cumulative probabilities:

$$P(X \le x) = \sum_{k=0}^{x} b(k;\, n, p) \ \approx\ P\left(Z \le \frac{x+0.5-np}{\sqrt{npq}}\right).$$

The $+0.5$ is the **continuity correction** — it accounts for the width of each discrete bar.

6.7.2 Example: Children Living in Poverty

21% of children under age six live in households below the poverty level. A study selects a simple random sample of 300 children.

Let X = number of children in the sample who live in poverty. $X \sim \text{Binom}(n=300,\ p=0.21)$.

(a) Mean and standard deviation:

$$\mu = np = 300 \times 0.21 = 63, \qquad \sigma = \sqrt{npq} = \sqrt{300 \times 0.21 \times 0.79} \approx 7.055.$$

(b) Find $P(X \ge 80)$:

$$P(X \ge 80) = 1 - P(X \le 79) \approx 1 - P\left(Z \le \frac{79+0.5-63}{7.055}\right) = 1 - P(Z \le 2.34) \approx 0.01.$$

Feb23

Exercise 6.27

Problem: In a city, 4% of the adolescents are alcoholic. Out of 100 adolescents randomly selected, what is the probability that (1) between 8 and 18 of them are alcoholics? (2) fewer than 5 are alcoholics?

Solution: $\mu = 4$, $\sigma = \sqrt{3.84} = 1.96$. Normal approx with continuity correction:

(1) $P(7.5 < X < 18.5) : Z_1 = (7.5-4)/1.96 = 1.79$, $Z_2 = (18.5-4)/1.96 = 7.40$. $P \approx 1 - \Phi(1.79) = 1 - 0.9633 = \boxed{0.0367}$.

(2) $P(X < 4.5) : Z = (4.5-4)/1.96 = 0.255$. $P \approx \Phi(0.255) \approx \boxed{0.6006}$.

Exercise 6.36

Problem: A common practice of airline companies is to sell more tickets for a particular flight than there are seats on the plane, because customers who buy tickets do not always show up for the flight. Suppose that the percentage of no-shows at flight time is 2%. For a particular flight with 197 seats, a total of 200 tickets were sold. What is the probability that the airline overbooked this flight?

Solution: $X = \#$ who show up. $X \sim \text{Bin}(200, 0.98)$. $\mu = 196$, $\sigma = \sqrt{3.92} = 1.98$. Overbooked if $X > 197$. $P(X \geq 198)$: with continuity correction, $P(X > 197.5)$. $Z = (197.5 - 196)/1.98 = 0.758$. $P(Z > 0.758) = 1 - 0.7757 = \boxed{0.2243}$.

6.8 Gamma & Exponential Distributions

6.8.1 Motivation: Connection to the Poisson Process



On a timeline of arrivals:

- $\#$ of arrivals $\sim \text{Poisson}(\lambda)$, where $\lambda =$ average $\#$ of arrivals per unit time.
- Time between arrivals (or until first arrival) $\sim \text{Exponential}\big(\beta = \frac{1}{\lambda}\big)$, where $\beta =$ average time until first arrival or between arrivals.
- Time until the αth arrival $\sim \text{Gamma}(\alpha, \beta)$.

6.8.2 The Gamma Function

<u>Def:</u> The **Gamma Function** is defined as

$$\boxed{\Gamma(\alpha) = \int_0^\infty x^{\alpha-1} e^{-x}\, dx, \qquad \text{for } \alpha > 0.}$$

Key properties:

- **Recursive:** $\Gamma(\alpha) = (\alpha - 1)\,\Gamma(\alpha - 1)$
- **Integer values:** $\Gamma(n) = (n-1)!$ if n is a positive integer
- **Special values:** $\Gamma\big(\frac{1}{2}\big) = \sqrt{\pi}$, $\Gamma(1) = 1$

6.8.3 Gamma Distribution

Def: A continuous r.v. X has the **gamma distribution** with parameters α and β if its pdf is

$$\boxed{f(x;\,\alpha,\beta) = \frac{1}{\beta^{\alpha}\,\Gamma(\alpha)}\,x^{\alpha-1}\,e^{-x/\beta}, \qquad x > 0,}$$

where $\alpha > 0,\ \beta > 0$.

Mean and variance:

$$\mu = \alpha\beta, \qquad \sigma^2 = \alpha\beta^2.$$

6.8.4 Graphs of the Gamma Distribution



The shape of the gamma pdf changes with α and β:

- $\alpha = 1,\ \beta = 1$: decreasing curve starting at $f(0) = 1$
- $\alpha = 2,\ \beta = 1$: rises then falls, peak shifts right
- $\alpha = 4,\ \beta = 1$: more bell-shaped, peak shifts further right

As α increases (with β fixed), the distribution becomes more symmetric and the peak moves to the right.

6.8.5 Exponential Distribution

If $\alpha = 1$, the gamma distribution reduces to the **exponential distribution**.

Def: $X \sim \mathrm{Exp}(\beta)$ if its pdf is

$$f(x;\,\beta) = \frac{1}{\beta}\,e^{-x/\beta}, \qquad x > 0,$$

where $\beta > 0$.

Mean and variance:

$$\mu = \beta, \qquad \sigma^2 = \beta^2.$$

6.8.6 Relation to the Poisson Process

Proof for time until the first occurrence.

Let $X = \#$ of arrivals in time t, so $X \sim \text{Poisson}(\lambda t)$. Recall the Poisson pdf:

$$P(X = x,\, \lambda t) = \frac{e^{-\lambda t}(\lambda t)^x}{x!}, \qquad x = 0, 1, 2, \ldots$$

The probability of *no* arrivals until time t:

$$P(X = 0,\, \lambda t) = \frac{e^{-\lambda t}(\lambda t)^0}{0!} = e^{-\lambda t}.$$

Now let Y = time until the first Poisson event. Then:

$$P(Y > y) = e^{-\lambda y}, \qquad y > 0 \quad \text{(prob. of no arrivals until time } y\text{)}.$$

CDF:

$$P(Y \le y) = 1 - e^{-\lambda y}, \qquad y > 0.$$

PDF (take derivative w.r.t. y):

$$f(y) = \lambda e^{-\lambda y}, \qquad y > 0.$$

This is the exponential distribution with $\lambda = \frac{1}{\beta}$.

6.8.7 Example: Cars at an Intersection

The number of cars arriving at a certain intersection per minute has a Poisson distribution with mean $\lambda = 5$.

(a) Probability that more than 10 cars appear during any given minute:

Let $X = \#$ of cars in one minute, so $X \sim \text{Poisson}(\mu = 5)$.

$$P(X > 10) = 1 - P(X \le 10) = 1 - \sum_{x=0}^{10} \frac{e^{-5}\,(5)^x}{x!} \approx 0.0137.$$

(b) Probability that more than 2 minutes are required before 10 cars arrive:

Let X = time until 10 events $\sim \text{Gamma}\big(\alpha = 10,\ \beta = \frac{1}{5}\big)$.

$$P(X > 2) = 1 - P(X \le 2) = 1 - \int_0^2 \frac{1}{\beta^\alpha\,\Gamma(\alpha)}\, x^{\alpha-1}\, e^{-x/\beta}\, dx$$

Change of variable: Let $y = x/\beta$, so $x = \beta y$ and $dx = \beta\, dy$, in order to utilize Table A.23 in the book.

$$P(X \le 2) = P(Y \le 10) = \int_0^{10} \frac{y^{\alpha-1}e^{-y}}{\Gamma(\alpha)}\, dy = \int_0^{10} \frac{y^9\, e^{-y}}{\Gamma(10)}\, dy = 0.542 \quad \text{(from Table A.23)}$$

Therefore:

$$P(X > 2) = 1 - 0.542 = 0.458.$$

Exercise 6.41

Problem: $X \sim \text{Gamma}(\alpha = 1, \beta = 1) = \text{Exp}(1)$. Find $P(1.6 < X < 2.8)$.
Solution: $P(1.6 < X < 2.8) = e^{-1.6} - e^{-2.8} = 0.2019 - 0.0608 = \boxed{0.1411}$.

6.9 Weibull Distribution, Reliability, Failure Rates & Chi-Squared Distribution

6.9.1 Review: Poisson Process and Related Distributions

Recall from the previous lecture that the **Poisson process** connects several important distributions. Here is a quick summary of the "big picture":

Poisson Process Summary

Consider events ("arrivals") occurring randomly along a time axis.

- **Number of arrivals** in a fixed time interval of length t: $X \sim \text{Poisson}(\lambda t)$, with pmf

$$P(X = x) = \frac{e^{-\lambda t}(\lambda t)^x}{x!}, \qquad x = 0, 1, 2, \ldots$$

  Here λ is the *rate* of arrivals (average number per unit time).

- **Time until the first arrival** (or any *interarrival* time): $\text{Exp}(\beta)$ where $\beta = 1/\lambda$.

$$f(x) = \frac{1}{\beta} e^{-x/\beta}, \qquad x > 0, \quad \beta > 0.$$

- **Time until the α-th arrival**: $\text{Gamma}(\alpha, \beta)$ with $\beta = 1/\lambda$.

Intuition: If you're sitting at a bus stop and buses arrive according to a Poisson process at rate $\lambda = 3$ buses per hour, then the number of buses in one hour is Poisson(3), the time until the next bus is Exponential with mean $\beta = 1/3$ hour (i.e. 20 minutes), and the time until the 5th bus is $\text{Gamma}(5, 1/3)$.

6.9.2 Example: Lifetime of Switches

Problem: The life, in years, of a certain type of switch has an exponential distribution with average life $\beta = 2$ years. If 100 of them are installed independently, what is the probability that at most 30 fail in the first year?

Step 1: Find the probability that a single switch fails within 1 year.

Let X = time until failure of a single switch. Then $X \sim \text{Exp}(\beta = 2)$, so:

$$P(X < 1) = \int_0^1 \frac{1}{2} e^{-x/2}\, dx = \left[-e^{-x/2}\right]_0^1 = 1 - e^{-1/2} \approx 0.3935.$$

So each switch has roughly a 39.35% chance of failing in the first year.

Step 2: Model the number of failures among 100 switches.

Let Y = number of switches (out of 100) that fail within 1 year. Since each switch fails independently with the same probability $p = 0.3935$:

$$Y \sim \text{Binomial}(n = 100,\ p = 0.3935).$$

We want $P(Y \leq 30)$. This is a sum of binomial probabilities:

$$P(Y \leq 30) = \sum_{y=0}^{30} \binom{100}{y} p^y (1-p)^{100-y}.$$

Step 3: Use the normal approximation.

Since $n = 100$ is large, we can approximate using the normal distribution. Compute:

$$\mu = np = 100 \times 0.3935 = 39.35, \qquad \sigma^2 = npq = 100 \times 0.3935 \times 0.6065 \approx 23.86.$$

Using the **continuity correction** (adding 0.5 because we are approximating a discrete distribution with a continuous one):

$$P(Y \leq 30) \approx P\left(Z \leq \frac{30 + 0.5 - np}{\sqrt{npq}}\right) = P\left(Z \leq \frac{30.5 - 39.35}{\sqrt{23.86}}\right) = P(Z \leq -1.81) \approx 0.035.$$

So there is only about a 3.5% chance that at most 30 switches fail in the first year.

Why the Continuity Correction?

When approximating a discrete distribution (binomial) with a continuous one (normal), the value "$Y \leq 30$" in the discrete world corresponds to "$Y \leq 30.5$" in the continuous world. This small adjustment of ± 0.5 is called the **continuity correction** and improves the accuracy of the approximation.

6.9.3 Memoryless Property of the Exponential Distribution

Theorem: Memoryless Property

The exponential distribution is **memoryless**. That is, for $X \sim \text{Exp}(\beta)$:

$$P(X \geq t) = P(X \geq t_0 + t \mid X \geq t_0).$$

What does this mean in plain English?

Suppose X represents the lifetime of a light bulb. The memoryless property says:

> If a light bulb has already survived t_0 hours, the probability of it lasting an *additional* t hours is the same as the probability of a brand-new bulb lasting t hours from the start.

In other words, the component does not "age" — its past history gives you no information about how much longer it will last. This is a very special property: among all continuous distributions, *only* the exponential distribution has it.

Quick proof sketch:

Recall that the survival function for $X \sim \text{Exp}(\beta)$ is:

$$P(X \geq t) = e^{-t/\beta}.$$

Then:

$$P(X \geq t_0 + t \mid X \geq t_0) = \frac{P(X \geq t_0 + t)}{P(X \geq t_0)} = \frac{e^{-(t_0+t)/\beta}}{e^{-t_0/\beta}} = e^{-t/\beta} = P(X \geq t). \qquad \square$$

6.9.4 Weibull Distribution

The exponential distribution assumes a *constant* failure rate (memoryless property). But in real life, many components either wear out over time (increasing failure rate) or "burn in" and improve (decreasing failure rate). The **Weibull distribution** generalizes the exponential to handle these cases.

Definition: Weibull Distribution

A continuous random variable X has a **Weibull distribution** with parameters $\alpha > 0$ and $\beta > 0$ if its pdf is:

$$f(x;\, \alpha,\, \beta) = \begin{cases} \alpha\beta\, x^{\beta-1}\, e^{-\alpha x^{\beta}}, & x > 0, \\ 0, & \text{otherwise.} \end{cases}$$

CDF:

$$F(x) = 1 - e^{-\alpha x^{\beta}}, \qquad x \geq 0.$$

Mean and Variance:

$$\mu = \alpha^{-1/\beta}\, \Gamma\left(1 + \frac{1}{\beta}\right), \qquad \sigma^2 = \alpha^{-2/\beta} \left\{ \Gamma\left(1 + \frac{2}{\beta}\right) - \left[\Gamma\left(1 + \frac{1}{\beta}\right)\right]^2 \right\}.$$

Here $\Gamma(\cdot)$ is the gamma function from the previous lecture.

Special Case: β

When $\beta = 1$, the Weibull pdf simplifies to $f(x) = \alpha\, e^{-\alpha x}$, which is exactly the exponential distribution with rate α (i.e. $\beta_{\text{exp}} = 1/\alpha$). So the exponential is a special case of the Weibull.

6.9.5 Reliability and Failure Rate (Hazard Rate)

In engineering, we often care about how long a product will function before failing. This leads to the concepts of **reliability** and **failure rate**.

6.9.5.1 Reliability Function

Definition: Reliability

Reliability is the probability that a product will function properly for at least a specified time under specified conditions.

Let T be the random variable representing the *time until failure* of a component, with pdf $f(t)$ and cdf $F(t)$. The **reliability function** is:

$$R(t) = P(T > t) = \int_t^\infty f(s)\,ds = 1 - F(t).$$

Interpretation: $R(t)$ tells you the probability that the component is still working at time t. For example, $R(5) = 0.90$ means there is a 90% chance the component survives past 5 years.

6.9.5.2 Conditional Failure Probability

Now suppose we know a component has survived until time t. What is the probability it fails in the small interval from t to $t + \Delta t$?

$$P(t < T < t + \Delta t \mid T > t) = \frac{P(t < T < t + \Delta t)}{P(T > t)} = \frac{F(t + \Delta t) - F(t)}{R(t)}.$$

This is a *conditional probability*: given that the component has survived to time t, how likely is it to fail in the next Δt units of time?

6.9.5.3 Failure Rate (Hazard Rate)

The **failure rate** (also called the **hazard rate**) measures the rate of change of this conditional probability as $\Delta t \to 0$:

$$z(t) = \lim_{\Delta t \to 0} \frac{F(t + \Delta t) - F(t)}{R(t) \cdot \Delta t} = \frac{F'(t)}{R(t)} = \frac{f(t)}{R(t)} = \frac{f(t)}{1 - F(t)}.$$

Intuition: The hazard rate $z(t)$ tells you the "instantaneous risk of failure" at time t, given that the component has survived up to time t. A high $z(t)$ means the component is very likely to fail soon; a low $z(t)$ means it's relatively safe.

6.9.5.4 Failure Rate for the Weibull Distribution

For the Weibull distribution with parameters α and β:

$$z(t) = \alpha\beta\, t^{\beta-1}, \qquad t > 0, \quad \alpha, \beta > 0.$$

This is one of the most important results in reliability theory, because the parameter β controls how the failure rate changes over time:

1. **If** $\beta = 1$: $z(t) = \alpha$ (constant). The failure rate does not change over time. This is the exponential distribution — the component does not age (memoryless property).
2. **If** $\beta > 1$: $z(t) = \alpha\beta\, t^{\beta-1}$ is an *increasing* function of t. The failure rate gets higher over time. This models components that **wear out** (e.g. brake pads, batteries, tires).
3. **If** $\beta < 1$: $z(t)$ is a *decreasing* function of t. The failure rate gets lower over time. This models components that **strengthen over time** or experience "infant mortality" (early failures are weeded out, and survivors become more reliable).

The Bathtub Curve

In practice, many products exhibit a failure rate that combines all three phases: high early failure ($\beta < 1$), then a stable period ($\beta \approx 1$), then increasing failures due to wear-out ($\beta > 1$). When plotted, the hazard rate looks like a bathtub — hence the name "bathtub curve."

6.9.6 Chi-Squared Distribution

The **chi-squared distribution** is another special case of the gamma distribution. It will become extremely important later in the course when we study hypothesis testing and confidence intervals.

Definition: Chi-Squared Distribution

The chi-squared distribution with ν degrees of freedom is a **special case of the gamma distribution** with:

$$\alpha = \frac{\nu}{2}, \qquad \beta = 2.$$

We write $X \sim \chi^2(\nu)$.

Mean and Variance:

Since it is a gamma distribution with $\alpha = \nu/2$ and $\beta = 2$:

$$\mu = \alpha\beta = \frac{\nu}{2} \cdot 2 = \nu, \qquad \sigma^2 = \alpha\beta^2 = \frac{\nu}{2} \cdot 4 = 2\nu.$$

Key facts:

- ν must be a *positive integer* and is called the **degrees of freedom**.
- The mean equals the degrees of freedom: $\mu = \nu$.

- The variance is twice the degrees of freedom: $\sigma^2 = 2\nu$.
- As ν increases, the chi-squared distribution becomes more symmetric and approaches a normal distribution (by the Central Limit Theorem).

Why Does This Matter?

The chi-squared distribution arises naturally when you square standard normal random variables. If $Z_1, Z_2, \ldots, Z_\nu$ are independent standard normal random variables, then:

$$Z_1^2 + Z_2^2 + \cdots + Z_\nu^2 \sim \chi^2(\nu).$$

This connection makes the chi-squared distribution fundamental in statistics — it appears in goodness-of-fit tests, tests of independence, and confidence intervals for the variance.

Exercise 6.47

Problem: Suppose that the service life, in years, of a hearing aid battery is a random variable having a Weibull distribution with $\alpha = 1/2$ and $\beta = 2$. (1) How long can such a battery be expected to last? (2) What is the probability that such a battery will be operating after 2 years?

Solution: Weibull mean: $E(X) = \beta\,\Gamma(1 + 1/\alpha) = 2\,\Gamma(3) = 2 \times 2 = \boxed{4}$ years.

(2) $P(X > 2) = e^{-(2/2)^{1/2}} = e^{-1} = \boxed{0.3679}$.

Chapter 7

Chapter 7: Transformations of Random Variables

7.1 Functions of Random Variables

7.1.1 Motivation

A very common problem in probability and statistics is: if we know the distribution of a random variable X, and we define a new random variable $Y = u(X)$ (some function of X), what is the distribution of Y?

Real-world example: Suppose X is the radius of a randomly manufactured ball bearing, and we know the distribution of X. The cross-sectional area is $Y = \pi X^2$. What is the distribution of Y?

7.1.2 Discrete Case: One-to-One Transformations

Theorem: Discrete One-to-One Transformation

Suppose X is a discrete random variable with pmf $f(x)$. Let $Y = u(X)$ define a **one-to-one** transformation, so that $y = u(x)$ can be uniquely solved for x:

$$x = w(y) \qquad \text{(the inverse function).}$$

Then the pmf of Y is:

$$g(y) = f[w(y)]\,.$$

What does "one-to-one" mean? A function is one-to-one if each output value y comes from exactly one input value x. For example, $y = 2x + 3$ is one-to-one (you can solve back to get $x = (y - 3)/2$), but $y = x^2$ is *not* one-to-one (both $x = 2$ and $x = -2$ give $y = 4$).

Intuition: If the transformation is one-to-one, then each value y of the new variable corresponds to exactly one value x of the old variable. So the probability of $Y = y$ is the same as the probability of $X = w(y)$, which is just $f(w(y))$.

7.1.2.1 Example: Geometric Distribution

Suppose X has a geometric distribution:

$$f(x) = \frac{3}{4}\left(\frac{1}{4}\right)^{x-1}, \qquad x = 1, 2, 3, \ldots$$

Let $Y = X^2$. Since X only takes positive integer values, the map $x \mapsto x^2$ is one-to-one on this domain (distinct positive integers have distinct squares). The inverse is $x = \sqrt{y}$, so:

$$g(y) = f(\sqrt{y}) = \frac{3}{4}\left(\frac{1}{4}\right)^{\sqrt{y}-1}, \qquad y = 1, 4, 9, 16, \ldots$$

We can verify: $P(Y = 1) = P(X = 1) = 3/4$, $P(Y = 4) = P(X = 2) = 3/16$, $P(Y = 9) = P(X = 3) = 3/64$, and so on.

7.1.3 Discrete Case: Non-One-to-One Transformations

When the transformation is *not* one-to-one, multiple values of X can map to the same value of Y. In this case, we need to sum over all x-values that produce the same y.

7.1.3.1 Example: $Y = X^2$ with Symmetric Support

Suppose X takes values $\{-2, -1, 0, 1, 2\}$ with some pmf, and let $Y = X^2$.



Notice that both $X = 2$ and $X = -2$ map to $Y = 4$, and both $X = 1$ and $X = -1$ map to $Y = 1$. So:

$$g(y) = \begin{cases} P(X = \sqrt{y}) + P(X = -\sqrt{y}), & \text{if } y > 0, \\ P(X = 0), & \text{if } y = 0, \\ 0, & \text{if } y < 0. \end{cases}$$

Key idea: For non-one-to-one transformations, collect all x-values that produce the same y, and add up their probabilities.



The parabola $y = x^2$ maps both x and $-x$ to the same y-value. This is why we must sum probabilities from both branches.

7.1.4 Discrete Case: Joint Transformations (Two Variables)

Theorem: Joint Discrete Transformation

Suppose X_1, X_2 are discrete random variables with joint pmf $f(x_1, x_2)$. Let:

$$Y_1 = u_1(X_1, X_2), \qquad Y_2 = u_2(X_1, X_2)$$

be a **one-to-one** transformation. That is, given (y_1, y_2), we can uniquely solve back:

$$x_1 = w_1(y_1, y_2), \qquad x_2 = w_2(y_1, y_2).$$

Then the joint pmf of (Y_1, Y_2) is:

$$g(y_1, y_2) = f[w_1(y_1, y_2),\ w_2(y_1, y_2)]\,.$$

Why do we need two new variables? Even if we only care about the distribution of Y_1, the technique requires us to define a second variable Y_2 so that the transformation is invertible (one-to-one). After finding the joint pmf $g(y_1, y_2)$, we can then "marginalize out" Y_2 by summing over all its possible values to get the marginal pmf of Y_1.

7.1.4.1 Example: Sum of Independent Poissons

Problem: X_1 and X_2 are independent Poisson random variables with means μ_1 and μ_2 respectively. Find the pmf of $Y = X_1 + X_2$.

Step 1: Set up the transformation.

Define:

$$Y_1 = X_1 + X_2, \qquad Y_2 = X_2.$$

The inverse transformation is:

$$X_1 = Y_1 - Y_2, \qquad X_2 = Y_2.$$

The domain is $y_1 = 0, 1, 2, \ldots$ and $y_2 = 0, 1, \ldots, y_1$ (since $X_1 = Y_1 - Y_2 \geq 0$).

Step 2: Write the joint pmf.

Since X_1 and X_2 are independent:

$$f(x_1, x_2) = \frac{e^{-\mu_1}\,\mu_1^{x_1}}{x_1!} \cdot \frac{e^{-\mu_2}\,\mu_2^{x_2}}{x_2!}, \qquad x_1, x_2 = 0, 1, 2, \ldots$$

Substituting $x_1 = y_1 - y_2$ and $x_2 = y_2$:

$$g(y_1, y_2) = \frac{e^{-\mu_1}\,\mu_1^{y_1 - y_2}}{(y_1 - y_2)!} \cdot \frac{e^{-\mu_2}\,\mu_2^{y_2}}{y_2!}.$$

Step 3: Find the marginal pmf of Y_1 by summing over Y_2.

$$h(y_1) = \sum_{y_2=0}^{y_1} g(y_1, y_2) = e^{-(\mu_1+\mu_2)} \sum_{y_2=0}^{y_1} \frac{\mu_1^{y_1-y_2}\,\mu_2^{y_2}}{(y_1 - y_2)!\,y_2!}.$$

Multiply and divide by $y_1!$:

$$h(y_1) = \frac{e^{-(\mu_1+\mu_2)}}{y_1!} \sum_{y_2=0}^{y_1} \binom{y_1}{y_2} \mu_1^{y_1-y_2}\,\mu_2^{y_2}.$$

By the **binomial theorem**, $\displaystyle\sum_{y_2=0}^{y_1} \binom{y_1}{y_2} \mu_1^{y_1-y_2} \mu_2^{y_2} = (\mu_1 + \mu_2)^{y_1}$. Therefore:

$$\boxed{h(y_1) = \frac{e^{-(\mu_1+\mu_2)}\,(\mu_1 + \mu_2)^{y_1}}{y_1!}, \qquad y_1 = 0, 1, 2, \ldots}$$

This is a Poisson$(\mu_1 + \mu_2)$ distribution!

Important Result

The sum of independent Poisson random variables is again Poisson, with the mean equal to the sum of the individual means. This is an example of the **reproductive property** of the Poisson distribution.

7.1.5 Continuous Case: CDF Method

For continuous random variables, we often find the distribution of $Y = u(X)$ by first finding the **CDF** of Y, and then differentiating to get the pdf. This approach is called the **CDF method** (or **distribution function method**).

7.1.5.1 Example: $X \sim$ Uniform$(0, 1)$, $Y = X^2$

Suppose $X \sim \text{Uniform}(0, 1)$, so $f(x) = 1$ for $0 < x < 1$. Let $Y = X^2$.





Step 1: Find the CDF of Y.

For $0 < y < 1$:

$$P(Y \leq y) = P(X^2 \leq y) = P(X \leq \sqrt{y}) \quad \text{(since } X > 0\text{)}.$$

The CDF of X is $P(X \leq x) = \int_0^x 1\,dt = x$ for $0 < x < 1$. So:

$$P(Y \leq y) = \sqrt{y}, \qquad 0 < y < 1.$$

Step 2: Differentiate to get the pdf.

$$f_Y(y) = \frac{d}{dy}\sqrt{y} = \frac{1}{2\sqrt{y}}, \qquad 0 < y < 1.$$

Sanity check: We can verify: $P(X < 0.5) = 0.5$, and $P(Y < 0.25) = \sqrt{0.25} = 0.5$. Since $Y = X^2$, the event $\{X < 0.5\}$ is the same as $\{Y < 0.25\}$, so the probabilities match. ✓

7.1.5.2 Example: $f(x) = (x+1)/2$, $Y = X^2$

Let $f(x) = (x+1)/2$ for $-1 < x < 1$, and $Y = X^2$.

For $0 < y < 1$:

$$P(Y \leq y) = P(X^2 \leq y) = P(-\sqrt{y} \leq X \leq \sqrt{y}) = \int_{-\sqrt{y}}^{\sqrt{y}} \frac{x+1}{2}\,dx = \sqrt{y}, \qquad 0 < y \leq 1.$$

Therefore:

$$f_Y(y) = \frac{d}{dy}\sqrt{y} = \frac{1}{2\sqrt{y}}, \qquad 0 < y < 1.$$

7.1.6 Continuous Case: Change-of-Variable (Jacobian) Method

The CDF method always works but can be tedious. For one-to-one transformations, there is a more direct approach using the **Jacobian**.

Theorem: Continuous One-to-One Transformation

Suppose X is a continuous random variable with pdf $f(x)$. Let $Y = u(X)$ be a one-to-one transformation with inverse $x = w(y)$. Then the pdf of Y is:

$$g(y) = f[w(y)] \cdot |J|,$$

where $J = w'(y) = \dfrac{dx}{dy}$ is the **Jacobian** of the transformation.

Why the absolute value of the derivative? When we change variables, the "stretching" or "compression" of the x-axis by the transformation needs to be accounted for so that probabilities are preserved. The Jacobian $|dx/dy|$ captures this scaling factor.

7.1.6.1 Example: $f(x) = x^3/4$, $Y = X^6$

Let $f(x) = x^3/4$ for $0 < x < 2$, and $Y = X^6$.

Using the CDF method:

For $0 < y < 64$ (since $x \in (0, 2)$ means $y = x^6 \in (0, 64)$):

$$P(Y \leq y) = P(X^6 \leq y) = P(X \leq y^{1/6}) = \int_0^{y^{1/6}} \frac{x^3}{4}\, dx = \frac{x^4}{16}\bigg|_0^{y^{1/6}} = \frac{(y^{1/6})^4}{16} = \frac{y^{2/3}}{16}.$$

Differentiating:

$$g(y) = \frac{d}{dy}\frac{y^{2/3}}{16} = \frac{2}{48}\, y^{-1/3} = \frac{y^{-1/3}}{24}, \qquad 0 < y < 64.$$

Using the Jacobian method:

The inverse transformation is $x = y^{1/6} = w(y)$. The Jacobian is:

$$J = w'(y) = \frac{1}{6}\, y^{-5/6}.$$

Then:

$$g(y) = f(y^{1/6}) \cdot |J| = \frac{(y^{1/6})^3}{4} \cdot \frac{1}{6}\, y^{-5/6} = \frac{y^{1/2}}{4} \cdot \frac{1}{6}\, y^{-5/6} = \frac{y^{-1/3}}{24}, \qquad 0 < y < 64.$$

Both methods give the same answer, as expected.

7.1.7 Continuous Case: Non-One-to-One Transformations

When the transformation $Y = u(X)$ is *not* one-to-one on the support of X, different x-values can produce the same y-value. We handle this by partitioning the domain.

Theorem: Continuous Non-One-to-One Transformation

Suppose X is a continuous r.v. with pdf $f(x)$. Let $u = Y(X)$ be **not** a one-to-one transformation. If the interval over which X is defined can be partitioned into k mutually disjoint sets such that each inverse function

$$x_i = w_i(y), \qquad i = 1, \ldots, k$$

of $y = u(x)$ defines a one-to-one correspondence, then the pdf of Y is:

$$g(y) = \sum_{i=1}^{k} f[w_i(y)]\ |J_i|, \qquad \text{where } J_i = w_i'(y),\ i = 1, \ldots, k.$$

Intuition: We split the domain of X into pieces where u is one-to-one on each piece, apply the Jacobian method to each piece separately, and add up the contributions.

7.1.7.1 Example: Showing $Y = \left(\frac{X-\mu}{\sigma}\right)^2 \sim \chi^2(1)$ when $X \sim N(\mu, \sigma^2)$

If $X \sim N(\mu, \sigma^2)$, let $Z = (X - \mu)/\sigma \sim N(0, 1)$ and $Y = Z^2$.

The standard normal pdf is $f(z) = \frac{1}{\sqrt{2\pi}}\, e^{-z^2/2}$ for $-\infty < z < \infty$.

The transformation $y = z^2$ is *not* one-to-one: both z and $-z$ give the same y. We partition into two sets:

- For $z < 0$: $z_1 = -\sqrt{y}$, so $J_1 = \dfrac{dz_1}{dy} = -\dfrac{1}{2\sqrt{y}}$.
- For $z > 0$: $z_2 = \sqrt{y}$, so $J_2 = \dfrac{dz_2}{dy} = \dfrac{1}{2\sqrt{y}}$.

Applying the theorem:

$$g(y) = f(-\sqrt{y}) \cdot \left|\frac{-1}{2\sqrt{y}}\right| + f(\sqrt{y}) \cdot \left|\frac{1}{2\sqrt{y}}\right| = \frac{1}{\sqrt{2\pi}}\, e^{-y/2} \cdot \frac{1}{2\sqrt{y}} + \frac{1}{\sqrt{2\pi}}\, e^{-y/2} \cdot \frac{1}{2\sqrt{y}}.$$

$$g(y) = \frac{1}{\sqrt{2\pi}}\, y^{1/2-1}\, e^{-y/2}, \qquad y > 0.$$

This is a Gamma($\alpha = 1/2,\ \beta = 2$) distribution, which is exactly $\chi^2(1)$.

Key Result

If $X \sim N(\mu, \sigma^2)$, then $\left(\dfrac{X-\mu}{\sigma}\right)^2 \sim \chi^2(1)$. This fundamental result is used extensively in hypothesis testing and confidence intervals.

7.2 Moments and Moment Generating Functions (MGFs)

7.2.1 Moments About Zero

Def: The r**-th moment about zero** of a random variable X is:

$$\mu'_r = E(X^r) = \begin{cases} \sum_x x^r f(x), & X \text{ discrete}, \\ \int_{-\infty}^{\infty} x^r f(x)\, dx, & X \text{ continuous}. \end{cases}$$

Important special cases:

- **First moment:** $\mu'_1 = E(X) = \mu$ (the mean)
- **Second moment:** $\mu'_2 = E(X^2) = \sigma^2 + \mu^2$

7.2.2 Central Moments

Def: The r**-th central moment** (moment about the mean μ) is:

$$E[(X-\mu)^r]\,.$$

Important special cases:

- **2nd central moment:** $E[(X-\mu)^2] = \sigma^2$ (the variance)
- **3rd central moment:** related to **skewness** (asymmetry of the distribution)
- **4th central moment:** related to **kurtosis** (heaviness of tails)

7.2.3 Moment Generating Functions

Def: The **moment generating function (MGF)** of X is:

$$\boxed{M_X(t) = E(e^{tX}) = \begin{cases} \sum_x e^{tx} f(x), & X \text{ discrete}, \\ \int_{-\infty}^{\infty} e^{tx} f(x)\, dx, & X \text{ continuous}. \end{cases}}$$

The MGF exists if the sum or integral converges in a neighborhood of $t = 0$.

Theorem: MGF Generates Moments

If the MGF $M_X(t)$ exists, then the r-th moment about zero is obtained by differentiating r times and evaluating at $t = 0$:

$$\frac{d^r}{dt^r} M_X(t)\bigg|_{t=0} = \mu'_r = E(X^r).$$

Why "moment generating"? Expanding e^{tX} as a Taylor series gives $e^{tX} = 1 + tX + \frac{t^2X^2}{2!} + \cdots$, so $M_X(t) = 1 + t\,E(X) + \frac{t^2}{2!}E(X^2) + \cdots$. Taking derivatives at $t = 0$ peels off successive moments.

7.2.4 Example: MGF of the Exponential Distribution

Let $X \sim \text{Exp}(\beta)$ with $f(x) = \frac{1}{\beta}\,e^{-x/\beta}$ for $x > 0$.

$$M_X(t) = E(e^{tX}) = \int_0^\infty e^{tx}\,\frac{1}{\beta}\,e^{-x/\beta}\,dx = \frac{1}{\beta}\int_0^\infty e^{(t-1/\beta)x}\,dx.$$

This integral converges when $t < 1/\beta$:

$$M_X(t) = \frac{1}{\beta}\cdot\frac{1}{1/\beta - t} = \frac{1}{1-\beta t}, \qquad t < \frac{1}{\beta}.$$

Finding the mean:

$$M_X'(t) = \frac{\beta}{(1-\beta t)^2} \quad\Longrightarrow\quad E(X) = M_X'(0) = \beta.$$

Finding the variance:

$$M_X''(t) = \frac{2\beta^2}{(1-\beta t)^3} \quad\Longrightarrow\quad E(X^2) = M_X''(0) = 2\beta^2.$$

$$\sigma^2 = E(X^2) - \mu^2 = 2\beta^2 - \beta^2 = \beta^2.$$

This confirms $\mu = \beta$ and $\sigma^2 = \beta^2$ for the exponential distribution.

7.2.5 Uniqueness Theorem

Theorem: MGF Uniqueness

Let X and Y be two random variables with MGFs $M_X(t)$ and $M_Y(t)$ respectively. If $M_X(t) = M_Y(t)$ for all values of t in some neighborhood of 0, then X and Y have the **same distribution**.

Why this matters: The MGF *uniquely characterizes* a distribution. If we can show that a random variable's MGF matches a known distribution's MGF, we can conclude it *is* that distribution — without needing to derive the pdf directly.

7.2.6 Properties of MGFs

Property 1: Linear transformation. If $Y = a + bX$ where $a, b \in \mathbb{R}$, then:

$$M_Y(t) = e^{at}\,M_X(bt).$$

Proof: $M_Y(t) = E(e^{tY}) = E(e^{t(a+bX)}) = E(e^{ta}\cdot e^{btX}) = e^{ta}\,E(e^{btX}) = e^{at}\,M_X(bt)$. $\square$

Property 2: Sum of independent random variables. If $X_1, \ldots, X_n$ are **independent** r.v.'s with MGFs

$M_{X_1}(t), \ldots, M_{X_n}(t)$, and $Y = X_1 + \cdots + X_n$, then:

$$\boxed{M_Y(t) = M_{X_1}(t) \cdot M_{X_2}(t) \cdots M_{X_n}(t).}$$

Proof (for $n = 2$, the general case follows by induction):

$$\begin{aligned} M_Y(t) &= E(e^{tY}) = E\Big(e^{t(X_1+X_2)}\Big) = E\big(e^{tX_1} \cdot e^{tX_2}\big) \\ &= E(e^{tX_1}) \cdot E(e^{tX_2}) \qquad \text{(by independence)} \\ &= M_{X_1}(t) \cdot M_{X_2}(t). \qquad \square \end{aligned}$$

For the continuous case with joint pdf factoring as $f_1(x_1) \cdot f_2(x_2) \cdots f_n(x_n)$:

$$M_Y(t) = \int_{x_1} \cdots \int_{x_n} e^{t(x_1+\cdots+x_n)} f_1(x_1) \cdots f_n(x_n)\, dx_1 \cdots dx_n = \prod_{i=1}^{n} \int_{x_i} e^{tx_i} f_i(x_i)\, dx_i = \prod_{i=1}^{n} M_{X_i}(t).$$

7.2.7 Example: Sum of Independent Poissons via MGFs

Let $X_1 \sim \text{Poisson}(\lambda_1)$ and $X_2 \sim \text{Poisson}(\lambda_2)$, independent. Find the distribution of $Y = X_1 + X_2$.

The MGF of a Poisson(λ) random variable is:

$$M_X(t) = E(e^{tX}) = e^{\lambda(e^t - 1)}.$$

By Property 2:

$$M_Y(t) = M_{X_1}(t) \cdot M_{X_2}(t) = e^{\lambda_1(e^t-1)} \cdot e^{\lambda_2(e^t-1)} = e^{(\lambda_1+\lambda_2)(e^t-1)}.$$

By the uniqueness theorem, this is the MGF of Poisson($\lambda_1 + \lambda_2$).

MGFs of Common Distributions

- $X \sim N(\mu, \sigma^2)$: $\quad M_X(t) = \exp\big(\mu t + \frac{1}{2}\sigma^2 t^2\big)$
- $X \sim \chi^2(\nu)$: $\quad M_X(t) = (1 - 2t)^{-\nu/2}$
- $X \sim \text{Poisson}(\lambda)$: $\quad M_X(t) = e^{\lambda(e^t-1)}$
- $X \sim \text{Exp}(\beta)$: $\quad M_X(t) = (1 - \beta t)^{-1}, \ t < 1/\beta$
- $X \sim \text{Gamma}(\alpha, \beta)$: $\quad M_X(t) = (1 - \beta t)^{-\alpha}, \ t < 1/\beta$

7.3 Linear Combinations of Normal Random Variables

One of the most powerful results in probability is that linear combinations of independent normal random variables are again normal.

Theorem: Linear Combinations of Independent Normals

If $X_1, \ldots, X_n$ are **independent** normal random variables with means $\mu_1, \ldots, \mu_n$ and variances $\sigma_1^2, \ldots, \sigma_n^2$, and $a_1, \ldots, a_n$ are real constants, then:

$$Y = a_1X_1 + a_2X_2 + \cdots + a_nX_n \sim N\big(a_1\mu_1 + \cdots + a_n\mu_n,\ \ a_1^2\sigma_1^2 + \cdots + a_n^2\sigma_n^2\big)\,.$$

Proof (for $n = 2$): Let $Y = a_1X_1 + a_2X_2$. Using the MGF:

$$\begin{aligned} M_Y(t) &= M_{X_1}(a_1t) \cdot M_{X_2}(a_2t) \\ &= \exp\big(a_1\mu_1 t + \tfrac{1}{2}a_1^2\sigma_1^2t^2\big) \cdot \exp\big(a_2\mu_2 t + \tfrac{1}{2}a_2^2\sigma_2^2t^2\big) \\ &= \exp\big[(a_1\mu_1 + a_2\mu_2)t + \tfrac{1}{2}(a_1^2\sigma_1^2 + a_2^2\sigma_2^2)t^2\big]\,. \end{aligned}$$

This is the MGF of $N(a_1\mu_1 + a_2\mu_2,\ a_1^2\sigma_1^2 + a_2^2\sigma_2^2)$, so by the uniqueness theorem, Y has this distribution. □

7.3.1 Special Cases

1) Sum of independent normals: If $X_i \sim N(\mu_i, \sigma_i^2)$ independently, with all $a_i = 1$:

$$X_1 + X_2 + \cdots + X_n \sim N\big(\mu_1 + \cdots + \mu_n,\ \ \sigma_1^2 + \cdots + \sigma_n^2\big)\,.$$

2) IID normals — sum and sample mean: If $X_1, \ldots, X_n \overset{\text{iid}}{\sim} N(\mu, \sigma^2)$:

$$Y = X_1 + X_2 + \cdots + X_n \sim N(n\mu,\ n\sigma^2),$$

$$\bar{X} = \frac{X_1 + \cdots + X_n}{n} \sim N\left(\mu,\ \frac{\sigma^2}{n}\right).$$

Interpretation: The sample mean $\bar{X}$ is also normally distributed, centered at μ, but with variance σ^2/n. As the sample size n increases, the variance of $\bar{X}$ decreases — the sample mean becomes a more precise estimator of μ.

7.3.2 Chi-Squared Additivity and Corollaries

Theorem: Sum of Independent Chi-Squared R.V.s

If $X_1, \ldots, X_n$ are independent chi-squared random variables with degrees of freedom $\nu_1, \ldots, \nu_n$, then:

$$Y = X_1 + \cdots + X_n \sim \chi^2(\nu_1 + \cdots + \nu_n).$$

Proof: Using MGFs, $M_Y(t) = \prod_{i=1}^n (1-2t)^{-\nu_i/2} = (1-2t)^{-(\nu_1+\cdots+\nu_n)/2}$, which is the MGF of $\chi^2(\nu_1+\cdots+\nu_n)$. □

Corollary 1: If $X_1, \ldots, X_n \overset{\text{iid}}{\sim} N(\mu, \sigma^2)$, then:

$$Y = \sum_{i=1}^{n} \left(\frac{X_i - \mu}{\sigma} \right)^2 \sim \chi^2(n).$$

Why? Each $\left(\frac{X_i-\mu}{\sigma}\right)^2 \sim \chi^2(1)$ (as shown earlier), and they are independent. By the additivity theorem, their sum is $\chi^2(n)$.

Corollary 2 (more general): If $X_i \sim N(\mu_i, \sigma_i^2)$ independently (not necessarily identically distributed), then:

$$Y = \sum_{i=1}^{n} \left(\frac{X_i - \mu_i}{\sigma_i} \right)^2 \sim \chi^2(n).$$

Exercise 7.1

Problem: X uniform on $\{1, \ldots, 6\}$. Find distribution of $Y = 2X + 1$.
Solution: Y takes values $3, 5, 7, 9, 11, 13$ each with probability $1/6$. $P(Y = y) = 1/6$ for $y \in \{3, 5, 7, 9, 11, 13\}$.

Exercise 7.2

Problem: $X \sim \text{Bin}(3, 1/3)$. Find distribution of $Y = X^2$.
Solution: $P(X = 0) = (2/3)^3 = 8/27$, $P(X = 1) = 3(1/3)(2/3)^2 = 12/27$, $P(X = 2) = 3(1/3)^2(2/3) = 6/27$, $P(X = 3) = 1/27$.
$Y = X^2$: $P(Y = 0) = 8/27$, $P(Y = 1) = 12/27$, $P(Y = 4) = 6/27$, $P(Y = 9) = 1/27$.

Exercise 7.4

Problem: $f(x, y) = \frac{x+2y}{27}$ for $x, y \in \{0, 1, 2\}$. Find distribution of $Z = X + Y$.
Solution: Z takes values $0, 1, 2, 3, 4$. Sum $f(x, y)$ over all (x, y) with $x + y = z$:
$P(Z = 0) = f(0, 0) = 0/27 = 0$. $P(Z = 1) = f(0, 1) + f(1, 0) = 2/27 + 1/27 = 3/27$. $P(Z = 2) = f(0, 2) + f(1, 1) + f(2, 0) = 4/27 + 3/27 + 2/27 = 9/27$. $P(Z = 3) = f(1, 2) + f(2, 1) = 5/27 + 4/27 = 9/27$. $P(Z = 4) = f(2, 2) = 6/27$.
Check: $0 + 3 + 9 + 9 + 6 = 27$. ✓

Exercise 7.9

Problem: $Y = X + 4$ where $f(x) = \frac{32}{(x+4)^3}$ for $x > 0$. (1) PDF of Y. (2) $P(Y > 8)$.
Solution: (1) $x = y - 4$, $dx/dy = 1$. For $y > 4$: $g(y) = f(y - 4) \cdot |1| = \frac{32}{y^3}$.
(2) $P(Y > 8) = \int_8^\infty \frac{32}{y^3}\, dy = \left[-\frac{16}{y^2}\right]_8^\infty = \frac{16}{64} = \boxed{1/4}$.

Exercise 7.13

Problem: Current $f(i) = 6i(1 - i)$ for $0 < i < 1$. Resistance $g(r) = 2r$ for $0 < r < 1$. Independent. Find distribution of $W = I^2R$.
Solution: This requires finding the pdf of W via a two-variable transformation. Define $W = I^2R$,

$V = R$. Then $I = \sqrt{W/V}$, $R = V$.
Jacobian: $|J| = \frac{1}{2\sqrt{WV}}$. Domain: $0 < W/V < 1$ and $0 < V < 1$, so $W < V$ and $0 < V < 1$ with $W > 0$.
$h(w, v) = f(\sqrt{w/v})\, g(v)\, \frac{1}{2v\sqrt{w/v}} = 6\sqrt{w/v}(1 - \sqrt{w/v}) \cdot 2v \cdot \frac{1}{2\sqrt{wv}}$.
Marginalizing over v gives the pdf of W. The integral is complex; the result can be evaluated numerically or left in integral form.

Exercise 7.15

Problem: $f(x) = \frac{2(x+1)}{9}$ for $-1 < x < 2$. Find distribution of $Y = X^2$.
Solution: Y ranges from 0 to 4. For $0 < y < 1$: two roots $x = \pm\sqrt{y}$. $g(y) = \frac{f(\sqrt{y})}{2\sqrt{y}} + \frac{f(-\sqrt{y})}{2\sqrt{y}} = \frac{1}{2\sqrt{y}}\left[\frac{2(\sqrt{y}+1)}{9} + \frac{2(1-\sqrt{y})}{9}\right] = \frac{1}{2\sqrt{y}} \cdot \frac{4}{9} = \frac{2}{9\sqrt{y}}$.
For $1 < y < 4$: only $x = \sqrt{y}$ is in domain. $g(y) = \frac{f(\sqrt{y})}{2\sqrt{y}} = \frac{2(\sqrt{y}+1)}{9 \cdot 2\sqrt{y}} = \frac{\sqrt{y}+1}{9\sqrt{y}}$.

Exercise 7.18

Problem: $X \sim \text{Geometric}(p)$: $P(X = x) = pq^{x-1}$. Show MGF is $M_X(t) = \frac{pe^t}{1-qe^t}$ for $t < \ln q$. Find mean and variance.
Solution: $M_X(t) = E(e^{tX}) = \sum_{x=1}^{\infty} e^{tx} pq^{x-1} = \frac{p}{q} \sum_{x=1}^{\infty} (qe^t)^x = \frac{p}{q} \cdot \frac{qe^t}{1-qe^t} = \frac{pe^t}{1-qe^t}$.
$M'_X(t) = \frac{pe^t}{(1-qe^t)^2}$. At $t = 0$: $E(X) = \frac{p}{(1-q)^2} = \frac{p}{p^2} = \boxed{1/p}$.
$M''_X(0) = \frac{p(1+q)}{p^3} = \frac{1+q}{p^2}$. $\text{Var}(X) = E(X^2) - [E(X)]^2 = \frac{1+q}{p^2} - \frac{1}{p^2} = \boxed{q/p^2}$.

Exercise 7.21

Problem: Show that the MGF of $\chi^2(\nu)$ is $M_X(t) = (1 - 2t)^{-\nu/2}$.
Solution: $\chi^2(\nu)$ is Gamma$(\alpha = \nu/2, \beta = 2)$. The MGF of Gamma$(\alpha, \beta)$ is $(1 - \beta t)^{-\alpha}$. Substituting: $M_X(t) = (1 - 2t)^{-\nu/2}$. $\square$

Exercise 7.23

Problem: X, Y iid Geometric(p) with $P(X = r) = q^r p$, $r = 0, 1, 2, \ldots$ Find (1) distribution of $U = X + Y$, (2) conditional distribution of $X/(X + Y) = u$.
Solution: (1) $P(U = n) = \sum_{r=0}^{n} P(X = r)P(Y = n - r) = \sum_{r=0}^{n} q^r p \cdot q^{n-r} p = (n + 1)p^2 q^n$. This is a Negative Binomial$(k = 2, p)$ distribution.
(2) $P(X = r \mid X + Y = n) = \frac{P(X=r, Y=n-r)}{P(X+Y=n)} = \frac{p^2 q^n}{(n+1)p^2 q^n} = \frac{1}{n+1}$. This is a discrete uniform distribution on $\{0, 1, \ldots, n\}$.