

# Hyperbolic string art

Elliot Kienzle<sup>1</sup>

<sup>1</sup>Mathematics, University of California, Berkeley; ekienzle@berkeley.edu

## Abstract

*String art* are emergent patterns arising by stretching straight lines across a circle according to mathematical rules. We re-contextualize string art, envisioning the circle as the circle at infinity of the hyperbolic plane. The strings stretch across the Beltrami-Klein model of the hyperbolic plane, each line a hyperbolic geodesic. We examine the string art coming from Möbius transforms. The string art is described by its envelope, a curve tangent to every string stretching across a circle. We describe the envelopes of a Möbius transform in terms of the underlying hyperbolic symmetry: Elliptic Möbius transforms give hyperbolic circle envelopes, parabolic transforms give horocycles, and hyperbolic transforms give hypercircles. We conclude with a purely aesthetic application, showing a hyperbolic string art fractal.

## Introduction

*String art* (or *curve stitching*) is an art form composed of thin stings stretched between points. Placing the strings according to mathematical rules results in aesthetic designs that illustrate the mathematics of the generating rule. For example, in a classic classroom activity, students place 12 equally spaced pins around a circle, labeled 1 to 12. Each student stretches strings from the pin labeled  $i$  to that labeled  $i + 3$  modulo 12 (See Figure 1a). The strings accumulate around a smaller central circle, manifesting the rotational symmetry of modular addition. The smaller circle is a curve built from straight lines. It is tangent to each string and “envelopes” the family of strings, so is called the *envelope*. Modular multiplication is similarly illustrated by stretching strings from  $i$  to  $2i$ . The strings once again accumulate around their envelope, the heart-shaped cardioid (Figure 1b).


Unfortunately, the exercise seldom goes much further. Other usual functions (squaring, exponential, etc.) produce noisy results without an envelope (see Figure 1c). These functions are not natural with respect to the modular structure, so their string art lacks obvious mathematical or aesthetic patterns. So, We change perspective, replacing the integers labeling the pegs with their angle around the circle. The strings are defined by maps from the circle to itself. Modular addition (Figure 1a) is a rigid rotation, while modular multiplication by two (Figure 1b) is the angle doubling map. To make string art, choose a set of input points, and stretch a line from each point to its image under the defining map. This viewpoint suggests several other natural functions, like sine and cosine, which produce aesthetic envelopes (See Figure 2).

In this paper, we explore a further leap in perspective, exploring the geometry of the disc that the strings stretch across. We introduce *Hyperbolic string art*, which identifies the disc with the hyperbolic plane. Strings are straight lines across the disc, representing geodesics (hyperbolic lines<sup>1</sup>) in the Beltrami-Klein model of the hyperbolic plane. The bounding circle is now hyperbolic infinity (or “the circle at infinity”) <sup>2</sup>, the set of possible directions a geodesic can point. Hyperbolic geometry suggests several natural functions mapping the circle at infinity to itself. In particular, symmetries of the hyperbolic plane extend as Möbius transforms to the circle at infinity. We will explore the string art coming from these maps, understanding their envelopes and relating them to the underlying symmetries of the hyperbolic plane. Through this, string art artistically visualizes properties and themes of hyperbolic geometry. We also show purely artistic applications, such as fractal string art.


---

<sup>1</sup>In this paper, we will refer to hyperbolic lines as geodesics, and Euclidean lines as lines

<sup>2</sup>hyperbolic infinity is sometimes called the set of *ideal points*



**Figure 1:** String art illustrating modular arithmetic.




**Figure 2:** String art for the function  $\theta \rightarrow 2(\theta + \frac{1}{2} \sin(\theta))$

### String art in the hyperbolic plane

Let us first recap the necessary aspects of hyperbolic geometry (For reference, see chapter 2 of [1]). The hyperbolic plane is a two-dimensional plane with constant negative curvature. Like the Euclidean plane, there are lines, angles, distances, and an axiomatic approach to geometry analogous to Euclid's. However, the negative curvature forces geodesics to curve away from one another, possibly never intersecting. This is a failure of Euclid's fifth postulate, as there are infinitely many geodesics thru a given point, parallel to a given geodesic. Despite this, we can still draw the hyperbolic plane on Euclidean paper using *models*. These are akin to flat maps of the spherical earth, and like maps, every model has its own uses. Let us start with the *Beltrami-Klein model*, where the hyperbolic plane is drawn as the interior of a unit disc, and geodesics are drawn as straight lines. A fixed hyperbolic distance looks smaller the closer it is to the boundary of the disc, vanishing at the boundary. In fact, the bounding circle is infinitely far from any point inside the disc, hence its moniker "the circle at infinity".

Note that every line passing thru the interior of the unit disc intersects the boundary circle twice, and is uniquely defined by its intersection points. Translating to hyperbolic geometry, each geodesic is uniquely defined by its two asymptotic directions, each a point at hyperbolic infinity. This manifests a creed of hyperbolic geometry: *an object in the hyperbolic plane is uniquely determined by its behavior at hyperbolic infinity*. This stands in contrast to Euclidean space, where the forward and backward directions of a line are



**Figure 3:** String art of a Möbius transform, shown in two different models of hyperbolic space.

necessarily opposite, and many lines share the same asymptotic direction. The core insight of hyperbolic string art is to treat the boundary of the disc as hyperbolic infinity and to treat strings as hyperbolic geodesics.


The hyperbolic plane carries similar symmetries (or isometries) to the Euclidean plane, including rotation around a point and translation along a direction. Unlike Euclidean space, hyperbolic translations and rotations mix together, as translation along a loop results in a net rotation. This mixing is captured in the symmetry group of the hyperbolic plane  $SL(2, \mathbb{R})$ , which consists of  $2 \times 2$  real matrices with determinant one. Hyperbolic symmetries send geodesics to other geodesics, hence naturally acts on the space of asymptotic directions. They restrict to *Möbius transforms* on the circle at infinity. In fact, hyperbolic symmetries are uniquely defined by their action at infinity, manifesting the above hyperbolic creed.

Let us form string art from a hyperbolic symmetry. First, we choose a symmetry, such as rotation around a point. This acts on the circle at infinity via a Möbius transform. Then we draw the string art for this map, shown in Figure 3a. The envelope forms an ovoid shape, surrounding the pivot point of the rotation.

However, the Beltrami-Klein model obfuscates hyperbolic shapes, because it distorts angles. To understand the envelope's shape, we switch to the *Poincare disc model*, which preserves hyperbolic angles. The Poincare disc model is based on the unit disc, but geodesics are drawn as circular arcs intersecting the boundary circle at right angles. Using these circle arcs to render the strings from Figure 3a results in Figure 3b. In the Poincare disc model, the envelope appears to be a perfect circle, and the center of the envelope circle is the original pivot point of the hyperbolic rotation. This is not a coincidence. In the next section, we will explain why the envelope is a hyperbolic circle, and discuss the envelopes for all other types of Möbius transforms.

### Möbius transform taxonomy through string art

Symmetries of the hyperbolic plane fall into three classes: Rotations, translations, and parabolic transforms. Their associated Möbius transforms are called elliptic, hyperbolic, and parabolic respectively. These are illustrated in Figure 4: The Orange arrows circumnavigating the border circles show how the Möbius transform acts on each point. An elliptic Möbius transform has no fixed points at infinity and comes from a hyperbolic rotation about the point marked in purple. Moving to column 2, a parabolic Möbius transform has one fixed point at infinity, shown by the orange dot. The corresponding symmetry of the hyperbolic plane




**Figure 4:** Description of string art for each type of Möbius transform.

does not have a description familiar to Euclidean beings. Shown in column 3, a hyperbolic Möbius transform pulls points at infinity away from a repelling fixed point, and towards an attracting fixed point. This is the boundary action of hyperbolic translation along a geodesic which starts at the repelling point and ends at the attracting point called the *axis of translation* (marked in purple).

Algebraically, this trichotomy arises from the conjugacy classes of the associated elements of  $SL(2, \mathbb{R})$ . Each conjugacy class is uniquely defined by its trace. When the trace is  $> 2$  (resp.  $= 2, < 2$ ), the associated Möbius transform is elliptical (resp. parabolic, hyperbolic). Geometrically, this represents symmetries of the hyperbolic plane up to change in perspective. All hyperbolic rotations by  $\theta$  degrees are conjugate to one another by a translation: Translate one pivot point to another, then rotate by  $\theta$ , then translate back.  $\theta$  defines a conjugacy class and is uniquely determined by the trace. Similarly, hyperbolic translations are conjugate by rotations, and the only invariant under conjugacy is the length of translation. Finally, all parabolic transforms are conjugate. Each class of Möbius transforms can be understood from a single family of basic transforms, such as rotation around a fixed point or translation along a fixed line. We will use this technique to study the string art of Möbius transforms. These are shown in Figure 4, with their envelopes shown in green. The descriptions of each envelope are defined and justified below.

### ***Elliptic Möbius transforms***

As mentioned above, elliptic Möbius transforms all arise from hyperbolic rotations. We start with the simplest rotation, pivoting around the center of the Poincaré disc. This acts on the boundary circle by rigid rotation. All the strings are thus circular arcs of the same radius. The envelope is a circle with radius equal to the smallest distance between one string and the center. To see this, consider the distance function to the center of the Poincaré disc. At the minimum distance along one string, the string must be tangent to the level set of the distance function, which is a circle (this is a geometric realization of the method of Lagrange multipliers). By rotational symmetry, every string has the same minimal distance, so all strings are tangent to the same



**Figure 5:** String art for a parabolic Möbius transform in the Poincaré disc (left) and half-plane (right) models. The rainbow boundary is hyperbolic infinity, the red curves are geodesic strings, the green curve is the envelope of the strings, and the white shows normal geodesics to the envelope.

level set. Hence, the envelope is a circle. Note that this argument also justifies the circular envelope in string art for modular addition, shown in Figure 1a. The Poincaré projection preserves angles, so hyperbolic circles are drawn as Euclidean circles. Thus, the envelope is the image of a hyperbolic circle.

Now we extend the simple case of rigid rotation  $R$  to all elliptic Möbius transforms. Let us translate the string art of rigid rotation by a hyperbolic translation  $T_p$  sending the center of the Poincaré disc to a point  $p$ . Since  $T_p$  is a hyperbolic symmetry, it sends geodesics to geodesics. After translating, the geodesics form string art from a new function  $f$ , which maps the start of each translated geodesic to its endpoint. Symbolically, if  $d$  is a point at hyperbolic infinity, the function  $f$  maps  $T_p(d)$  to  $T_p(R(d))$ . Labeling  $T_p(d)$  as  $d'$ , we see

$$f : d' \rightarrow T_p R T_p^{-1}(d')$$

So, the image is the string art for the Möbius transform  $T_p R T_p^{-1}$ , which is a hyperbolic rotation around the point  $p$ . Every elliptic Möbius transform is conjugate to  $R$  in this way.

The image under the translation  $T_p$  of the envelope for rigid rotation  $R$  gives the envelope for rotation around  $p$ ,  $T_p R_0 T_p^{-1}$ . However, the envelope for  $R$  is a hyperbolic circle. The translation  $T_p$  is an isometry, so it sends hyperbolic circles to hyperbolic circles. Thus, *The envelope of any elliptic Möbius transform is a hyperbolic circle*. This is drawn in the Poincaré disc as a Euclidean circle, as seen in column 1 of Figure 4.

### Parabolic Möbius transforms

We establish the string art for parabolic transforms following the blueprint of the above argument. First, we choose a specific model of hyperbolic space and specific parabolic transform with a very simple description. In particular, it will be realized by a Euclidean isometry of the model. Then, we find the envelope and describe it intrinsically with hyperbolic geometry. Finally, we describe the string art of all other parabolic Möbius transforms by conjugating the simple case. We notice that our intrinsic description of the envelope is preserved by isometries, and conclude that all envelopes of parabolic transforms share the same intrinsic description.

We will describe our base parabolic transform in the Poincaré half-plane model, shown in the right half of Figure 5. This consists of the complex numbers with imaginary part  $> 0$ , endowed with a constant

negative curvature metric which scales down distances near the real line (visualized in Figure 5 by the scale of the background pattern). This metric is invariant under translation by a real number. Hyperbolic infinity consists of the real line (drawn in rainbow), along with the point at infinity in the complex plane. Geodesics consist of semicircles intersecting the real line orthogonally. Möbius transforms act on hyperbolic infinity by fractional linear transforms:

$$A = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{R}) \quad A : x \rightarrow \frac{ax + b}{cx + d}$$

All parabolic Möbius transforms are conjugate to those of the form  $x \rightarrow x + 1$ , which are induced by a translation of the complex plane.

The strings for this action are semicircles stretching from  $x$  to  $x + 1$  on the real line (some examples are drawn in red in Figure 5). The envelope is a horizontal line at height 1/2 (drawn in green): Indeed, every string has maximal height 1/2, so the level set of the height function at 1/2 is tangent to every string. Returning to the hyperbolic plane, this horizontal line is a *Horocycle*: A curve whose perpendicular geodesics all converge to the same point at hyperbolic infinity. Indeed, the perpendicular geodesics of a horizontal line are vertical lines (infinitely large semicircles, drawn in white), which converge at infinity in the half-plane model. Convergence is clear after converting back to the Poincare disc model, where the horocycle is drawn as a circle tangent to the top of the boundary circle (See left half of Figure 5).

All parabolic Möbius transforms are conjugate to that described above. Following the argument for elliptic transforms, the string art of every parabolic transform is related by hyperbolic symmetries. These symmetries preserve horocycles, so *the envelope of a parabolic Möbius transform is a horocycle*. In general, horocycles are drawn in the Poincare disc model as circles tangent to the boundary circle, as seen in column two of Figure 4.

### *Hyperbolic Möbius transforms*


For hyperbolic Möbius transforms, we follow the same strategy as above. We describe the base hyperbolic translation in the *band model*, shown on the right side of Figure 6. This consists of the strip of complex numbers  $z$  with  $\text{Im } z \in (-1, 1)$ , endowed with a constant negative curvature metric that scales distances to zero at the top and bottom line (visualized again by the scaling of the background pattern). The metric is translation invariant under  $z \rightarrow z + b$  for real  $b$ . Hyperbolic infinity consists of top and bottom lines  $\text{Im } z = \pm 1$ , along with the two points at infinity of the real line, denoted  $\pm\infty$ . A geodesic stretching between two points on the same boundary line is not a semicircle, but is a convex shape that does not cross the real axis (some examples are drawn in red). In particular, each geodesic has a unique point closest to the middle of the band.

Every hyperbolic translation is conjugate to a Euclidean translation  $z \rightarrow z + b$  in the band model. The fixed points of this action are  $\pm\infty$ , and the geodesic connecting the two fixed points is the real axis, drawn in purple in Figure 6.<sup>3</sup> The string art consists of geodesics connecting  $z$  with  $z + b$  with  $\text{Im}(z) = \pm 1$ . Every string is the same shape, so has the same minimal distance to the real axis. Hence, the envelope consists of two parallel lines equidistant to the real axis (drawn in green). This envelope is the locus of points of fixed hyperbolic distance to the geodesic connecting  $\pm\infty$ , a shape called a *hypercircle*.

Converting back to the Poincare disc model (Figure 6, left), this hyperbolic Möbius transform has fixed points at the left and right extremes of the unit disc, corresponding to  $-\infty$  and  $+\infty$  respectively. The geodesic connecting these two points is again the real axis. The hypercircle is drawn as two circular arcs connecting the fixed points at infinity of the translation, mirror to one another about the real axis, as depicted in green. In

---

<sup>3</sup>Compare this to the parabolic case, where the parabolic transform was realized as a Euclidean translation with a single fixed point at Euclidean infinity. The band model realizes a hyperbolic transform as a Euclidean translation, with two fixed points at Euclidean infinity.



**Figure 6:** String art for a hyperbolic Möbius transform in the Poincare disc model (left) and the band model (right). The rainbow boundary is hyperbolic infinity, the red curves are geodesic strings, the green curve is the envelope of the strings, and the purple geodesic is the axis of the hyperbolic translation.

general, hypercircles about a given geodesic in the Poincare disc are drawn as a pair of circular arcs sharing the endpoints of that geodesic. See the string art for the hyperbolic Möbius transform in Figure 4.

Once again, we realize the string art of all hyperbolic Möbius transforms by applying a hyperbolic symmetry to the string art above. Since hyperbolic symmetries preserve the hyperbolic distance to geodesics, the image of the envelope is still a hypercircle. Hence, *the envelope of a hyperbolic Möbius transform is a hypercircle*.<sup>4</sup> The only conjugacy invariant of hyperbolic translations is the translation amount, which controls the angle of the envelope at infinity. For example, small translations will yield strings that stay far from the translation axis, so the envelope will have a wide angle.

### Generalities about Möbius transforms

In this section, we argued that an elliptic (resp. parabolic, hyperbolic) Möbius transform yields string art with an envelope forming a hyperbolic circle (resp. horocycle, hypercircle). This is summarized in Figure 4. The arguments above can be formalized into a proof without much trouble. For an intrinsic proof that does not rely on specific models of hyperbolic geometry, see the extended online version of this paper [2]. In the Poincare disc model, all three envelope types share a uniform description:

- Elliptic transforms have circular envelopes that remain within the boundary circle.
- Parabolic transforms have circular envelopes tangent to the boundary circle.
- Hyperbolic transforms have circular envelopes intersecting the boundary circle.

The points of intersection with the boundary circle are the fixed points of the transform. The origin of the two-component envelope in hyperbolic transforms is clarified by extending the strings to full circles (as in Figure 4). Then, the envelope of any transform has two circular components, one being the inversion of the other about the boundary circle. For elliptic and parabolic transforms, this second envelope lies outside the disc, but for hyperbolic transforms both envelopes intersect the disc.

---

<sup>4</sup>If the transform is not orientation-preserving, then the string art does not have an envelope


## Aesthetic Applications

On top of illustrating mathematics, we can pick string art functions purely for their artistic merit. Figure 7 shows string art inspired by the Mandelbrot set, a fractal built from the repeated dynamics of the map  $z \rightarrow z^2 + c$ . This squares a complex number and then applies a parabolic Möbius transform (shifting by  $c$ ). Restricting from the Riemann sphere to the unit circle,  $z \rightarrow z^2$  multiplies angles by 2, and shifting by  $c$  is a Möbius transform. A hyperbolic analog is the map  $\theta \rightarrow 2 \text{ Möbius}(\theta)$ . Figure 7 shows the string art for this function, iterated various numbers of times.

## Summary and Future Work

Hyperbolic string art recontextualizes string art in the framework of hyperbolic geometry. In this paper, we explored only symmetries, the simplest functions. These produced envelopes with natural hyperbolic descriptions, reflecting the symmetry of the hyperbolic plane. String art visualizes Möbius transforms and hyperbolic geometry in a novel, artistic way. However, hyperbolic geometry cares about more than mere symmetries.

Maps from infinity to itself capture information about hyperbolic tessellations, as a deformed hyperbolic lattice characteristically deforms infinity. Circle maps thus parameterize all hyperbolic lattices, and hence all hyperbolic surfaces [1]. We hope for string art to visualize the boundaries of this space, elucidating the appearance of geodesic laminations. The space of circle maps is also a core object in, ironically enough, string theory: Perhaps one day, we could use string art to visualize string theory.



**Figure 7:** String art for iterations of the map  $\theta \rightarrow 2 \text{ Möbius}(\theta)$ . From the top down, we depict 1, 3, 5, and 7 iterations.

## Acknowledgements

Thanks to Chaim Goodman-Strauss, Fran Herr, Sabetta Matsumoto, and Henry Segerman for helpful discussions. All string art was generated using a web app <https://chessapig.github.io/sketch/strings>. The app was coded in P5.js, a JavaScript library designed for and creative coding.

## References

- [1] J. Hubbard. *Teichmüller Theory: And Applications to Geometry, Topology, and Dynamics*. Matrix Editions. no. v. 1. <https://books.google.com/books?id=azpNswEACAAJ>.
- [2] E. Kienzle. “Hyperbolic string art (extended).” [https://chessapig.github.io/files/hyperbolic\\_string\\_art.pdf](https://chessapig.github.io/files/hyperbolic_string_art.pdf).