

# Antiandrogenic Effect of Spirolactones: Mechanism of Action

PIERRE CORVOL, ANNIE MICHAUD, JOËL MENARD, MARTIN FREIFELD,  
AND JACQUES MAHOUDEAU

INSERM U-36 75005 Paris, France

**ABSTRACT.** Spirolactones are aldosterone antagonists which inhibit the binding of aldosterone to the renal mineralocorticoid receptor. These molecules also possess an antiandrogenic effect which could be due, among other possibilities, to a peripheral antagonism of androgens. This hypothesis has been tested in the present study. From *in vivo* experiments, spironolactone and K<sup>+</sup> canrenoate appear to inhibit the binding of [<sup>3</sup>H]5 $\alpha$ -dihydrotestosterone [<sup>3</sup>H]DHT to the cytosolic and nuclear receptor of the rat ventral prostate. The doses used are in the same range as those used for demonstrating the antimineralocorticoid effect of

these molecules. *In vitro* incubations and *in vitro* displacement studies show that spironolactone and K<sup>+</sup> canrenoate are respectively about 20 and 100 times less effective than DHT in displacing 50% of  $5 \times 10^{-10}$  M [<sup>3</sup>H]DHT from its receptor. Spirolactones are also able to compete with [<sup>3</sup>H]DHT for the specific 8 S cytosolic receptor. Neither spironolactone nor K<sup>+</sup> canrenoate decreases prostatic 5 $\alpha$ -reductase activity, even at a concentration as high as  $10^{-5}$ M. It seems likely that spirolactones, besides their action on testosterone biosynthesis, exert their antiandrogenic activity via a peripheral androgen antagonism. (*Endocrinology* 97: 52, 1975)

**S**PIROLACTONES are aldosterone antagonists which are currently used in the treatment of primary hyperaldosteronism and essential hypertension. Kagawa first showed the antimineralocorticoid effect of spirolactones in adrenalectomized rats treated by deoxycorticosterone acetate (1). Spirolactones act *in vitro* by competitively inhibiting active Na<sup>+</sup> transport induced by aldosterone (2). At the molecular level, spirolactones have been shown to inhibit the first step of the mechanism of action of aldosterone, that is, its binding to kidney cytosol mineralocorticoid receptor (3–6).

One of the side effects most frequently encountered during chronic spirolactone therapy is the occurrence of sexual disorders induced by these drugs. Several mechanisms could account for these effects: Menard *et al.* (7) have recently reported that spironolactone decreased the activity of the 17 $\alpha$ -hydroxylase and the content of cytochrome P-450 in the rat testis resulting in an impairment in testosterone biosynthesis.

Another possible mechanism could be a peripheral antiandrogenic effect of spirolactones at the level of the target organs.

Such a mechanism is suggested by the experiments of Steelman *et al.* (8) and Rasmussen *et al.* (9) who have demonstrated that spironolactone antagonizes the androgen-stimulated weight gain of the seminal vesicle and ventral prostate. Such an androgen antagonism could be due either to the inhibition of prostatic 5 $\alpha$ -reductase activity or to competition with 5 $\alpha$ -dihydrotestosterone<sup>1</sup> (DHT) for the androgenic cytosolic receptor. The purpose of this study was to investigate these two possible effects of spirolactones on prostatic tissue.

## Materials and Methods

### Animals

Adult male Wistar rats (200 g) were castrated by the scrotal route 24 h before sacrifice. On the day of experiment, the prostate was removed and rinsed with isotonic saline at 4 C. [1,2-<sup>3</sup>H]5 $\alpha$ -Dihydrotestosterone (<sup>3</sup>H-DHT), 47 Ci/mmol, and [1,2-<sup>3</sup>H]testosterone, 48 Ci/mmol, were purchased from Amersham and their purity checked

<sup>1</sup> The terminology used is: 5 $\alpha$ -dihydrotestosterone (DHT), 5 $\alpha$ -androstane-17 $\beta$ -ol-3-one; cyproterone acetate, 6 $\alpha$ -chloro-17 $\alpha$ -hydroxy-1 $\alpha$ ,2 $\alpha$ -methylene-4,6-pregnadiene-3,20-dione-17-acetate.

Received July 19, 1974.

before use as previously described (10). Spironolactone (SC 9420) and  $K^+$  canrenoate (SC 14266) were kindly given by Searle, and  $5\alpha$ -dihydrotestosterone (Ikapharm) was recrystallized before use. Pure sucrose was obtained from Sigma and all other chemical products were analytical grade and purchased from Merck (Germany), unless otherwise indicated.

#### *In vivo experiments*

There were 3 rats in each experimental group. Animals in the control group (group I) were injected intraperitoneally with 0.5 ml of a mixture of ethanol, propanediol and physiologic saline. Rats in the other groups received either 200  $\mu$ g of DHT (group II), 5 mg of spironolactone (group III) or 5 mg of  $K^+$  canrenoate (group IV), dissolved in the same mixture. One hour later, all rats were injected intravenously with 20  $\mu$ Ci ( $4.3 \times 10^{-10}$  M) of [ $^3$ H]DHT. Fifteen minutes after the last injection, the animals were killed and the prostate glands removed and homogenized in 4 ml of Tris HCl buffer 0.01M, EDTA 1 mM, pH 7.4, containing 10% glycerol and 0.05 mM of dithiothreitol. The homogenate was centrifuged at  $700 \times g$  for 15 min and the cytosol ( $105,000 \times g$ ) was prepared from the supernatant. The pellet obtained from the low speed centrifugation of the homogenate was resuspended in 2.2M sucrose and the nuclei purified by centrifugation for 1 h at  $78,000 \times g$  on 2.2M sucrose. The bound and free radioactivity in the cytosol were separated using a dextran-charcoal method: 200  $\mu$ l of cytosol were incubated for 90 min under agitation with 500  $\mu$ l of dextran-coated charcoal (dextran 0.5%, charcoal 5%) and centrifuged at  $2,500 \times g$  for 25 min; the supernatant was then counted. The nuclear pellet was resuspended in 2 ml of 0.4M KCl for 15 min, then centrifuged at  $20,000 \times g$  for 20 min. An aliquot of the supernatant was taken for radioactivity measurement and another aliquot was taken for protein determination by the method of Lowry *et al.* (11).

#### *In vitro incubations*

Prostate slices (280  $\mu$ m) were incubated with  $5 \times 10^{-9}$  M [ $^3$ H]DHT with or without competitors at 25 C for 25 min in incubation buffer A:  $Na^+ = 133$ ,  $K^+ = 6$ ,  $Ca^{++} = 1$ ,  $Mg^{++} = 0.5$ ,  $Cl^- = 134$ ,  $H_2PO_4^- = 6$ , Tris HCl = 5, and glucose = 5 mM, pH 7.4. At the end of the incubation, the slices were homogenized at 4 C

in Tris-HCl buffer, 0.01M, pH 7.4, containing 1 mM EDTA and 10% glycerol. The cytosol and the nuclei were then prepared as described above. Bound and free radioactivity in the cytosol were separated by dextran-charcoal and the nuclear radioactivity was extracted and counted as already described.

#### *In vitro displacement studies*

Twelve rat prostates were homogenized in 12 ml of Tris-HCl buffer, 0.05M, pH 7.4, containing 1 mM EDTA and 0.1 mM mercaptoethanol. The homogenate was then centrifuged at  $1,000 \times g$  for 15 min and the resulting supernatant was incubated at 4 C for 180 min with  $5 \times 10^{-10}$  M [ $^3$ H]DHT in the presence or the absence of various competitors (DHT, spironolactone,  $K^+$  canrenoate, cyproterone acetate<sup>1</sup>). Bound [ $^3$ H]DHT was separated from free steroid by the dextran-charcoal technique.

Linear sucrose gradients (5–20%) were prepared in the homogenization buffer containing 10% glycerol; 0.2 ml cytosol samples were then layered at the top of the gradients. Ultracentrifugation, fraction collection and the use of standard proteins were performed as already described (12).

#### *Effects on $5\alpha$ -reductase activity*

Ventral prostates were removed from ether-anesthetized 200–250 g rats. Prostatic tissue was separated from connective tissue, then minced with scissors and pooled in cold buffer A (4 C). About 150 mg of tissue was incubated at 37 C for 2 h under oxygen in 2 ml of buffer A, pH 7.4, containing 2 U glucose-6-phosphate dehydrogenase, 10 pmol glucose-6-phosphate and 2.5 pmol TPN (Sigma) per incubation vial. Incubations were performed with  $1.7 \times 10^{-9}$  M [ $^3$ H]testosterone with or without  $10^{-7}$  to  $10^{-5}$  M spironolactone or  $K^+$  canrenoate. After addition of 100  $\mu$ g testosterone, 100  $\mu$ g DHT and [ $^{14}$ C]DHT tracer for recovery, the tissue was homogenized and the steroids were extracted and purified by thin layer chromatography (TLC): benzene, then benzene: methanol (95:5). After acetylation, DHT acetate was chromatographed by TLC in benzene: ethyl acetate (90:10). The  $^3H/^{14}C$  ratio was calculated.

#### *Radioactivity measurements*

Aqueous samples were counted after addition of 10 ml of Unisolve (Koch and Light) in a

TriCarb Scintillation spectrometer 3380 with an efficiency of 34%.  $^{14}\text{C}$ - and  $^3\text{H}$ -labeled compounds eluted from thin-layer plates were counted in a standard counting solution (toluene, 960 ml. Liquifluor (New England Nuclear), 40 ml).

## Results

### *In vivo experiments*

Fifteen minutes after *in vivo* injection of  $4.3 \times 10^{-10}$  M [ $^3\text{H}$ ]DHT, radioactivity was found in the cytosol and nuclei of rat ventral prostate. The specifically bound radioactivity in cytosol was measured after removal of free radioactivity by dextran-charcoal. Displacement experiments showed that 200  $\mu\text{g}$  DHT and 5 mg spironolactone could equally displace the bound radioactivity in both cytosol and nuclei (Fig. 1).  $\text{K}^+$  canrenoate (5 mg) also inhibited the binding of the radioactivity to the cytosolic and nuclear receptor.

### *In vitro displacement studies*

Very similar results were obtained by using *in vitro* incubation of rat ventral prostate slices (Table 1). A 5,000-fold ex-

TABLE 1. Binding of [ $^3\text{H}$ ]DHT in cytosol and nuclei during *in vitro* incubation of rat ventral prostate slices in the presence of competitors

Steroids	Cytosol dpm/mg protein	Nuclei dpm/mg protein
$5 \times 10^{-9}$ M [ $^3\text{H}$ ]DHT	4,580 4,183	39,928 39,133
+ $2.5 \times 10^{-6}$ M DHT	1,195 1,077	11,390 9,542
+ $2.5 \times 10^{-5}$ M Spironolactone (SC 9420)	1,223 1,038	9,661 10,041
+ $2.5 \times 10^{-5}$ M $\text{K}^+$ Canrenoate (SC 14 266)	2,211 2,570	17,381 20,948

cess of spironolactone displaced 75% of the bound radioactivity as did 500-fold excess of unlabelled DHT. In this experiment, as for the *in vivo* experiment,  $\text{K}^+$  canrenoate appeared to be less potent than spironolactone since a 5,000-fold excess of this compound displaced only 50% the bound radioactivity.

Figure 2 shows complete *in vitro* studies: the  $1,000 \times g$  supernatant was incubated with  $5 \times 10^{-10}$  M [ $^3\text{H}$ ]DHT and various competitors. Fifty per cent displacement of the bound radioactivity was obtained by  $3 \times 10^{-8}$  M spironolactone and




FIG. 1. Effect of DHT and of spironolactones on binding of radioactivity in prostatic cytosol and nuclei after injection of [ $^3\text{H}$ ]DHT in orchidectomized rats. Rats were injected ip with solvent (control group), 200  $\mu\text{g}$  DHT, 5 mg of spironolactone (SC 9420) or 5 mg  $\text{K}^+$  canrenoate (SC 14266). One hour later the rats were injected iv with 10  $\mu\text{Ci}$  [ $^3\text{H}$ ]DHT and killed 15 min later. The specifically bound radioactivity in the prostatic cytosol was counted after separation of bound from free steroid by the dextran-charcoal method. Nuclear bound radioactivity was counted after 0.4M KCl extraction. Each bar represents the mean of two experiments. There were 3 rats in each experimental group. De-

termination of cytosolic radioactivity was performed in quadruplicate (mean  $\pm$  SD) and nuclear radioactivity was measured in duplicate.

$4 \times 10^{-6}$  M  $K^+$  canrenoate. Cyproterone acetate produced about the same displacement of  $[^3H]$ DHT as spironolactone. When cytosol incubated with  $5 \times 10^{-9}$  M of  $[^3H]$ DHT was analyzed by sucrose gradient centrifugation,  $[^3H]$ DHT was specifically bound to an 8 S protein peak (Fig. 3). This peak was displaced by a 1,000-fold excess of unlabelled DHT or a 10,000-fold excess of spironolactone.  $K^+$  canrenoate (10,000-fold excess) could also displace  $[^3H]$ DHT from its 8 S binding peak.

#### *Effect of spiro lactones on $5\alpha$ -reductase activity*

Finally, the possible effect of spironolactone and  $K^+$  canrenoate on the  $5\alpha$ -reductase activity of rat ventral prostate was tested. As shown in Table 2, neither spironolactone nor  $K^+$  canrenoate decreased  $5\alpha$ -reductase activity of rat ventral

TABLE 2. Effect of spiro lactone on prostatic  $5\alpha$ -reductase

Steroids (M)	mg Prostatic tissue/incubation	% of Radioactivity recovered as $[^3H]$ DHT/incubation
Control ( $1.7 \times 10^{-9}$ M $[^3H]$ testosterone)	$164 \pm 24^*$	$48 \pm 6^*$
Spironolactone (SC 9420) $10^{-7}$	151 141	49 47
$10^{-6}$	147 170	49 45
$10^{-5}$	148 139	46 48
$K^+$ Canrenoate (SC 14 266) $10^{-7}$	115 158	36 45
$10^{-6}$	89 122	47 52
$10^{-5}$	144 156	42 45

\* Mean of 4 values  $\pm$  SD.




FIG. 2. *In vitro* incubation of prostatic supernatant with  $5 \times 10^{-10}$  M  $[^3H]$ DHT in the presence of various competitors. Bound DHT was separated from free DHT by the dextran-charcoal technique. The results are expressed as a percent of control. Each point is the mean of 3 experiments.

prostate, even at concentrations as high as  $10^{-5}$  M.

#### Discussion

Ever since the clinical introduction of spiro lactones in diuretic therapy, there have been reports of associated sexual disorders. Gynecomastia is a common complication in spironolactone treated men, as well as decreased libido (13-16). Menstrual cycle abnormalities are very often observed in females (16). All of these effects disappear on withdrawal of the drug.

Spiro lactones can interfere at different levels with the metabolism of androgens. Dymling *et al.* (17) have shown that 100 to 200 mg of  $K^+$  canrenoate iv was able to lower the plasma level of testosterone significantly in normal men. Such an effect could be due to a change in testosterone metabolism or to reduced testicular secretion of testosterone. In favor of this latter hypothesis is the recent report of Menard




FIG. 3. Sedimentation patterns of prostatic cytosol after incubation with  $5 \times 10^{-9}$  M [ $^3\text{H}$ ]DHT in the absence or in the presence of  $5 \times 10^{-6}$  M DHT or  $5 \times 10^{-5}$  M spironolactone. Five to 20% linear sucrose gradients were made in 0.05M Tris-HCl, 1.0 mM EDTA, 0.1 mM mercaptoethanol, pH 7.4, containing 10% glycerol. Samples were centrifuged at 55,000 rpm for 15 h at 2 C.

*et al.* (7) which demonstrates that daily treatment of male rats with spironolactone (200 mg/kg) for three days induces a 72% decrease in the *in vitro* formation of testosterone from progesterone.

However other mechanisms could account in part for the antiandrogenic effect of spironolactones. Steelman *et al.* (8) and Rasmusson *et al.* (9) have shown that spironolactone can antagonize the peripheral

action of androgens in the castrate rat. One to 3 mg of spironolactone was able to significantly decrease the seminal vesicle and ventral prostate weights in the immature male castrate rat treated with a single dose of 0.5 mg of testosterone enanthate (9).

The aim of the present studies was to investigate further the peripheral antiandrogenic properties of spironolactones. Evi-

dence has been presented that spirolactones are able to compete *in vivo* and *in vitro* with DHT for the intracellular androgen receptor sites. *In vitro* experiments confirmed that both spironolactone and  $K^+$  canrenoate were able to decrease the formation of cytosolic- and  $[^3H]$ DHT-chromatin-receptor complexes. That spironolactone was effectively able to compete with  $[^3H]$ DHT for the specific 8 S cytosolic receptor (18,19) was shown by the sucrose gradient experiment.

The doses used in the current study for demonstrating the inhibitory effect of spironolactone on DHT binding are similar to those used for proving its antiandrogenic and antimineralocorticoid activity. Inhibition of  $[^3H]$ DHT binding to cytosolic and nuclear androgen receptors *in vivo* was obtained with 5 mg of spironolactone, a dose which is close to that used by Rasmussen *et al.* (9) for testing the *in vivo* antiandrogenic effect (1 and 3 mg). Similarly, Kagawa showed that 1.6 mg of spironolactone was necessary to completely block the effect of 6 and 12  $\mu g$  of deoxycorticosterone acetate in the adrenalec-tomized rat (1).

From complete *in vitro* displacement studies (Fig. 2), spironolactone appears to be about one-twentieth as effective as DHT in displacing 50% of  $5 \times 10^{-10}$  M  $[^3H]$ DHT bound to the receptor.  $K^+$  canrenoate, on the other hand, appears to be an even less potent competitor than spironolactone for  $[^3H]$ DHT binding sites; *in vitro* competition experiments on a  $1,000 \times g$  supernatant receptor (Fig. 2) show that  $K^+$  canrenoate is about 100 times less effective than spironolactone in displacing 50% of  $[^3H]$ DHT binding.

Similarly, spirolactones inhibit aldosterone action by competing with aldosterone for the mineralocorticoid receptor. Funder *et al.* (20) found also a relatively low affinity of spironolactone for this receptor, since  $4 \times 10^{-8}$  M spironolactone inhibited the binding of  $2 \times 10^{-9}$  M aldosterone by only 54%.  $K^+$  canrenoate ap-

peared to be about 100 times less effective than spironolactone in displacing 50% of  $[^3H]$ aldosterone bound to the mineralocorticoid receptor (20). It is interesting to note that, as in the case of the androgenic receptor, the effect of opening the  $\gamma$ -lactone ring appears to decrease the affinity to the receptor.

The results of the present study should be compared with those obtained in similar experiments using cyproterone or cyproterone acetate, a well-known antiandrogen (21). This molecule has been previously reported to prevent  $[^3H]$ DHT binding to the cytosolic and nuclear androgen receptors, at about the same doses as those used in the present study (22,23). As in the case of spirolactones, it has no inhibitory effect on prostatic  $5\alpha$ -reductase (20). Spirolactones thus appear to produce their peripheral anti-androgenic effect via competition for the androgenic receptor.

The daily blood production rate of testosterone in men is about 5 mg and that of DHT is 0.3 mg (24). The doses of spirolactones commonly used in men (100–400 mg) represent, therefore, about 20–80 and 330–1330 times the production rate of testosterone and DHT, respectively. This study shows that the affinity of spironolactone for the androgenic receptor is about one twentieth of that of DHT. Therefore it seems likely that spirolactones, besides their action on testosterone synthesis, exert their antiandrogenic activity via a peripheral antagonism.

### Acknowledgments

This work has been supported in part by a grant from Institut National de la Santé et de la Recherche Médicale and by the Centre National de la Recherche Scientifique.

### References

1. Kagawa, C. M., *Endocrinology* **67**: 125, 1960.
2. Porter, G. A., *Mol Pharmacol* **4**: 224, 1968.
3. Sharp, G. W. G., C. L. Komack, and A. Leaf, *J Clin Invest* **45**: 450, 1966.
4. Herman, T. S., G. M. Fimognari, and I. S. Edelman, *J Biol Chem* **243**: 3949, 1968.

5. Fanestil, D. D., *Biochem Pharmacol* **17**: 2240, 1968.
6. Swaneck, G. E., L. L. H. Chu, and I. S. Edelman, *J Biol Chem* **245**: 5382, 1971.
7. Menard, R. H., B. Stripp, and J. R. Gillette, *Endocrinology* **94**: 1628, 1974.
8. Steelman, S. L., J. R. Brooks, E. R. Morgan, and D. J. Pananelli, *Steroids* **14**: 449, 1969.
9. Rasmusson, G. H., A. Chen, G. F. Reynolds, D. J. Patanelli, A. A. Patchett, and G. E. Arth, *J Med Chem* **15**: 1165, 1972.
10. Mahoudeau, J. A., C. W. Bardin, and M. B. Lipsett, *J Clin Invest* **50**: 1338, 1971.
11. Lowry, O. H., N. J. Rosebrough, A. L. Farr, and R. J. Randall, *J Biol Chem* **19**: 3, 1951.
12. Corvol, P., R. Falk, M. Freifeld, and C. W. Bardin, *Endocrinology* **90**: 1464, 1972.
13. Stokes, J. F., *Lancet* **2**: 911, 1962.
14. Clark, E., *JAMA* **193**: 163, 1965.
15. Froment, A., and H. Milon, *In Extra Renal Activity of Aldosterone and its Antagonists*, Excerpta Medica, Amsterdam, 1972, p. 64.
16. Brown, J. J., J. B. Ferris, R. Fraser, A. F. Lever, and J. I. S. Robertson, *In the Medical Uses of Spironolactone*, Excerpta Medica, Amsterdam, 1970, p. 27.
17. Dymling, J. F., K. O. Nilsson, and B. Hökfelt, *Acta Endocrinol (Kbh)* **70**: 104, 1972.
18. Mainwaring, W. I. P., *J Endocrinol* **45**: 531, 1969.
19. Baulieu, E. E., and I. Jung, *Biochem Biophys Res Comm* **38**: 599, 1970.
20. Funder, J. W., D. Feldman, E. Highland, and I. S. Edelman, *Biochem Pharmacol* **23**: 1493, 1974.
21. Neumann, F., and R. von Berswordt-Wallrabe, *J Endocrinol* **35**: 363, 1966.
22. Fang, S., and S. Liao, *Mol Pharmacol* **5**: 420, 1969.
23. Stern, J. M., and A. J. Eisenfeld, *Science* **166**: 223, 1969.
24. Ito, T., and R. Horton, *J Clin Invest* **50**: 1621, 1971.