

RUNS Test

1) $H_0: R_i \sim \text{Independence}$

$H_1: R_i \not\sim \text{Independence}$

2) write sequence of runs above & below mean

3) count the number of observation above the mean (n_1) and the no of observation below the mean (n_2)

max no. of possible runs (N) = $n_1 + n_2$

min no. of possible runs = one

Total no. of runs (b)

4) calculate mean & variance of b

$$\mu_b = \frac{2n_1 + n_2}{N} + \frac{1}{2}$$

$$\sigma_b^2 = \frac{2n_1 n_2 (2n_1 n_2 - N)}{N^2 (N - 2)} \quad \text{then } n_1 \text{ or } n_2 > 20$$

the distribution of b will be approximated by a N.D.

5) Standard normal statistics

$$Z_0 = \frac{b - \mu_b}{\sigma_b}$$

6) $-Z_{\alpha/2} \leq Z_0 \leq Z_{\alpha/2} = H_0 \text{ accepted.}$

$\alpha = \text{level of significance.}$

CHI SQUARE TEST

class (n)	N (range)	O_i	$E_i = \frac{N}{n}$	$O_i - E_i$	$(O_i - E_i)^2$	$\frac{(O_i - E_i)^2}{E_i}$
1	0.01 - 0.10	2	3	-1	1	1/3
2	0.11 - 0.20	3	3	0	0	0
3	0.21 - 0.30	4	3	1	1	1/3
4	0.31 - 0.40	4	3	1	1	1/3
5	0.41 - 0.50	3	3	0	0	0

$N = \text{no. of random nos (interval)}$

$n = \text{class}$

KS Test

- 1) Arrange the sequence in ascending order.
- 2) Calculate:

$$D^+ = \max_{1 \leq i \leq N} \left\{ \frac{i}{N} - R_i \right\}$$

$$D^- = \max_{1 \leq i \leq N} \left\{ R_i - \left(i - \frac{1}{N} \right) \right\}$$

level of significance, α will be mentioned.

N = total number of random nos.

i	1	2	3	4	5
R_i					

← substitute values.

- 3) calculate:

$$D = \max \{ D^+, D^- \} \quad / \quad D = \{ \max D^+, \max D^- \}$$

D^+					
D^-					

} upon calculating.

$$4) D < D_\alpha \Rightarrow H_0 \text{ accepted}$$

AUTOCORRELATION TEST

M = largest integer such that $i + (M+1)m \leq N$

$H_0: \rho_{im} = 0$, if nos are independent

$H_1: \rho_{im} \neq 0$ if nos are dependent

$$\hat{\rho}_{im} = \frac{1}{M+1} \left[\sum R_{i+km} R_{i+(k+1)m} \right] \text{ or } \left[\sum_{k=0}^M R_{i+km} \cdot R_{i+(k+1)m} \right] - 0.25$$

$$\hat{\sigma}_\rho = \sqrt{\frac{18M+7}{12(M+1)}}$$

$$Z_0 = \frac{\hat{\rho}_{im}}{\hat{\sigma}_\rho \hat{\rho}_{im}}$$

$Z_0 < z_{init}$
then accept
 H_0