Climate Risk and Human Development

Are the regions with the most extreme climate also the least developed? A pilot on France, India and the United States

DS01 - Climate Risk & Human Development Observatory

31 October 2026

Table of contents

Summary **2**

1 Introduction **2**
1.1 Context . . . 2
1.2 Research question and hypotheses . . . 3
1.3 Scope . . . 3

2 Data **3**
2.1 Sources . . . 3
2.2 Climate data . . . 4
2.3 Population . . . 4
2.4 Development . . . 4

3 Method **5**
3.1 The risk index . . . 5
3.2 Three versions of the index . . . 5
3.3 Statistical tests . . . 6

4 Results: the maps **6**
4.1 Hazard . . . 6
4.2 Exposure and development . . . 7
4.3 Risk . . . 8
4.4 High hazard and low development together . . . 10

5 Results: the hypotheses **10**
5.1 H1: more hazard, less development . . . 10
5.2 H2: is it still true inside countries? . . . 11
5.3 H3: is the ranking stable? . . . 13

6 Discussion and limitations **14**
6.1 What the results mean . . . 14
6.2 Limitations . . . 14

7 Technical implementation **15**
7.1 Architecture . . . 15
7.2 Running the pipeline . . . 15
7.3 Quality checks . . . 15
7.4 Dashboard and deployment . . . 16

8 Conclusion and next steps **16**

9 References **17**

Summary

This project asks a simple question: **are the regions with the most extreme climate also the regions with the lowest human development?** We answer it for 105 admin-1 regions (states, provinces and regions) in France, India and the United States, which together have 1.8 billion people.

We build a Python pipeline that downloads open data, computes a climate risk index for each region, tests three hypotheses, and shows the results in maps, a report and a web dashboard.

The main results are:

1. **H1 - More hazard goes with less development.** On all regions, the Spearman correlation between the climate hazard index and the Subnational Human Development Index (SHDI) is −0.59. But most of this comes from the difference between India and the two other countries.
2. **H2 - The link is weaker inside countries, but it is still there.** When we add a country effect to the regression, the hazard coefficient goes from −0.39 to −0.10. The link is clear in India and the USA, and absent in France.
3. **H3 - The ranking is stable.** Three different versions of the index give almost the same top 10. All ten highest-risk regions are in India.

These results are **associations, not causes**.

> **i** Note
>
> **Key numbers**
>
> - **105** regions in **3** countries, **1803.1 million** people; SHDI available for **99** regions.
> - H1: Spearman correlation hazard-SHDI = **-0.59** (median SHDI 0.786 vs 0.936).
> - H2: with a country effect, hazard coefficient = **-0.10** (95% CI -0.14 to -0.06).
> - H3: at least **9** of the top 10 regions are the same in the three index versions.

1 Introduction

1.1 Context

Climate risk is often discussed country by country. But a country is not uniform: in India, Rajasthan has about 170 days above 35 °C per year, while Kerala has almost none. Human development also

changes a lot inside a country. To know where help is most needed, we need to work at a smaller scale.

We work at **admin-1 level** (states, provinces, regions). This is the smallest level where climate, population and development data are all available for free.

1.2 Research question and hypotheses

> Are the regions with the most extreme climate also the regions with the lowest human development?

We test three hypotheses:

- **H1.** Regions with a higher climate hazard have a lower SHDI.
- **H2.** This is still true when we compare regions of the same country.
- **H3.** The risk ranking does not change much when we change the way the index is built.

H2 is important. India is both hotter and poorer than France and the USA, so a correlation on all regions could only say "India is different". Comparing regions inside the same country is a fairer test.

1.3 Scope

The pilot covers three countries, chosen because they are very different: France (rich, temperate, small), the USA (rich, very large, many climates) and India (lower-middle income, hot, very populated). The code can run on other countries by changing one line in `config/config.yaml`.

What this report does not claim. All results are associations. We do not prove that climate causes low development. History, geography and institutions can explain both.

2 Data

2.1 Sources

Layer	Source	Resolution	Years	Licence
Region boundaries	geoBoundaries (gbOpen)	admin-1 polygons	2024 release	CC-BY 4.0
Daily climate	NASA POWER (MERRA-2)	points	1991-2023	Open, no key
Population	GHS-POP R2023A	~1 km grid	2020	CC-BY 4.0
Development	Global Data Lab SHDI	admin-1	latest year	Free for research
National context	World Bank WDI	country	2010-2023	CC-BY 4.0

2.2 Climate data

NASA POWER gives one daily time series per point. Inside each region we choose between 1 and 8 sample points (one point per 40,000 km²), spread over the whole region. This gives **308 points** in total. For each point we download every day from 1991 to 2023.

From the daily data we compute six indicators per year, then take the mean over 2004-2023:

Code	Indicator	Group
HD35	days with maximum temperature of 35 °C or more	heat
TR25	nights with minimum temperature of 25 °C or more	heat
CDD	longest run of days with less than 1 mm of rain	dry
WBAL	rain minus evapotranspiration (Hargreaves formula)	dry
Rx1day	heaviest rain in one day	wet
R50	days with more than 50 mm of rain	wet

Table 2. Mean climate indicators by country, 2004-2023.

Indicator	FRA	IND	USA
hd35	2.170	60.420	11.250
tr25	0.470	96.230	31.600
cdd	23.320	59.670	24.830
wbal	-126.720	-5.530	10.780
rx1d	28.890	66.930	50.470
r50	0.050	2.370	0.870

A quick check: Rajasthan (desert) has 169 heat days per year and Kerala (monsoon coast) has 0. This is what we expect, so the data look correct.

2.3 Population

GHS-POP gives the number of people in each 1 km cell. We sum the cells inside each region. We also give each person to the nearest climate point. The two totals agree (1,803.1 million against 1,805.5 million, a 0.1% difference).

2.4 Development

The SHDI comes from the Global Data Lab. Its region names are not always the same as in geoBoundaries (for example *Odisha / Orissa*). We matched names in three steps: exact match, a short manual list, then fuzzy matching. **99 of the 105 regions** have an SHDI. The 6 others (Ladakh and five small US territories) are kept on the climate maps but are not used in the tests. We do not fill them with the country average, because this would create fake data.

3 Method

3.1 The risk index

We follow the IPCC idea: risk needs a **hazard**, people **exposed** to it, and a **vulnerability**. Each part is a score between 0 and 1.

To build each score we use **percentile ranks**: the region with the lowest value gets a score close to 0 and the region with the highest value gets 1. Ranks let us mix days, millimetres and index points, and one extreme region does not squash all the others.

Hazard is the mean of three sub-scores:

$$\text{Hazard} = \tfrac{1}{3}\left(H_{\text{heat}} + H_{\text{dry}} + H_{\text{wet}}\right)$$

H_{heat} is the mean rank of HD35 and TR25, H_{dry} the mean rank of CDD and (reversed) WBAL, and H_{wet} the mean rank of Rx1day and R50.

Exposure is the rank of the region's population. We do not multiply population by heat days here, because heat is already in the hazard score: it would be counted twice.

Vulnerability is $1 - \text{rank(SHDI)}$. Low development means less money, health and education to face a climate shock.

Risk is the product of the three:

$$\text{Risk} = \text{Hazard} \times \text{Exposure} \times \text{Vulnerability}$$

We use a product because a region needs all three to be at risk: a hot desert with nobody living there is not a disaster.

3.2 Three versions of the index

Version	Formula	Why
`equal` (main)	Hazard × Exposure × Vulnerability	the main result
`additive`	mean of the three scores	does the product matter?
`per_person`	Hazard × Vulnerability	removes population: is the ranking only a size effect?

The `per_person` version is useful because big regions always get a high exposure score. If the top regions stay the same without population, the ranking is not only about size.

3.3 Statistical tests

- **H1:** Spearman rank correlation between hazard and SHDI, on all regions and then country by country. We use Spearman because it works on ranks and does not need a normal distribution. We also compare the median SHDI of the more hazardous half and the less hazardous half.
- **H2:** two linear regressions (OLS) with SHDI as the outcome:
  - M1: SHDI $= a + b \cdot$ Hazard
  - M2: the same, plus one dummy variable per country ("country effect").

  The country dummies remove the average difference between countries, so b in M2 only uses differences between regions of the same country. We also fit M1 in each country separately. We use robust (HC3) standard errors.
- **H3:** Spearman correlation between the rankings of the three versions, and the number of regions that are in the top 10 of both versions.

We use a 5% significance level.

4 Results: the maps

4.1 Hazard




Figure 1: Climate hazard index

India has the highest hazard (mean 0.67), then the USA (0.45) and France (0.32). In the USA, the south and south-west are much more hazardous than the north-east.

4.2 Exposure and development




Figure 2: Exposure (population rank)




Figure 3: Subnational Human Development Index

SHDI goes from 0.614 (Bihar) to 0.961 (top US states). India goes from 0.61 to 0.81, the USA from 0.88 to 0.96 and France from 0.87 to 0.96. The differences inside India are larger than the difference between the USA and France.

Table 1. Summary by country.

Country	Regions	Population (m)	Hazard (mean)	Exposure (mean)	Vulnerability (mean)	Risk (median)	Person-days of heat (bn/yr)
ALL	105	1803.072	0.505	0.505	0.495	0.089	152.292
FRA	13	64.250	0.317	0.472	0.481	0.059	0.152
IND	36	1399.785	0.666	0.629	0.818	0.430	146.240
USA	56	339.036	0.445	0.432	0.277	0.041	5.900

4.3 Risk




Figure 4: Composite climate risk

Table 3. The ten highest-risk regions.

Region	Country	Population (m)	Hazard	Exposure	Vulnerability	Risk
Bihār	IND	125.300	0.849	0.981	0.990	0.825
Uttar Pradesh	IND	232.200	0.844	1.000	0.965	0.815
Madhya Pradesh	IND	84.600	0.860	0.962	0.950	0.786
Gujarāt	IND	70.900	0.913	0.933	0.889	0.757
Jhārkhand	IND	39.200	0.830	0.876	0.980	0.713
Odisha	IND	46.700	0.814	0.905	0.965	0.711
Andhra Pradesh	IND	53.400	0.833	0.914	0.899	0.685
West Bengal	IND	100.400	0.763	0.971	0.909	0.674
Mahārāshtra	IND	126.400	0.871	0.991	0.773	0.667
Chhattīsgarh	IND	29.700	0.833	0.829	0.919	0.635

All ten highest-risk regions are in India. Bihar, Uttar Pradesh and Madhya Pradesh are hot, very populated and have a low SHDI. On this shared scale France and the USA look pale. Inside their own country, the highest-risk regions are Texas, Alabama and Louisiana in the USA, and Auvergne-Rhône-Alpes, Occitanie and Nouvelle-Aquitaine in France (see the figure below and the `rank_in_country` column).




Figure 5: The five highest-risk regions in each country




Figure 6: Risk without population (hazard × vulnerability)

The map without population looks very similar. Small Indian states such as Tripura move up, but the big northern and central states stay at the top.

4.4 High hazard and low development together




Figure 7: Regions with high hazard and low development

We split hazard and vulnerability at their median. 33 regions are both above the median hazard and above the median vulnerability: 28 in India and 5 in the USA, none in France. These are the regions where adaptation is most urgent.

5 Results: the hypotheses

5.1 H1: more hazard, less development

Table 4. H1: Spearman correlation between hazard and SHDI.

Group	Regions	Spearman rho	p-value	Median SHDI, more hazard	Median SHDI, less hazard
ALL	99	-0.590	< 0.001	0.786	0.936
FRA	13	0.180	0.566	0.908	0.898
IND	35	-0.480	0.003	0.686	0.737
USA	51	-0.480	< 0.001	0.923	0.942

On all regions, the correlation is -0.59 and very significant. The median SHDI is 0.786 in the more hazardous half and 0.936 in the other half.




Figure 8: Hazard and development, one line per country

The figure shows why we must be careful. The points form **three groups**: India at the bottom right, the USA and France at the top. A large part of the correlation is simply the gap between India and the other two countries. Inside each country, the correlation is smaller: −0.48 in India and in the USA (both significant), and +0.18 in France (not significant).

H1 is supported, but mainly because of the difference between countries.

5.2 H2: is it still true inside countries?

Table 5. H2: hazard coefficient in the two regression models.

Model	Coefficient (95% CI)	p-value	N	R squared
M1 all regions	-0.392 (-0.463 to -0.321)	< 0.001	99	0.480
M2 country effects	-0.101 (-0.145 to -0.056)	< 0.001	99	0.910




Figure 9: Hazard coefficient with and without country effects

How to read the coefficient: hazard goes from 0 to 1, so the coefficient is the change in SHDI from the least to the most hazardous region.

- Without country effects (M1), the coefficient is −0.39.
- With country effects (M2), it falls to **−0.10** (95% CI −0.14 to −0.06). In simple words: in the same country, +0.1 of hazard goes with about −0.01 of SHDI.

So about three quarters of the link is between countries, and one quarter remains inside countries.

Table 6. H2: slope of SHDI on hazard in each country.

Country	Regions	Slope (95% CI)	p-value	R squared
FRA	13	0.007 (-0.140 to 0.154)	0.930	0.000
IND	35	-0.127 (-0.204 to -0.049)	0.001	0.180
USA	51	-0.081 (-0.125 to -0.037)	< 0.001	0.230




Figure 10: Slope inside each country

The slope is negative and significant in India and the USA, and close to zero in France. France has only 13 regions and small climate differences, so we cannot conclude anything there.

H2 is supported for India and the USA, with a smaller effect than in H1.

5.3 H3: is the ranking stable?

Table 7. H3: agreement between the three index versions.

Version A	Version B	Regions	Spearman rho	Same in top 10
additive	equal	99	0.970	10
additive	per_person	99	0.890	9
equal	per_person	99	0.860	9




Figure 11: Rank correlation between index versions

The three versions are strongly correlated (Spearman correlation of at least 0.86), and at least 9 of the top 10 regions are the same in every pair. Even the version without population keeps the same leaders, so the ranking is not only a size effect.

H3 is supported.

6 Discussion and limitations

6.1 What the results mean

Climate hazard and low human development overlap, mostly between countries but also inside India and the USA. The effect inside countries is moderate: climate is one factor among many. For policy, this means that help should target **regions**, not whole countries, and that the regions with both high hazard and low development (map above) should come first.

6.2 Limitations

1. **Association, not causation.** History, geography and the economy can explain both climate and development.

2. **Neighbouring regions are similar.** The regressions treat each region as independent, but neighbours share climate and economy. So the p-values are probably too small. A spatial regression model would be a good next step.
3. **Only 3 countries and 99 regions.** France has only 13 regions, so its results are uncertain.
4. **One development indicator.** SHDI does not show inequality inside a region.
5. **Fixed thresholds.** 35 °C and 50 mm are choices. France has almost no days above 35 °C, so a lower threshold would change its map.
6. **Static population.** Population is for 2020, but the climate mean is for 2004-2023.
7. **Coarse sampling.** At most 8 climate points per region: enough for Bihar, thin for Alaska.

7 Technical implementation

7.1 Architecture

The project is a Python package (`cro`) with one command-line tool. Each step reads and writes Parquet files, so a step can be run again without downloading everything.

```
config/config.yaml        all settings (countries, thresholds, years)
src/cro/ingest/           download boundaries, climate, population, development
src/cro/features/build.py hazard, exposure, vulnerability, risk
src/cro/analysis/         descriptive statistics, H1/H3 tests, H2 regression
src/cro/viz/              static maps, charts, interactive maps
src/cro/db/               PostGIS schema and loader (optional)
src/cro/app/              Dash dashboard + summary and report pages (EN/FR)
src/cro/report/           report tables + Quarto rendering
tests/                    unit tests and data checks (pytest)
```

7.2 Running the pipeline

```
cro all          # download, build the index, run the analysis, export the figures
cro validate     # 8 data checks (exit code 1 if one fails)
cro report       # build the report tables and render this report
cro load         # optional: copy the tables into PostGIS (docker compose up -d)
cro app          # dashboard at http://127.0.0.1:8050
```

7.3 Quality checks

- `cro validate` checks the data: unique regions, valid geometry, population totals, physical ranges of the climate values, scores between 0 and 1, SHDI coverage and output tables.
- `pytest` runs unit tests on small hand-made examples (percentile ranks, hazard, exposure, vulnerability, Spearman test, rank comparison) and smoke tests for the dashboard.
- The report and the dashboard read their numbers from the same CSV tables, so they cannot show different values.

Data checks

- Checks run: **8**
- Passed: **8**
- Failed: **0**
- `PASS` **boundaries** - 105 regions, 3 countries, CRS EPSG:4326
- `PASS` **climate sample points** - 308 points covering 105/105 regions
- `PASS` **population reconciles** - region total 1,803.1M vs point total 1,805.5M (0.1% difference)
- `PASS` **climate ranges** - 105 regions checked
- `PASS` **index bounds** - all index columns within 0..1
- `PASS` **master table** - 105 rows, 0 duplicates, 0 regions missing
- `PASS` **development coverage** - SHDI available for 99/105 regions (94%)
- `PASS` **output tables** - 7 tables present

7.4 Dashboard and deployment

The dashboard is built with Plotly Dash and Flask. It shows the maps, the ranking, the hazard-SHDI chart and a profile for each region, in English and French. It reads PostGIS when the database is running and Parquet files otherwise. It is deployed for free on Render.

Interactive maps (no Python needed):

- **Risk**: https://climate-risk-observatory.onrender.com/maps/map_web_risk.html
- **Hazard**: https://climate-risk-observatory.onrender.com/maps/map_web_hazard.html
- **Exposure (population)**: https://climate-risk-observatory.onrender.com/maps/map_web_exposure.html
- **SHDI**: https://climate-risk-observatory.onrender.com/maps/map_web_shdi.html

8 Conclusion and next steps

The project shows that, in France, India and the USA, regions with a more extreme climate tend to have lower human development. Most of this link is between countries, but a smaller link remains inside India and the USA. The ranking of the most at-risk regions is stable when we change the index formula.

Possible next steps:

- add more countries (the code is ready, it is a change in the config file);
- use a spatial regression model to take neighbours into account;
- add a poverty indicator to see inequality inside regions;
- test other heat thresholds (for example 30 °C).

9 References

Global Data Lab. *Subnational Human Development Index (SHDI)*. Radboud University. https://globaldatalab.org/shdi/

Hargreaves, G. H., & Samani, Z. A. (1985). Reference crop evapotranspiration from temperature. *Applied Engineering in Agriculture*, 1(2), 96-99.

IPCC (2022). *Climate Change 2022: Impacts, Adaptation and Vulnerability*. Contribution of Working Group II to the Sixth Assessment Report.

NASA Langley Research Center. *POWER Project Data Access Viewer*. https://power.larc.nasa.gov/

Pesaresi, M., et al. (2023). *GHS-POP R2023A - GHS population grid multitemporal (1975-2030)*. European Commission, JRC.

Runfola, D., et al. (2020). geoBoundaries: A global database of political administrative boundaries. *PLOS ONE*, 15(4), e0231866.

World Bank. *World Development Indicators*. https://databank.worldbank.org/source/world-development-indicators