

Complete Genome Sequence of *Rhodococcus* sp. Strain WMMA185, a Marine Sponge-Associated Bacterium

Navid Adnani,^a Doug R. Braun,^a Bradon R. McDonald,^b Marc G. Chevrete,^{b,c} Cameron R. Currie,^b Tim S. Bugni^a

Pharmaceutical Sciences Division, University of Wisconsin–Madison, Madison, Wisconsin, USA^a; Department of Bacteriology, University of Wisconsin–Madison, Madison, Wisconsin, USA^b; Laboratory of Genetics, University of Wisconsin–Madison, Madison, Wisconsin, USA^c

The *Rhodococcus* strain WMMA185 was isolated from the marine sponge *Chondrilla nucula* as part of ongoing drug discovery efforts. Analysis of the 4.44-Mb genome provides information regarding interspecies interactions as pertains to regulation of secondary metabolism and natural product biosynthetic potentials.

Received 24 October 2016 Accepted 25 October 2016 Published 15 December 2016

Citation Adnani N, Braun DR, McDonald BR, Chevrete MG, Currie CR, Bugni TS. 2016. Complete genome sequence of *Rhodococcus* sp. strain WMMA185, a marine sponge-associated bacterium. *Genome Announc* 4(6):e01406-16. doi:10.1128/genomeA.01406-16.

Copyright © 2016 Adnani et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Tim S. Bugni, tim.bugni@wisc.edu.

Drug resistance is now commonplace for microbial human pathogens and approaches to discovering new antibiotics, indeed, completely new antibacterial chemotypes, continue to be developed to counter this problem (1). Accordingly, the number of natural products (as possible drug leads) available for screening, their producers and their production conditions must necessarily increase. In support of this tenet, the production of natural products by one producing organism in the presence of another organism has garnered tremendous interest (2–6). Such co-culturing of microorganisms has proven extremely effective for coaxing microbes into making natural products that would not otherwise be produced. This approach to new natural product generation underscores the importance of having genomic information available for co-cultured organisms (6). Importantly, the structural diversity of natural products enabled by co-culturing stems, in large part, from the diversity of co-cultured organisms.

The genus *Rhodococcus* is a Gram-positive bacterium within the subgroup actinobacteria whose members are commonly associated with bioremediation and biocatalytic processes (7–9); steroids, nitriles, lignins, and organosulfur agents are but a few compound classes degraded by rhodococci (10). Additionally, although rare, select *Rhodococcus* spp. are human pathogens; pathogenicity has been associated with similarities to *Mycobacterium* spp. (11, 12).

Genome analyses of the few reported *Rhodococcus* spp. highlight tremendous biosynthetic potential despite a scarcity of isolated secondary metabolites (13). Recently, cocultures of *Rhodococcus* spp. with other actinobacteria, including *Streptomyces* spp. (14, 15) and *Micromonosporaceae* (2) have been shown to produce otherwise undetectable secondary metabolites. In particular, marine invertebrate-associated *Rhodococcus* sp. WMMA185 induced biosynthesis in other marine actinobacteria via interspecies interactions; the precise nature of these interactions awaits further investigation. Using genomic data from WMMA185, mechanisms of biosynthetic regulation and interspecies communication may be deciphered in an effort to access unexploited (or cryptic) bio-

synthetic potentials from actinobacteria or, for that matter, WMMA185 itself.

Rhodococcus sp. strain WMMA185 was isolated in 2011 from a marine sponge *Chondrilla nucula* collected off the coast of the Florida Keys. WMMA185 was isolated from a plate prepared using R2A medium supplemented with 50% artificial seawater (ASW).

The complete genome of *Rhodococcus* sp. WMMA185 was sequenced at the Duke Center for Genomic and Computational Biology (GCB) using PacBio RS II (Pacific Biosciences) technology. Reads were assembled using the HGAP assembler (16) into a single contig. Open reading frames were predicted by Prodigal (17) and annotated using HMMer models for the TIGRfam (18), KEGG (19, 20), and PFAM (21, 22) databases. The genome was found to be 4.44 Mb in length and has 64.08% G+C, and 90.39% coding density. The organism's secondary metabolic content/potential was assessed using anti-SMASH (23, 24), PRISM (25), and custom pipelines. Among other cluster types, a total of two type I polyketide (PKS), eight nonribosomal peptide (NRPS), and two terpene biosynthetic gene clusters were identified within the WMMA185 genome.

Accession number(s). The complete genome sequence of *Rhodococcus* sp. strain WMMA185 has been deposited at the DDBJ/EMBL/GenBank under the sequence GenBank accession no. CP017014.

ACKNOWLEDGMENTS

This work was supported by National Institutes of Health grants R01-GM104192 and U19 A1109673.

FUNDING INFORMATION

This work, including the efforts of Timothy S. Bugni, was funded by HHS | National Institutes of Health (NIH) (R01-GM104192). This work, including the efforts of Cameron R. Currie, was funded by HHS | National Institutes of Health (NIH) (U19 A1109673).

REFERENCES

- Centers for Disease Control and Prevention. 2013. Antibiotic resistance threats in the United States, 2013. U.S. Department of Health and Human Services, CDC.

2. Adnani N, Vazquez-Rivera E, Adibhatla SN, Ellis GA, Braun DR, Bugni TS. 2015. Investigation of interspecies interactions within marine *Micromonosporaceae* using an improved co-culture approach. *Mar Drugs* 13:6082–6098. <http://dx.doi.org/10.3390/md13106082>.
3. Bertrand S, Bohni N, Schnee S, Schumpp O, Gindro K, Wolfender JL. 2014. Metabolite induction via microorganism co-culture: A potential way to enhance chemical diversity for drug discovery. *Biotechnol Adv* 32:1180–1204. <http://dx.doi.org/10.1016/j.biotechadv.2014.03.001>.
4. Netzker T, Fischer J, Weber J, Mattern DJ, König CC, Valiante V, Schroeckh V, Brakhage AA. 2015. Microbial communication leading to the activation of silent fungal secondary metabolite gene clusters. *Front Microbiol* 6:299. <http://dx.doi.org/10.3389/fmicb.2015.00299>.
5. Demain AL. 2014. Importance of microbial natural products and the need to revitalize their discovery. *J Ind Microbiol Biotechnol* 41:185–201. <http://dx.doi.org/10.1007/s10295-013-1325-z>.
6. Pettit RK. 2009. Mixed fermentation for natural product drug discovery. *Appl Microbiol Biotechnol* 83:19–25. <http://dx.doi.org/10.1007/s00253-009-1916-9>.
7. Larkin MJ, Kulakov LA, Allen CCR. 2005. Biodegradation and *Rhodococcus*—masters of catabolic versatility. *Curr Opin Biotechnol* 16: 282–290. <http://dx.doi.org/10.1016/j.copbio.2005.04.007>.
8. Larkin MJ, Kulakov LA, Allen CCR. 2006. Biodegradation by members of the genus *Rhodococcus*: biochemistry, physiology, and genetic adaptation. *Adv Appl Microbiol* 59:1–29. [http://dx.doi.org/10.1016/S0065-2164\(06\)59001-X](http://dx.doi.org/10.1016/S0065-2164(06)59001-X).
9. De Carvalho CCCR, Costa SS, Fernandes P, Couto I, Viveiros M. 2014. Membrane transport systems and the biodegradation potential and pathogenicity of genus *Rhodococcus*. *Front Physiol* 5:133. <http://dx.doi.org/10.3389/fphys.2014.00133>.
10. Yam KC, Okamoto S, Roberts JN, Eltis LD. 2011. Adventures in *Rhodococcus*—from steroids to explosives. *Can J Microbiol* 57:155–168. <http://dx.doi.org/10.1139/W10-115>.
11. Wang X, Coulson GB, Miranda-CasoLuengo AA, Miranda-CasoLuengo R, Hondalus MK, Meijer WG. 2014. IcgA is a virulence factor of *Rhodococcus equi* That modulates intracellular growth. *Infect Immun* 82: 1793–1800. <http://dx.doi.org/10.1128/IAI.01670-13>.
12. Letek M, González P, MacArthur I, Rodríguez H, Freeman TC, Valero-Rello A, Blanco M, Buckley T, Cherevach I, Fahey R, Hapeshi A, Holdstock J, Leadon D, Navas J, Ocampo A, Quail MA, Sanders M, Scortti MM, Prescott JF, Fogarty U, Meijer WG, Parkhill J, Bentley SD, Vázquez-Boland JA. 2010. The genome of a pathogenic *Rhodococcus*: cooptive virulence underpinned by key gene acquisitions. *PLoS Genet* 6:e1001145. <http://dx.doi.org/10.1371/journal.pgen.1001145>.
13. Kulakov LA, Larkin MJ. 2002. Genomic organization of rhodococcus in genomics of GC-rich Gram-positive bacteria, pp 15–46. A. Danchin (ed), Academic Press, Norfolk, United Kingdom.
14. Kurosawa K, Ghiviriga I, Sambandan TG, Lessard PA, Barbara JE, Rha C, Sinskey AJ. 2008. Rhodostreptomycins, antibiotics biosynthesized following horizontal gene transfer from *Streptomyces padanus* to *Rhodococcus fascians*. *J Am Chem Soc* 130:1126–1127. <http://dx.doi.org/10.1021/ja077821p>.
15. Onaka H, Mori Y, Igarashi Y, Furumai T. 2011. Mycolic acid-containing bacteria induce natural-product biosynthesis in streptomyces species. *Appl Environ Microbiol* 77:400–406. <http://dx.doi.org/10.1128/AEM.01337-10>.
16. Ge F, Wang LS, Kim J. 2005. The cobweb of life revealed by genome-scale estimates of horizontal gene transfer. *PLoS Biol* 3:e316. <http://dx.doi.org/10.1371/journal.pbio.0030316>.
17. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <http://dx.doi.org/10.1186/1471-2105-11-119>.
18. Haft DH, Selengut JD, Richter RA, Harkins D, Basu MK, Beck E. 2013. TIGRFAMs and genome properties in 2013. *Nucleic Acids Res* 41: D387–D395. <http://dx.doi.org/10.1093/nar/gks1234>.
19. Kanehisa M, Goto S, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2014. Data, information, knowledge and principle: back to metabolism in KEGG. *Nucleic Acids Res* 42:D199–D205. <http://dx.doi.org/10.1093/nar/gkt1076>.
20. Kanehisa M, Sato Y, Kawashima M, Furumichi M, Tanabe M. 2016. KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res* 44:D457–D462. <http://dx.doi.org/10.1093/nar/gkv1070>.
21. Finn RD, Bateman A, Clements J, Coghill P, Eberhardt RY, Eddy SR, Heger A, Hetherington K, Holm L, Mistry J, Sonnhammer ELL, Tate J, Punta M. 2014. Pfam: the protein families database. *Nucleic Acids Res* 42:D222–D230. <http://dx.doi.org/10.1093/nar/gkt1223>.
22. Finn RD, Coghill P, Eberhardt RY, Eddy SR, Mistry J, Mitchell AL, Potter SC, Punta M, Qureshi M, Sangrador-Vegas A, Salazar GA, Tate J, Bateman A. 2016. The Pfam protein families database: towards a more sustainable future. *Nucleic Acids Res* 44:D279–D285. <http://dx.doi.org/10.1093/nar/gkv1344>.
23. Medema MH, Blin K, Cimermancic P, de Jager V, Zakrzewski P, Fischbach MA, Weber T, Takano E, Breitling R. 2011. antiSMASH: rapid identification, annotation and analysis of secondary metabolite biosynthesis gene clusters in bacterial and fungal genome sequences. *Nucleic Acids Res* 39:W339–W346. <http://dx.doi.org/10.1093/nar/gkr466>.
24. Weber T, Blin K, Duddela S, Krug D, Kim HU, Brucoleri R, Lee SY, Fischbach MA, Müller R, Wohlleben W, Breitling R, Takano E, Medema MH. 2015. antiSMASH 3.0—a comprehensive resource for the genome mining of biosynthetic gene clusters. *Nucleic Acids Res* 43: W237–W243. <http://dx.doi.org/10.1093/nar/gkv437>.
25. Skinnider MA, Dejong CA, Rees PN, Johnston CW, Li H, Webster ALH, Wyatt MA, Magarvey NA. 2015. Genomes to natural products PRediction informatics for secondary metabolomes (PRISM). *Nucleic Acids Res* 43:9645–9662. <http://dx.doi.org/10.1093/nar/gkv1012>.