

Dissipative dynamics of coupled quantum dots under quantum measurement

Shi-Hua Ouyang,¹ Chi-Hang Lam,² and J. Q. You¹

¹*Department of Physics and Surface Physics Laboratory (National Key Laboratory), Fudan University, Shanghai 200433, China*

²*Department of Applied Physics, Hong Kong Polytechnic University, Hung Hom, Hong Kong, China*

(Dated: November 16, 2005)

We have studied the dissipative dynamics of a solid-state qubit with an extra electron confined to either one of two coupled quantum dots. Previous theoretical work based on Bloch-type rate equations gave an unphysical uniform occupation probability of the electron in the quantum dots even for non-identical dots. We show that this is due to neglecting higher order interactions in the analysis. By including higher order terms, we obtain expected asymmetric occupation probabilities for non-identical dots. Our work demonstrates that the high order interaction terms can lead to important qualitative impacts on the operation of the qubit.

I. INTRODUCTION

Quantum measurement has attracted much attention because of its fundamental importance as well as its relevance to quantum computing. However, there are still vastly different points of view on quantum measurement. On one hand, the Zeno effect predicts that continuous measurement can freeze the state of a quantum system.¹ On the other hand, interaction with the outside environment such as a detector is expected to lead to the collapse of the state of the measured system. Therefore, reconciling these views by detailing the real mechanisms for selected model systems is very important.

Many experiments on two-level quantum systems (see, e.g., Refs. 2,3,4) have been performed to study this problem. Together with experimental investigations, some theoretical studies were also devoted to this important topic. Gurvitz *et al.*^{5,6} have theoretically studied a mesoscopic qubit-plus-detector system. The qubit consists of two coupled-quantum dots (CQD's) while the detector is a quantum point contact (QPC) interacting locally with one of the dots. The authors derived a set of Bloch-type rate equations to describe the measurement effects. Goan *et al.*⁷ considered the same setup and developed a Lindblad quantum master equation using the quantum trajectory approach for dissipative processes. However, while reasonable result was obtained for a pair of identical quantum dots in both Refs. 5 and 7, these works predicted unreasonable equal occupation probabilities in the dots when their energy levels are different. Subsequently, Li *et al.*⁸ developed a unified quantum master equation with the Markovian approximation and obtained distinct occupation probabilities for non-identical quantum dots as expected.

In this work, we further study this qubit-plus-detector system following the original approach of Gurvitz *et al.*⁵ We have found that the unphysical result of identical occupation probabilities in non-identical dots mentioned above is only due to neglecting higher-order terms in their derivation. We will show that higher-order terms are of fundamental importance. Retaining additional terms in our calculations leads to physically valid results consistent with those of Li *et al.*⁸ Our analysis also illustrates

the detailed mechanism of quantum measurement.

II. BLOCH-TYPE RATE EQUATIONS

The system we study is shown schematically in Fig. 1. It can be modeled by the Hamiltonian $H = H_{\text{CQD}} + H_{\text{QPC}} + H_{\text{int}}$, with

$$\begin{aligned} H_{\text{CQD}} &= E_1 a_1^\dagger a_1 + E_2 a_2^\dagger a_2 + \Omega_0 (a_1^\dagger a_2 + a_2^\dagger a_1), \\ H_{\text{QPC}} &= \sum_l E_l a_l^\dagger a_l + \sum_r E_r a_r^\dagger a_r + \sum_{l,r} \Omega_{lr} (a_l^\dagger a_r + \text{H.c.}), \\ H_{\text{int}} &= \sum_{l,r} \delta\Omega_{lr} a_1^\dagger a_1 (a_l^\dagger a_r + a_r^\dagger a_l). \end{aligned} \quad (1)$$

Here, H_{CQD} is the Hamiltonian of the CQD's. The Hubbard terms are neglected because we consider only one extra electron in the CQD's. The parameter Ω_0 denotes the hopping amplitude of the extra electron between the two single-dot states (see Fig.1). H_{QPC} is the Hamiltonian of the QPC, and H_{int} is the interaction Hamiltonian between the CQD's and the QPC. Electrostatic effects are included in H_{int} : an extra electron in the left dot will lead to an effective variation $\delta\Omega_{lr}$ in the coupling Ω_{lr} between the states E_l and E_r in the two reservoirs of the QPC. The resulting coupling then becomes $\Omega'_{lr} = \Omega_{lr} + \delta\Omega_{lr}$.

In the occupation representation, the many-body wave function of the entire system (the CQD's plus the QPC) can be written as

$$\begin{aligned} |\psi(t)\rangle &= [b_1(t)a_1^\dagger + b_2(t)a_2^\dagger \\ &+ \sum_{l,r} b_{1lr}(t)a_1^\dagger a_r^\dagger a_l + \sum_{l,r} b_{2lr}(t)a_2^\dagger a_r^\dagger a_l \\ &+ \sum_{l<l',r<r'} b_{1ll'rr'}(t)a_1^\dagger a_r^\dagger a_{r'}^\dagger a_{l'} a_l a_{l'} \\ &+ \sum_{l<l',r<r'} b_{2ll'rr'}(t)a_2^\dagger a_r^\dagger a_{r'}^\dagger a_{l'} a_l a_{l'} + \cdots] |0\rangle, \end{aligned} \quad (2)$$

where $b_j(t)$, $j = 1, 2, 1lr, 2lr, \dots$, are the time-dependent probability amplitudes of finding the system in the corresponding states. The vacuum state $|0\rangle$ corresponds to the state of the entire system with no extra electron in the




FIG. 1: Two coupled quantum dots (CQD's) connected with a quantum point contact (QPC). Initially, all states up to the Fermi energy level (μ_L or μ_R) in each reservoir are filled by electrons. E_1 and E_2 denote the energy levels of an extra electron in the left and right quantum dots, respectively. The location of this extra electron will affect the current through the QPC. The quantum state of the CQD's can hence be read out by a current measurement.

CQD's and before the transfer of any electron between the reservoirs in the QPC. Then, $b_{1lr}(t)$ for instance denotes the probability amplitude that the extra electron is in the left dot and an electron has passed through the barrier in the QPC. The initial conditions are $b_1(0) = 1$ and $b_j(0) = 0$ for $j = 2, 1lr, 2lr, \dots$. Substituting the many-body wave function [Eq. (2)] and the Hamiltonian [Eq. (1)] into the time-dependent Schrödinger equation $i|\dot{\psi}(t)\rangle = H|\psi(t)\rangle$ gives an infinite set of linear equations for the probability amplitudes $b_j(t)$. Performing a Laplace transform to each equation, the transformed amplitude $b_j(E) = \int_0^{+\infty} e^{iEt} b_j(t) dt$ follows:

$$(E - E_1)b_1(E) - \Omega_0 b_2(E) - \sum_{l,r} \Omega'_{lr} b_{1lr}(E) = i, \quad (3a)$$

$$(E - E_2)b_2(E) - \Omega_0 b_1(E) - \sum_{l,r} \Omega_{lr} b_{2lr}(E) = 0, \quad (3b)$$

$$(E + E_l - E_1 - E_r)b_{1lr}(E) - \Omega'_{lr} b_1(E) - \Omega_0 b_{2lr}(E) - \sum_{l',r'} \Omega'_{l'r'} b_{1l'l'r'r'}(E) = 0, \quad (3c)$$

$$(E + E_l - E_2 - E_r)b_{2lr}(E) - \Omega_{lr} b_2(E) - \Omega_0 b_{1lr}(E) - \sum_{l',r'} \Omega_{l'r'} b_{2l'l'r'r'}(E) = 0, \quad (3d)$$

.....

To solve this set of equations, Gurvitz *et al.*^{5,6} introduced an approximation by keeping terms up to first order in Eqs. (3a) and (3b), second order in Eq. (3c) and (3d), and so on. When the degrees of freedom of the QPC are traced out, a coupled set of Bloch-type rate equations are derived. As shown in Fig. 3 of Ref. 5(b), the resulting occupation probabilities for states $|E_1\rangle$ and $|E_2\rangle$ are 0.5 when $t \rightarrow \infty$ for both symmetric and asymmetric CQD's. This is correct for the symmetric case with

identical quantum dots, but is unphysical in the latter case.

Here we show that this unphysical feature can be rectified if we keep terms up to second order in Eqs. (3a) and (3b), third order in Eqs. (3c) and (3d), and so on. Also, following Ref. 5, we consider the high voltage regime with $eV_d \gg \Omega^2 \rho$, where $V_d = \mu_L - \mu_R$, and ρ is the density of states in the reservoirs. The resulting equations for $b_j(E)$ are (see the Appendix)

$$[E - E_1 + \frac{iD'}{2V_d}(V_d + E - E_1)]b_1(E) - \Omega_0 b_2(E) - \frac{i\Lambda}{2V_d} b_2(E) = i, \quad (4a)$$

$$[E - E_2 + \frac{iD}{2V_d}(V_d + E - E_2)]b_2(E) - \Omega_0 b_1(E) - \frac{i\Lambda}{2V_d} b_1(E) = 0, \quad (4b)$$

$$[E + E_{lr1} + \frac{iD'}{2V_d}(V_d + E + E_{lr1})]b_{1lr}(E) - \Omega' b_1(E) - \left(\Omega_0 + \frac{i\Lambda}{2V_d}\right) b_{2lr}(E) = 0, \quad (4c)$$

$$[E + E_{lr2} + \frac{iD}{2V_d}(V_d + E + E_{lr2})]b_{2lr}(E) - \Omega b_2(E) - \left(\Omega_0 + \frac{i\Lambda}{2V_d}\right) b_{1lr}(E) = 0, \quad (4d)$$

.....

where $E_{lr1} = E_l - E_r - E_1$, $E_{lr2} = E_l - E_r - E_2$, $D' = 2\pi\rho_l\rho_r\Omega'^2 V_d$, $D = 2\pi\rho_l\rho_r\Omega^2 V_d$, and $\Lambda = 2\pi\rho_l\rho_r\Omega_0\Omega' V_d$. Now, we introduce the notation

$$\sigma_{ij}^{(n)} = \sum_{l \dots r \dots} b_{il \dots r \dots}(t) b_{jl \dots r \dots}^*(t).$$

For instance, $\sigma_{11}^{(1)} = \sum_{lr} b_{1lr}(t) b_{1lr}^*(t)$. Thus, the equations for the amplitudes $b_j(E)$, $j = 1, 2, 1lr, 2lr, \dots$, can be converted to a new set of equations for $\sigma_{ij}^{(n)}$ using the inverse Laplace transform:

$$\sigma_{ij}^{(n)} = \sum_{l \dots r \dots} \int \frac{dE dE'}{4\pi^2} b_{il \dots r \dots}(E) b_{jl \dots r \dots}^*(E') e^{i(E' - E)t}.$$

We now multiply Eq. (4c) by $b_{1lr}^*(E')$. The resulting equation is then subtracted by its complex conjugate after exchanging E with E' . We obtain

$$\begin{aligned} & \int \int \frac{dE dE'}{4\pi^2} \sum_{lr} \{ (E' - E - iD') b_{1lr}(E) b_{1lr}^*(E') \\ & - \frac{iD'}{2V_d} [(E' + E_{lr1}) + (E + E_{lr1})] b_{1lr}(E) b_{1lr}^*(E') \\ & + \frac{i\Lambda}{2V_d} [b_{2lr}(E) b_{1lr}^*(E') + b_{1lr}(E) b_{2lr}^*(E')] \\ & - \Omega_0 [b_{1lr}(E) b_{2lr}^*(E') - b_{2lr}(E) b_{1lr}^*(E')] \\ & - \Omega' [b_{1l}^*(E') b_{1lr}(E) - b_{1l}(E) b_{1lr}^*(E')] \} e^{i(E' - E)t} = 0. \end{aligned} \quad (5)$$

As shown in the Appendix, it can be derived that

$$\sum_{lr} \Omega' b_{1lr}(E) = -\frac{iD'}{2V_d}(V_d + E - E_1)b_1(E) + \frac{i\Lambda}{2V_d}b_2(E). \quad (6)$$

Using this result and Eqs. (4a) and (4c), we can obtain from Eq. (5), by keeping the leading terms, that

$$\begin{aligned} \dot{\sigma}_{11}^{(1)} = & -D'[\sigma_{11}^{(1)} - \sigma_{11}^{(0)}] + i\Omega_0[\sigma_{12}^{(1)} - \sigma_{21}^{(1)}] \\ & + \frac{\Lambda}{2V_d} \left(1 - \frac{\Omega'}{\Omega}\right) [\sigma_{12}^{(1)} + \sigma_{21}^{(1)} - \sigma_{12}^{(0)} - \sigma_{21}^{(0)}]. \end{aligned} \quad (7)$$

Similarly, Eq. (4d) gives

$$\begin{aligned} \dot{\sigma}_{22}^{(1)} = & -D[\sigma_{22}^{(1)} - \sigma_{22}^{(0)}] + i\Omega_0[\sigma_{21}^{(1)} - \sigma_{12}^{(1)}] \\ & + \frac{\Lambda}{2V_d} \left(1 - \frac{\Omega}{\Omega'}\right) [\sigma_{12}^{(1)} + \sigma_{21}^{(1)} - \sigma_{12}^{(0)} - \sigma_{21}^{(0)}]. \end{aligned} \quad (8)$$

To calculate the off-diagonal element σ_{12} , we subtract the product of Eq. (4c) and $b_{2lr}^*(E')$ by the product of the complex conjugate of Eq. (4d) after exchanging $E \leftrightarrow E'$ and $b_{1lr}(E)$. We obtain

$$\begin{aligned} \dot{\sigma}_{12}^{(1)} = & i(E_2 - E_1)\sigma_{12}^{(1)} - \frac{1}{2}(D' + D)\sigma_{12}^{(1)} + \sqrt{DD'}\sigma_{12}^{(0)} \\ & + i\Omega_0[\sigma_{11}^{(1)} - \sigma_{22}^{(1)}] - \frac{\Lambda}{2V_d} \left[\frac{\Omega'}{\Omega}\sigma_{11}^{(0)} + \frac{\Omega}{\Omega'}\sigma_{22}^{(0)} \right] \\ & + \frac{\Lambda}{2V_d} [\sigma_{11}^{(0)} + \sigma_{22}^{(0)}] - \frac{\Lambda}{2V_d} \left[\frac{\Omega}{\Omega'}\sigma_{11}^{(1)} + \frac{\Omega'}{\Omega}\sigma_{22}^{(1)} \right] \\ & + \frac{\Lambda}{2V_d} [\sigma_{11}^{(1)} + \sigma_{22}^{(1)}]. \end{aligned} \quad (9)$$

Similar procedures can be used for $\sigma_{ij}^{(n)}$ ($n \geq 2$) and the resulting equations are

$$\begin{aligned} \dot{\sigma}_{11}^{(n)} = & -D'[\sigma_{11}^{(n)} - \sigma_{11}^{(n-1)}] + i\Omega_0[\sigma_{12}^{(n)} - \sigma_{21}^{(n)}] \\ & + \frac{\Lambda}{2V_d} \left(1 - \frac{\Omega'}{\Omega}\right) [\sigma_{12}^{(n)} + \sigma_{21}^{(n)} - \sigma_{12}^{(n-1)} - \sigma_{21}^{(n-1)}], \\ \dot{\sigma}_{22}^{(n)} = & -D[\sigma_{22}^{(n)} - \sigma_{22}^{(n-1)}] + i\Omega_0[\sigma_{21}^{(n)} - \sigma_{12}^{(n)}] \\ & + \frac{\Lambda}{2V_d} \left(1 - \frac{\Omega}{\Omega'}\right) [\sigma_{12}^{(n)} + \sigma_{21}^{(n)} - \sigma_{12}^{(n-1)} - \sigma_{21}^{(n-1)}], \\ \dot{\sigma}_{12}^{(n)} = & i(E_2 - E_1)\sigma_{12}^{(n)} - \frac{1}{2}(D' + D)\sigma_{12}^{(n)} + \sqrt{DD'}\sigma_{12}^{(n-1)} \\ & + i\Omega_0[\sigma_{11}^{(n)} - \sigma_{22}^{(n)}] - \frac{\Lambda}{2V_d} \left[\frac{\Omega'}{\Omega}\sigma_{11}^{(n-1)} + \frac{\Omega}{\Omega'}\sigma_{22}^{(n-1)} \right] \\ & + \frac{\Lambda}{2V_d} [\sigma_{11}^{(n-1)} + \sigma_{22}^{(n-1)}] - \frac{\Lambda}{2V_d} \left[\frac{\Omega}{\Omega'}\sigma_{11}^{(n)} + \frac{\Omega'}{\Omega}\sigma_{22}^{(n)} \right] \\ & + \frac{\Lambda}{2V_d} [\sigma_{11}^{(n)} + \sigma_{22}^{(n)}]. \end{aligned} \quad (10)$$

Summing over n , we obtain the following Bloch-type rate




FIG. 2: (Color online) Occupation probability of the extra electron in each quantum dot as a function of time for $\Gamma_d = \Omega_0$ and (a) $\varepsilon = E_2 - E_1 = 0$, $\chi = 0$; (b) $\varepsilon = 0$, $\chi = 0.1\Omega_0$; (c) $\varepsilon = \Omega_0$, $\chi = 0$; and (d) $\varepsilon = \Omega_0$, $\chi = 0.1\Omega_0$.

equations for the density-matrix elements:

$$\begin{aligned} \dot{\sigma}_{11} &= i\Omega_0(\sigma_{12} - \sigma_{21}), \\ \dot{\sigma}_{22} &= i\Omega_0(\sigma_{21} - \sigma_{12}), \\ \dot{\sigma}_{12} &= i\varepsilon\sigma_{12} - \frac{\Gamma_d}{2}\sigma_{12} + i\Omega_0(\sigma_{11} - \sigma_{22}) \\ &\quad - \frac{\chi}{2}(\sigma_{11} + \sigma_{22}), \end{aligned} \quad (11)$$

with

$$\begin{aligned} \Gamma_d &= (\sqrt{D'} - \sqrt{D})^2, \\ \chi &= \left(\frac{\Lambda}{V_d}\right) \left(\frac{\Omega}{\Omega'} + \frac{\Omega'}{\Omega} - 2\right). \end{aligned} \quad (12)$$

Here $\sigma_{ij} = \sum_n \sigma_{ij}^{(n)}$ is the reduced density-matrix elements of the CQD system after tracing out the variables of the QPC, and $\varepsilon = E_2 - E_1$ is the energy-level difference of the two quantum dots. The decoherence rate Γ_d characterizes the exponential damping of the off-diagonal density-matrix element. These Bloch-type rate equations are the same as those from a theory based on master equations with the Markovian approximation [cf., Eq. (17) in Ref. 8]. When $\chi = 0$, these equations are reduced to the rate equations derived in Ref. 5(b), which ignores the effects of the higher-order terms.

III. DISCUSSION AND CONCLUSION

Figure 2 shows the time dependence of the occupation probability of the extra electron in each quantum dot. When $\chi = 0$, the occupation probability in each dot always decays to 0.5 at $t \rightarrow \infty$ [see Figs. (2a) and (2c)]. These results are identical to those in Ref. 5(b). However, after including higher order terms, χ becomes nonzero and is not negligible. This gives rise to an additional term $\frac{1}{2}\chi(\sigma_{11} + \sigma_{22})$ in the last equation of (11). From Fig. (2b), one can see that the occupation probabilities still decay to 0.5 as $t \rightarrow \infty$ for symmetrical CQD's, but decay more slowly than in the previous case with vanishing χ . Moreover, for asymmetrical CQD's, the occupation probability in each dot decays to a different value at $t \rightarrow \infty$ [see Fig. 2(d)]. This is reasonable because the two quantum dots are characterized by different parameters. Below we further explain these phenomena analytically.

Let us look at the stationary solution of Eq. (11) in the limit $t \rightarrow \infty$ so that $\dot{\sigma}_{ij}(t \rightarrow \infty) = 0$. Using $\sigma_{ij}(t \rightarrow \infty) = u_{ij} + iv_{ij}$, where u_{ij} and v_{ij} are real numbers, Eq. (11) can be rewritten as

$$\begin{aligned} 0 &= -2\Omega_0 v_{12} \\ 0 &= 2\Omega_0 v_{12} \\ 0 &= i\varepsilon(u_{12} + iv_{12}) - \frac{\Gamma_d}{2}(u_{12} + iv_{12}) \\ &\quad + i\Omega_0(\sigma_{11} - \sigma_{22}) - \frac{\chi}{2}. \end{aligned} \quad (13)$$

From these equations, one obtains

$$\sigma_{11} - \sigma_{22} = \frac{\chi\varepsilon}{\Omega_0\Gamma_d} = \frac{\varepsilon}{V_d}. \quad (14)$$

As expected, $\sigma_{11} = \sigma_{22}$ for $E_1 = E_2$ (i.e., $\varepsilon = 0$), while $\sigma_{11} \neq \sigma_{22}$ for asymmetrical CQD's with $E_1 \neq E_2$ (i.e., $\varepsilon \neq 0$).

In conclusion, we have presented a quantitative description of the dissipative dynamics of a CQD qubit con-

nected to a detector in the form of a QPC. Bloch-type rate equations for the reduced density matrix elements of the qubit are derived. In particular, the long-time probability distribution of the state of the qubit is found to depend on the energy levels of the single dot electron states. This corrects an earlier calculation which has predicted a distribution independent of the dot properties. The improvement results from taking into account higher order interaction terms in our analysis and it can be extended to other quantum systems.

APPENDIX: DERIVATION OF EQUATIONS FOR PROBABILITY AMPLITUDES

For the QPC, we assume that the hopping amplitude $\Omega_{lr}(E_l, E_r)$ between the left and right reservoirs depends only weakly on the energy levels E_l and E_r . The energy dependence is therefore neglected, i.e. $\Omega_{lr}(E_l, E_r) = \Omega$, and $\Omega'_{lr}(E_l, E_r) = \Omega'$. As high-order terms, $\sum_{l'r'} \Omega' b_{1l'r'r'}(E)$ and $\sum_{l'r'} \Omega b_{2l'r'r'}(E)$ in Eqs. (3c) and (3d) are neglected, they give

$$b_{1lr}(E) \approx \frac{1}{E + E_l - E_1 - E_r} \times \left[\Omega' b_1(E) + \frac{\Omega_0 \Omega' b_2(E)}{E + E_l - E_2 - E_r} \right]. \quad (A.1)$$

Because the energy levels in each electron reservoir of the QPC are dense, we can replace each sum over l and r in Eqs. (3a) and (3b) by an integral. For instance, $\sum_{lr} \rightarrow \int \int \rho_l(E_l) \rho_r(E_r) dE_l dE_r$, where $\rho_{l,r}$ is the density of states in the left and right reservoirs, respectively. This integral can be split into two parts: the principal and singular value parts. The principal part merely renormalizes the energy levels, while the singular value part plays an important role. When Eq. (A.1) is substituted into $\sum_{lr} \Omega' b_{1lr}(E)$, we thus obtain two terms. The first term is given by

$$\begin{aligned} \sum_{lr} \frac{\Omega'^2 b_1(E)}{E + E_l - E_1 - E_r} &= b_1(E) \int_{-\infty}^{\mu_L} dE_l \int_{\mu_R}^{+\infty} dE_r \frac{\rho_l(E_l) \rho_r(E_r) \Omega'^2}{E + E_l - E_1 - E_r} \\ &= b_1(E) \int_{-\infty}^{\mu_L} dE_l \int_{\mu_R}^{+\infty} dE_r \frac{\rho_l \rho_r \Omega'^2 (E + E_l - E_1 - E_r - i\eta)}{(E + E_l - E_1 - E_r)^2 + \eta^2} \\ &\approx b_1(E) \int_{-\infty}^{\mu_L} dE_l \int_{\mu_R}^{+\infty} dE_r (-i\pi \rho_l \rho_r \Omega'^2) \delta(E + E_l - E_1 - E_r) \\ &= -i \frac{D'}{2V_d} b_1(E) \int_{-\infty}^{\mu_L} dE_l \theta(E + E_l - E_1 - \mu_R) \\ &= -i \frac{D'}{2V_d} b_1(E) (V_d + E - E_1) \theta(V_d + E - E_1), \end{aligned} \quad (A.2)$$

where $\eta \rightarrow 0^+$, $V_d = \mu_L - \mu_R$, $D' = 2\pi \rho_l \rho_r \Omega'^2 V_d$, and $\theta(x)$ is the Heaviside step function. In the high voltage limit with $eV_d \gg \Omega^2 \rho$, the step function in Eq. (A.2) becomes one. Thus, the L.H.S. of Eq. (A.2) can be approximated by $-i(D'/2V_d)(V_d + E - E_1)b_1(E)$.

The second term is

$$\begin{aligned}
& \sum_{lr} \frac{\Omega' \Omega \Omega_0 b_2(E)}{(E + E_l - E_1 - E_r)(E + E_l - E_2 - E_r)} \\
&= b_2(E) \int_{-\infty}^{\mu_L} dE_l \int_{\mu_R}^{+\infty} dE_r \frac{\rho_l(E_l) \rho_r(E_r) \Omega' \Omega \Omega_0}{(E + E_l - E_1 - E_r)(E + E_l - E_2 - E_r)} \\
&= b_2(E) \int_{-\infty}^{\mu_L} dE_l \int_{\mu_R}^{+\infty} dE_r \rho_l \rho_r \Omega' \Omega \Omega_0 \frac{(E + E_l - E_1 - E_r - i\eta)(E + E_l - E_2 - E_r - i\eta)}{[(E + E_l - E_1 - E_r)^2 + \eta^2][(E + E_l - E_2 - E_r)^2 + \eta^2]} \\
&\approx b_2(E) \int_{-\infty}^{\mu_L} dE_l \int_{\mu_R}^{+\infty} dE_r (-i\pi \rho_l \rho_r \Omega' \Omega \Omega_0) \left\{ \frac{E + E_l - E_1 - E_r}{(E + E_l - E_1 - E_r)^2 + \eta^2} \delta(E + E_l - E_2 - E_r) \right. \\
&\quad \left. + \frac{E + E_l - E_2 - E_r}{(E + E_l - E_2 - E_r)^2 + \eta^2} \delta(E + E_l - E_1 - E_r) \right\} \\
&= -i \frac{\Lambda}{2V_d} b_2(E) \int_{-\infty}^{\mu_L} dE_l \frac{E_2 - E_1}{(E_2 - E_1)^2 + \eta^2} [\theta(E + E_l - E_2 - \mu_R) - \theta(E + E_l - E_1 - \mu_R)] \\
&= -i \frac{\Lambda}{2V_d} b_2(E) \frac{E_2 - E_1}{(E_2 - E_1)^2 + \eta^2} [(V_d + E - E_2)\theta(V_d + E - E_2) - (V_d + E - E_1)\theta(V_d + E - E_1)],
\end{aligned} \tag{A.3}$$

where $\Lambda = 2\pi\rho_l\rho_r\Omega'\Omega_0\Omega V_d$. In the high voltage limit, the two step functions in Eq. (A.3) become one and the L.H.S. of Eq. (A.3) is finally approximated by $i(\Lambda/2V_d)b_2(E)$. Thus, the sum of these two terms gives Eq. (6) in the high voltage limit. Substituting Eq. (6) into Eq. (3a), one obtains Eq. (4a). Also, one can derive Eqs. (4b)-(4d) using similar procedures.

ACKNOWLEDGMENTS

We would like to thank S. A. Gurvitz and Xinqi Li for valuable discussions. This work was supported by the National Natural Science Foundation of China Grant Nos. 10474013 and 10534060.

-
- ¹ M. Namiki, S. Pascazio, and H. Nakazato, *Decoherence and Quantum measurements* (World Scientific, Singapore, 1997).
² A. Yacoby, M. Heiblum, V. Umansky, H. Strikman, and D. Mahalu, Phys. Rev. Lett. **74**, 4047 (1995).
³ R. Shuster, E. Buks, M. Heiblum, D. Mahalu, V. Umansky, and H. Shtrikman, Nature (London) **385**, 417 (1997).
⁴ E. Buks, R. Schuster, M. Heiblum, D. Mahalu, V. Umansky, and H. Shtrikman, Phys. Rev. Lett. **77**, 4664 (1996).

- ⁵ (a) S. A. Gurvitz and Ya. S. Prager, Phys. Rev. B **53**, 15932 (1996); (b) S. A. Gurvitz, Phys. Rev. B **56**, 15215 (1997).
⁶ S. A. Gurvitz, L. Fedichkin, D. Mozyrsky, and G. P. Berman, Phys. Rev. Lett. **91**, 066801 (2003).
⁷ H. S. Goan, G. J. Milburn, H. M. Wiseman, and H. B. Sun, Phys. Rev. B **63**, 125326 (2001).
⁸ Xinqi Li, Wenkai Zhang, Ping Cui, Jiushu Shao, Zhongshui Ma, and Yijing Yan, Phys. Rev. B **69**, 085315 (2004).