ATLAS SUSY Searches* - 95% CL Lower Limits

ATLAS Preliminary

July 2018 $\sqrt{s} = 7, 8, 13 \text{ TeV}$ e, μ, τ, γ Jets $E_{\mathrm{T}}^{\mathrm{miss}}$ $\int \mathcal{L} \, dt [\mathrm{fb}^{-1}]$ Model **Mass limit** $\sqrt{s} = 7.8 \text{ TeV}$ $\sqrt{s} = 13 \text{ TeV}$ Reference $\tilde{q}\tilde{q}, \tilde{q} \rightarrow q\tilde{\chi}_1^0$ 1.55 1712.02332 2-6 jets Yes 36.1 [2x, 8x Degen. 0.9 $m(\tilde{\chi}_1^0) < 100 \text{ GeV}$ mono-jet 1-3 jets Yes 36.1 [1x, 8x Degen.] 0.43 0.71 $m(\tilde{q})-m(\tilde{\chi}_1^0)=5 \text{ GeV}$ 1711.03301 Inclusive Searches $\tilde{g}\tilde{g}, \, \tilde{g} \rightarrow q\bar{q}\tilde{\chi}_1^0$ 2-6 jets Yes 36.1 2.0 $m(\tilde{\chi}_{1}^{0})$ <200 GeV 1712.02332 0 0.95-1.6 Forbidden $m(\tilde{\chi}_{1}^{0}) = 900 \,\text{GeV}$ 1712.02332 4 jets $3e, \mu$ 36.1 1.85 $m(\tilde{\chi}_{1}^{0}) < 800 \,\text{GeV}$ 1706.03731 $\tilde{g}\tilde{g}, \tilde{g} \rightarrow g\bar{g}(\ell\ell)\tilde{\chi}_1^0$ ee, µµ 2 jets Yes 36.1 1.2 $m(\tilde{g})-m(\tilde{\tilde{\chi}}_1^0)=50 \text{ GeV}$ 1805.11381 $\tilde{g}\tilde{g}, \, \tilde{g} \rightarrow qqWZ\tilde{\chi}_1^0$ 7-11 jets 1.8 1708.02794 0 Yes 36.1 $m(\tilde{\chi}_1^0)$ <400 GeV 0.98 $3e, \mu$ 4 jets 36.1 $m(\tilde{g})-m(\tilde{\chi}_1^0)=200 \text{ GeV}$ 1706.03731 $\tilde{g}\tilde{g}, \, \tilde{g} \rightarrow t\bar{t}\tilde{\chi}_1^0$ 0-1 e, μ 2.0 3 b Yes 36.1 $m(\tilde{\chi}_1^0)$ <200 GeV 1711.01901 $3e, \mu$ 4 jets 36.1 1.25 $m(\tilde{g})-m(\tilde{\chi}_1^0)=300 \text{ GeV}$ 1706.03731 $\tilde{b}_1\tilde{b}_1,\,\tilde{b}_1{\rightarrow}b\tilde{\chi}_1^0/t\tilde{\chi}_1^{\pm}$ Multiple 36.1 \tilde{b}_1 Forbidden 0.9 $m(\tilde{\chi}_{1}^{0})=300 \text{ GeV}, BR(b\tilde{\chi}_{1}^{0})=1$ 1708.09266, 1711.03301 Multiple \tilde{b}_1 \tilde{b}_1 36.1 Forbidden 0.58-0.82 $m(\tilde{\chi}_1^0) = 300 \text{ GeV}, BR(b\tilde{\chi}_1^0) = BR(t\tilde{\chi}_1^{\pm}) = 0.5$ 1708.09266 Multiple 36.1 Forbidden 0.7 $m(\tilde{\chi}_{\perp}^{0}) = 200 \text{ GeV}, m(\tilde{\chi}_{\perp}^{\pm}) = 300 \text{ GeV}, BR(t\tilde{\chi}_{\perp}^{\pm}) = 1$ 1706.03731 Multiple $\tilde{b}_1 \tilde{b}_1, \tilde{t}_1 \tilde{t}_1, M_2 = 2 \times M_1$ 36.1 0.7 $m(\tilde{\chi}_1^0)=60 \text{ GeV}$ 1709.04183, 1711.11520, 1708.03247 \tilde{t}_1 Multiple 36.1 Forbidden 0.9 1709.04183, 1711.11520, 1708.03247 $m(\tilde{\chi}_{1}^{0})=200 \,\text{GeV}$ $\tilde{t}_1\tilde{t}_1, \, \tilde{t}_1 {\rightarrow} Wb\tilde{\chi}^0_1 \text{ or } t\tilde{\chi}^0_1$ 0-2 e, μ 0-2 jets/1-2 b Yes \tilde{t}_1 36.1 1.0 $m(\tilde{\chi}_1^0)=1 \text{ GeV}$ 1506.08616, 1709.04183, 1711.11520 $ilde{t}_1 ilde{t}_1,\, ilde{H}\,\mathsf{LSP}$ Multiple \tilde{t}_1 0.4-0.9 36.1 $m(\tilde{\chi}_1^0)=150 \text{ GeV}, m(\tilde{\chi}_1^{\pm})-m(\tilde{\chi}_1^0)=5 \text{ GeV}, \tilde{t}_1 \approx \tilde{t}_L$ 1709.04183, 1711.11520 3rd gen. direct pro Multiple Forbidden 0.6-0.8 $m(\tilde{\chi}_1^0)=300 \text{ GeV}, m(\tilde{\chi}_1^{\pm})-m(\tilde{\chi}_1^0)=5 \text{ GeV}, \tilde{t}_1 \approx \tilde{t}_L$ 1709.04183, 1711.11520 36.1 \tilde{t}_1 $\tilde{t}_1\tilde{t}_1$, Well-Tempered LSP Multiple \tilde{t}_1 0.48-0.84 $m(\tilde{\chi}_1^0)$ =150 GeV, $m(\tilde{\chi}_1^{\pm})$ - $m(\tilde{\chi}_1^0)$ =5 GeV, $\tilde{t}_1 \approx \tilde{t}_L$ 1709.04183, 1711.11520 36.1 $m(\tilde{\chi}_{1}^{0})=0 \text{ GeV}$ $m(\tilde{t}_{1},\tilde{c})-m(\tilde{\chi}_{1}^{0})=50 \text{ GeV}$ $\tilde{t}_1 \tilde{t}_1, \tilde{t}_1 \rightarrow c \tilde{\chi}_1^0 / \tilde{c} \tilde{c}, \tilde{c} \rightarrow c \tilde{\chi}_1^0$ 0 2cYes 36.1 0.85 1805.01649 0.46 1805.01649 $m(\tilde{t}_1,\tilde{c})-m(\tilde{\chi}_1^0)=5 \text{ GeV}$ 0 mono-jet Yes 36.1 0.43 1711.03301 $\tilde{t}_2\tilde{t}_2, \tilde{t}_2 \rightarrow \tilde{t}_1 + h$ 1-2 e, μ 4 b 36.1 0.32-0.88 $m(\tilde{\chi}_1^0)=0$ GeV, $m(\tilde{t}_1)-m(\tilde{\chi}_1^0)=180$ GeV 1706.03986 Yes \tilde{t}_2 $\begin{array}{c} \tilde{\chi}_1^{\pm}/\tilde{\chi}_2^0 \\ \tilde{\chi}_1^{\pm}/\tilde{\chi}_2^0 \end{array}$ $\tilde{\chi}_1^{\pm} \tilde{\chi}_2^0$ via WZ0.6 2-3 e, μ Yes 36.1 $m(\tilde{\chi}_1^0)=0$ 1403.5294, 1806.02293 ee, µµ ≥ 1 Yes 36.1 0.17 $m(\tilde{\chi}_1^{\pm})-m(\tilde{\chi}_1^{0})=10 \text{ GeV}$ 1712.08119 $\tilde{\chi}_{1}^{\pm}\tilde{\chi}_{2}^{0}$ via Wh $\ell\ell/\ell\gamma\gamma/\ell bb$ $\tilde{\chi}_1^{\pm}/\tilde{\chi}_2^0$ 0.26 $m(\tilde{\chi}_1^0)=0$ 1501.07110 Yes 20.3 $\begin{array}{c} \tilde{\chi}_1^{\pm}/\tilde{\chi}_2^0 \\ \tilde{\chi}_1^{\pm}/\tilde{\chi}_2^0 \end{array}$ $\tilde{\chi}_{1}^{\pm}\tilde{\chi}_{1}^{\mp}/\tilde{\chi}_{2}^{0}, \tilde{\chi}_{1}^{+} \rightarrow \tilde{\tau}\nu(\tau\tilde{\nu}), \tilde{\chi}_{2}^{0} \rightarrow \tilde{\tau}\tau(\nu\tilde{\nu})$ 2 τ Yes 36.1 0.76 $m(\tilde{\chi}_{1}^{0})=0, m(\tilde{\tau}, \tilde{\nu})=0.5(m(\tilde{\chi}_{1}^{\pm})+m(\tilde{\chi}_{1}^{0}))$ 1708 07875 0.22 $m(\tilde{\chi}_{1}^{\pm})-m(\tilde{\chi}_{1}^{0})=100 \text{ GeV}, m(\tilde{\tau}, \tilde{\nu})=0.5(m(\tilde{\chi}_{1}^{\pm})+m(\tilde{\chi}_{1}^{0}))$ 1708.07875 $\tilde{\ell}_{L,R}\tilde{\ell}_{L,R}, \, \tilde{\ell} \rightarrow \ell \tilde{\chi}_1^0$ $2e, \mu$ 0 Yes 36.1 0.5 $m(\tilde{\chi}_1^0)=0$ 1803.02762 $2e, \mu$ Yes 36.1 0.18 1712.08119 ≥ 1 $m(\tilde{\ell})-m(\tilde{\chi}_{1}^{0})=5 \text{ GeV}$ $\tilde{H}\tilde{H}, \tilde{H} \rightarrow h\tilde{G}/Z\tilde{G}$ 0 36.1 0.13-0.23 0.29-0.88 $\geq 3b$ Yes Ĥ $BR(\tilde{\chi}_1^0 \to h\tilde{G})=1$ 1806.04030 $4e, \mu$ 36.1 0.3 0 Yes \tilde{H} $BR(\tilde{\chi}_1^0 \to Z\tilde{G})=1$ 1804.03602 Direct $\tilde{\chi}_1^+ \tilde{\chi}_1^-$ prod., long-lived $\tilde{\chi}_1^{\pm}$ Disapp. trk 1 jet Yes 0.46 36.1 Pure Wino 1712.02118 0.15 Pure Higgsino ATL-PHYS-PUB-2017-019 Stable § R-hadron SMP 1606.05129 3.2 1.6 $[\tau(\tilde{g}) = 100 \text{ ns}, 0.2 \text{ ns}]$ Metastable \tilde{g} R-hadron, $\tilde{g} \rightarrow qq\tilde{\chi}_1^0$ Multiple 32.8 1.6 2.4 1710.04901, 1604.04520 $m(\tilde{\chi}_1^0)=100 \text{ GeV}$ GMSB, $\tilde{\chi}_1^0 \rightarrow \gamma \tilde{G}$, long-lived $\tilde{\chi}_1^0$ 0.44 2γ 20.3 $1 < \tau(\tilde{\chi}_1^0) < 3$ ns. SPS8 model 1409.5542 Yes $\tilde{g}\tilde{g}, \tilde{\chi}_1^0 \rightarrow eev/e\mu v/\mu\mu v$ displ. $ee/e\mu/\mu\mu$ 20.3 1.3 $6 < c\tau(\tilde{\chi}_1^0) < 1000 \text{ mm, m}(\tilde{\chi}_1^0) = 1 \text{ TeV}$ 1504.05162 LFV $pp \rightarrow \tilde{v}_{\tau} + X, \tilde{v}_{\tau} \rightarrow e\mu/e\tau/\mu\tau$ $e\mu$, $e\tau$, $\mu\tau$ 3.2 1.9 λ'_{311} =0.11, $\lambda_{132/133/233}$ =0.07 1607.08079 $\tilde{\chi}_{1}^{\pm}\tilde{\chi}_{1}^{\mp}/\tilde{\chi}_{2}^{0} \rightarrow WW/Z\ell\ell\ell\ell\nu\nu$ $4e, \mu$ 0 Yes 36.1 $[\lambda_{i33} \neq 0, \lambda_{12k} \neq 0]$ 0.82 1.33 $m(\tilde{\chi}_1^0)=100 \text{ GeV}$ 1804.03602 $\tilde{g}\tilde{g}, \, \tilde{g} \rightarrow qq\tilde{\chi}_1^0, \, \tilde{\chi}_1^0 \rightarrow qqq$ Large $\lambda_{112}^{\prime\prime}$ 4-5 large-R jets -36.1 $m(\tilde{\chi}_1^0) = 200 \text{ GeV}, 1100 \text{ GeV}]$ $\chi_{112}^0 = 2e-4, 2e-5]$ 1.3 1.9 1804.03568 Multiple 36.1 1.05 2.0 $m(\tilde{\chi}_1^0)$ =200 GeV, bino-like ATLAS-CONF-2018-003 $\tilde{g}\tilde{g}, \, \tilde{g} \to tbs \, / \, \tilde{g} \to t\bar{t}\tilde{\chi}_1^0, \, \tilde{\chi}_1^0 \to tbs$ Multiple 36.1 1.8 2.1 $m(\tilde{\chi}_1^0)=200$ GeV. bino-like ATLAS-CONF-2018-003 $\tilde{t}\tilde{t}, \tilde{t} \rightarrow t\tilde{\chi}_1^0, \tilde{\chi}_1^0 \rightarrow tbs$ $[\lambda_{222}^{"}=2e-4, 1e-2]$ 0.55 1.05 Multiple 36.1 $m(\tilde{\chi}_1^0)$ =200 GeV. bino-like ATLAS-CONF-2018-003 $\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow bs$ 2 jets + 2 b 0 36.7 [qq, bs]0.42 0.61 1710.07171 $\tilde{t}_1\tilde{t}_1, \tilde{t}_1 \rightarrow b\ell$ $2e, \mu$ 0.4-1.45 2 b 36.1 BR($\tilde{t}_1 \rightarrow be/b\mu$)>20% 1710.05544

 10^{-1}

^{*}Only a selection of the available mass limits on new states or phenomena is shown. Many of the limits are based on simplified models, c.f. refs. for the assumptions made.