Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Metropolis within Gibbs Algorithm: Application to Linear Geostatistical Model

Johnson, Olatunji Olugoke

o.johnson@lanacster.ac.uk

May 22, 2017

Outline

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introductio

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recomIntroduction

2 Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

4 Limitations, Conclusion and recommendation

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistica Model

Hastings-Within-Gibb Algorithm

Limitations, Conclusion and recommendation

What is Markov Chain Monte Carlo (MCMC)?

Markov Chain Monte Carlo (MCMC) methods are a class of algorithms for sampling from a probability distribution based on constructing a Markov chain that has the desired distribution as its stationary distribution.

What is the goal of MCMC?

The goal of MCMC is to draw samples from some probability distribution without having to know its exact **height at any point**. The way MCMC achieves this is to "wander around" on that distribution in such a way that the amount of time spent in each location is proportional to the height of the distribution. If the "wandering around" process is set up correctly, you can make sure that this proportionality (between time spent and height of the distribution) is achieved.

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations Conclusion and recom-

Applications of MCMC

- Sampling from any distribution.
- Heuristic way of Minimizing/Maximizing any function.
- Orytography: Breaking the code.
- Finding out the best arrangement of a DNA sequence
- 6

- Metropolis-Hasting Algorithm
- @ Gibbs Sampling

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations Conclusion and recommendation

Applications of MCMC

- Sampling from any distribution.
- **②** Heuristic way of **Minimizing/Maximizing any function**.
- Crytography: Breaking the code.
- Finding out the best arrangement of a DNA sequence.
- 5

- Metropolis-Hasting Algorithm
- @ Gibbs Sampling

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations Conclusion and recommendation

Applications of MCMC

- Sampling from any distribution.
- **②** Heuristic way of **Minimizing/Maximizing any function**.
- **3** Crytography: Breaking the code.
- Finding out the best arrangement of a DNA sequence.
- 5

- Metropolis-Hasting Algorithm
- @ Gibbs Sampling

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations Conclusion and recommendation

Applications of MCMC

- Sampling from any distribution.
- **②** Heuristic way of **Minimizing/Maximizing any function**.
- **3** Crytography: Breaking the code.
- Finding out the best arrangement of a DNA sequence.
- 5

- Metropolis-Hasting Algorithm
- @ Gibbs Sampling

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations Conclusion and recommendation

Applications of MCMC

- Sampling from any distribution.
- We Heuristic way of Minimizing/Maximizing any function.
- **3 Crytography**: Breaking the code.
- **1** Finding out the **best arrangement of a DNA sequence**.
- **3** ?

- Metropolis-Hasting Algorithm
- @ Gibbs Sampling

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations Conclusion and recommendation

Applications of MCMC

- Sampling from any distribution.
- Wear Heuristic way of Minimizing/Maximizing any function.
- **3** Crytography: Breaking the code.
- Finding out the **best arrangement of a DNA sequence**.
- **3** ?

- Metropolis-Hasting Algorithm
- Q Gibbs Sampling

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations Conclusion and recommendation

Applications of MCMC

- Sampling from any distribution.
- We Heuristic way of Minimizing/Maximizing any function.
- **3 Crytography**: Breaking the code.
- Finding out the **best arrangement of a DNA sequence**.
- **3** ?

- Metropolis-Hasting Algorithm
- @ Gibbs Sampling

Fundamental Mechanisms of MCMC

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Gibbs sampler

Given the target multivariate distribution $\pi(x) = \pi(x_1, \dots, x_p)$, the Gibbs samplers successively and repeatedly generates samples for each of the random variables, X_i , from the full conditional distribution $(X_i|X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_p)$.

Metropolis-Hastings algorithm

The Metropolis-Hastings (MH) algorithm simulates samples from a probability distribution by making use of the full joint density function and (independent) proposal distributions for each of the variables of interest.

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

- **•** RANDOM WALK PROPOSAL: $q(x^*, x) = q(x^*, -x)$
 - ② INDEPENDENCE PROPOSAL: $q(x^*, x) = q(y)$
- **3** TAILORED PROPOSAL: $q(\cdot) = f(\cdot | mode, H^{-1})$
- 4

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

- **•** RANDOM WALK PROPOSAL: $q(x^*, x) = q(x^*, -x)$
- **2** INDEPENDENCE PROPOSAL: $q(x^*, x) = q(y)$
- 3 TAILORED PROPOSAL: $q(\cdot) = f(\cdot|mode, H^{-1})$
- 4

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

- **1** RANDOM WALK PROPOSAL: $q(x^*, x) = q(x^*, -x)$
- ② INDEPENDENCE PROPOSAL: $q(x^*, x) = q(y)$
- **3** TAILORED PROPOSAL: $q(\cdot) = f(\cdot|mode, H^{-1})$
 - 4

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

- **1** RANDOM WALK PROPOSAL: $q(x^*, x) = q(x^*, -x)$
- ② INDEPENDENCE PROPOSAL: $q(x^*, x) = q(y)$
- **3** TAILORED PROPOSAL: $q(\cdot) = f(\cdot|mode, H^{-1})$
- **4** ?

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations Conclusion and recommendation

Classical geostatistical Model

$$Y(x_i) = d^{\top}(x_i)\beta + S(x_i) + Z(x_i)$$

Hierarchical framework

- ① Data Model: $Y|\beta, S(x), \tau^2 \sim MVN(D\beta + S(x), \tau^2 I)$
- Process Model: $S(x)|\theta \sim MVN(0, \Sigma(\theta))$
- O Parameter Model: $[\beta, \tau^2, \theta]$

- Data Model: $Y|\beta, \theta, \tau^2 \sim MVN(D\beta, \Sigma(\theta) + \tau^2 I)$
- **Parameter Model:** $[\beta, \tau^2, \theta]$

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Classical geostatistical Model

$$Y(x_i) = d^{\top}(x_i)\beta + S(x_i) + Z(x_i)$$

Hierarchical framework

- **1** Data Model: $Y|\beta, S(x), \tau^2 \sim MVN(D\beta + S(x), \tau^2 I)$
- **2** Process Model: $S(x)|\theta \sim MVN(0, \Sigma(\theta))$
- ③ Parameter Model: $[\beta, \tau^2, \theta]$

- ① Data Model: $Y|\beta, \theta, \tau^2 \sim MVN(D\beta, \Sigma(\theta) + \tau^2 I)$
- ② Parameter Model: $[\beta, \tau^2, \theta]$

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Classical geostatistical Model

$$Y(x_i) = d^{\top}(x_i)\beta + S(x_i) + Z(x_i)$$

Hierarchical framework

- **1** Data Model: $Y|\beta, S(x), \tau^2 \sim MVN(D\beta + S(x), \tau^2 I)$
- **2** Process Model: $S(x)|\theta \sim MVN(0, \Sigma(\theta))$
- **3** Parameter Model: $[\beta, \tau^2, \theta]$

- ① Data Model: $Y|\beta, \theta, \tau^2 \sim MVN(D\beta, \Sigma(\theta) + \tau^2 I)$
- ② Parameter Model: $[\beta, \tau^2, \theta]$

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Classical geostatistical Model

$$Y(x_i) = d^{\top}(x_i)\beta + S(x_i) + Z(x_i)$$

Hierarchical framework

- **1** Data Model: $Y|\beta, S(x), \tau^2 \sim MVN(D\beta + S(x), \tau^2 I)$
- **2** Process Model: $S(x)|\theta \sim MVN(0, \Sigma(\theta))$
- **3** Parameter Model: $[\beta, \tau^2, \theta]$

- **1** Data Model: $Y|\beta, \theta, \tau^2 \sim MVN(D\beta, \Sigma(\theta) + \tau^2 I)$
- 2 Parameter Model: $[\beta, \tau^2, \theta]$

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Classical geostatistical Model

$$Y(x_i) = d^{\top}(x_i)\beta + S(x_i) + Z(x_i)$$

Hierarchical framework

- **1** Data Model: $Y|\beta, S(x), \tau^2 \sim MVN(D\beta + S(x), \tau^2 I)$
- **2** Process Model: $S(x)|\theta \sim MVN(0, \Sigma(\theta))$
- **3** Parameter Model: $[\beta, \tau^2, \theta]$

- **1** Data Model: $Y|\beta, \theta, \tau^2 \sim MVN(D\beta, \Sigma(\theta) + \tau^2 I)$
- 2 Parameter Model: $[\beta, \tau^2, \theta]$

Full conditional distributions

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recom-

Set of full conditionals of the posterior

- $\beta|y,\tau^2,\theta\sim N(M,V)$
- $\theta|y,\beta,\theta\propto f(y|\beta,\theta,\tau^2)\pi(\beta,\tau^2,\theta)$
- $\tau^2 | y, \beta, \theta \propto f(y | \beta, \theta, \tau^2) \pi(\beta, \theta, \tau^2)$

where.

- $\pi(\beta) \sim N(\mu_{\beta}, \Sigma_{\beta})$
- $\pi(\theta) \sim IG(a,b)$
- $\pi(\tau^2) \sim IG(c,d)$
- $V = \left(\Sigma_{\beta}^{-1} + D'\Sigma_{\nu|\cdot}^{-1}D\right)^{-1}$
- $M = V\left(\Sigma_{\beta}^{-1}\mu_{\beta} + D'\Sigma_{\nu|\cdot}^{-1}y\right)$

Full conditional distributions

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recom-

Set of full conditionals of the posterior

- $\beta|y,\tau^2,\theta\sim N(M,V)$
- $\theta|y,\beta,\theta\propto f(y|\beta,\theta,\tau^2)\pi(\beta,\tau^2,\theta)$
- $\tau^2 | y, \beta, \theta \propto f(y | \beta, \theta, \tau^2) \pi(\beta, \theta, \tau^2)$

where.

- $\pi(\beta) \sim N(\mu_{\beta}, \Sigma_{\beta})$
- $\pi(\theta) \sim IG(a,b)$
- $\pi(\tau^2) \sim IG(c,d)$
- $\bullet \ V = \left(\Sigma_{\beta}^{-1} + D' \Sigma_{\gamma|\cdot}^{-1} D\right)^{-1}$
- $M = V\left(\Sigma_{\beta}^{-1}\mu_{\beta} + D'\Sigma_{\gamma|\cdot}^{-1}y\right)$

Full conditional distributions

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistica Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Set of full conditionals of the posterior

- $\beta|y,\tau^2,\theta\sim N(M,V)$
- $\theta|y,\beta,\theta\propto f(y|\beta,\theta,\tau^2)\pi(\beta,\tau^2,\theta)$
- $\tau^2 | y, \beta, \theta \propto f(y | \beta, \theta, \tau^2) \pi(\beta, \theta, \tau^2)$

where.

- $\pi(\beta) \sim N(\mu_{\beta}, \Sigma_{\beta})$
- $\pi(\theta) \sim IG(a,b)$
- $\pi(\tau^2) \sim IG(c,d)$
- $V = \left(\Sigma_{\beta}^{-1} + D'\Sigma_{\nu|\cdot}^{-1}D\right)^{-1}$
- $\bullet \ M = V \left(\Sigma_{\beta}^{-1} \mu_{\beta} + D' \Sigma_{\gamma|\cdot}^{-1} y \right)$

Limitation

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Limitation

- Hastings-within-Gibbs algorithm requires a lot of computational work.
- 2 slow mixing.
- inaccurate.

Limitation

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Limitation

- Hastings-within-Gibbs algorithm requires a lot of computational work.
- slow mixing.
- inaccurate.

Limitation

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Limitation

- Hastings-within-Gibbs algorithm requires a lot of computational work.
- slow mixing.
- 3 inaccurate.

Conclusions

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introductio

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Conclusions

- slow mixing and strong autocorrelation: the chains for β and the process would have strong autocorrelation.
- ② significant dependence: cross-correlations among the covariance parameters θ .
- 3 It is a good start for a new problem

Conclusions

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introductio

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Conclusions

- slow mixing and strong autocorrelation: the chains for β and the process would have strong autocorrelation.
- 2 significant dependence: cross-correlations among the covariance parameters θ .
- It is a good start for a new problem

Conclusions

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introductio

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Conclusions

- slow mixing and strong autocorrelation: the chains for β and the process would have strong autocorrelation.
- 2 significant dependence: cross-correlations among the covariance parameters θ .
- 3 It is a good start for a new problem

Recommendations

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introductio

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Recommendations

- Some properties must be study with care before conclusion.
- ② Ensure that the conditional distribution is derived from a valid joint distribution, that is the joint distribution must have a finite integral (though not always easy to validate

Recommendations

Markov Chain Monte Carlo

> Johnson, Olatunji Olugoke

Introductio

Hierarchical Gaussian Geostatistical Model

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recommendation

Recommendations

- Some properties must be study with care before conclusion.
- Ensure that the conditional distribution is derived from a valid joint distribution, that is the joint distribution must have a finite integral (though not always easy to validate).

Markov Chain Monte Carlo

Johnson, Olatunji Olugoke

Introduction

Hierarchical Gaussian Geostatistical

Hastings-Within-Gibbs Algorithm

Limitations, Conclusion and recom-

THANKS FOR YOUR ATTENTION !!!