

## STAT 610: Discussion 5

### 1 Summary

- Hypothesis test: after observing samples  $\mathbf{X} \sim \mathbb{P}_\theta$ , decide either to retain  $H_0 : \theta \in \Theta_0$  (null hypothesis) or to reject  $H_0$  and conclude  $H_1 : \theta \in \Theta_0^c$  (alternative hypothesis) is true.
- Basic definitions:
  - Type I error: reject  $H_0$  if  $H_0$  is true.
  - Type II error: retain  $H_0$  if  $H_0$  is false.
  - Let  $R$  denote the rejection region for a test. Then

$$\beta(\theta) = \mathbb{P}_\theta(\mathbf{X} \in R) = \begin{cases} \mathbb{P}(\text{Type I error}) & \text{if } \theta \in \Theta_0 \\ 1 - \mathbb{P}(\text{Type II error}) & \text{if } \theta \in \Theta_0^c \end{cases}$$

is called the power function.

- A test with power function  $\beta(\theta)$  is a size (level)  $\alpha$  test if  $\sup_{\theta \in \Theta_0} \beta(\theta) = \alpha$  ( $\leq \alpha$ ).
- The likelihood ratio test statistic for  $H_0 : \theta \in \Theta_0$  v.s.  $H_1 : \theta \in \Theta_0^c$  is

$$\lambda(\mathbf{x}) = \frac{\sup_{\theta \in \Theta_0} L(\theta | \mathbf{x})}{\sup_{\theta \in \Theta} L(\theta | \mathbf{x})}.$$

A *likelihood ratio test* (LRT) is any test with rejection region  $\{\mathbf{x} : \lambda(\mathbf{x}) \leq c\}$ , where  $0 \leq c \leq 1$ . Normally, we choose  $c$  such that the test

$$T(\mathbf{X}) = \begin{cases} 1 & \text{if } \lambda(\mathbf{X}) \leq c \\ 0 & \text{if } \lambda(\mathbf{X}) > c \end{cases}$$

has size  $\alpha$ , that is,  $\sup_{\theta \in \Theta_0} \beta(\theta) = \sup_{\theta \in \Theta_0} \mathbb{P}_\theta(\lambda(\mathbf{X}) \leq c) = \alpha$ .

- **Theorem 8.2.4.** If  $W(\mathbf{X})$  is a sufficient statistic for  $\theta$  and  $\lambda^*(w)$  and  $\lambda(\mathbf{x})$  are the LRT statistic based on  $W$  and  $\mathbf{X}$ , then  $\lambda^*(W(\mathbf{x})) = \lambda(\mathbf{x})$ .
- A test  $T^*$  of size  $\alpha$  is a *uniformly most powerful* (UMP) test if and only if  $\beta_{T^*}(\theta) \geq \beta_T(\theta)$  for all  $\theta \in \Theta_0^c$  and  $T$  of level  $\alpha$ .
- **Theorem 8.3.12. (Neyman-Pearson Lemma)** Consider  $H_0 : \theta = \theta_0$  v.s.  $H_1 : \theta = \theta_1$ , then

$$T^*(\mathbf{X}) = \begin{cases} 1 & f(\mathbf{X} | \theta_1) > cf(\mathbf{X} | \theta_0) \\ 0 & f(\mathbf{X} | \theta_1) < cf(\mathbf{X} | \theta_0) \end{cases},$$

is a UMP test of size  $\alpha_c := E(T^* | \theta_0)$ . Note that  $f(\mathbf{X} | \theta_1) = cf(\mathbf{X} | \theta_0)$  can be arbitrary.

## 2 Questions

1. Suppose that we observe  $m$  i.i.d  $\text{Ber}(\theta)$  random variables, denoted by  $Y_1, \dots, Y_m$ . Show that the LRT of  $H_0 : \theta \leq \theta_0$  v.s.  $H_1 : \theta > \theta_0$  will reject  $H_0$  if  $\sum_{i=1}^m Y_i > b$ .

2. A random sample  $X_1, \dots, X_n$  is drawn from a Pareto distribution with pdf

$$f(x | \theta, \nu) = \frac{\theta \nu^\theta}{x^{\theta+1}} I_{[\nu, \infty)}(x), \quad \theta > 0, \nu > 0.$$

- (a) Find the MLEs of  $\theta$  and  $\nu$ .  
(b) Show that the LRT of

$$H_0 : \theta = 1, \quad \text{v.s.} \quad H_1 : \theta \neq 1, \quad \nu \text{ unknown},$$

has reject region of the form  $\{\mathbf{x} : T(\mathbf{x}) \leq c_1 \text{ or } T(\mathbf{x}) \geq c_2\}$ , where  $0 < c_1 < c_2$  and

$$T = \log \left[ \frac{\prod_{i=1}^n X_i}{X_{(1)}^n} \right].$$

3. Suppose  $X$  is one observation from a population with distribution  $\text{Beta}(\theta, 1)$ .
- (a) For testing  $H_0 : \theta \leq 1$  v.s.  $H_1 : \theta > 1$ , find the size and the power function of the test that rejects  $H_0$  if  $X > 0.5$ .
- (b) Find the most powerful level  $\alpha$  test of  $H_0 : \theta = 1$  v.s.  $H_1 : \theta = 2$ .