

STAT 610: Discussion 1

1 Summary

- Order statistic: Let X_1, \dots, X_n be i.i.d sample and $X_{(1)}, \dots, X_{(n)}$ as order statistics, then

$$F_{X_{(j)}}(x) = \sum_{k=j}^n \binom{n}{k} \{F_X(x)\}^k \{1 - F_X(x)\}^{n-k}.$$

$$f_{X_{(j)}}(x) = \frac{n!}{(j-1)!(n-j)!} f_X(x) \{F_X(x)\}^{j-1} \{1 - F_X(x)\}^{n-j}.$$

$$f_{X_{(i)}, X_{(j)}}(u, v) = \frac{n!}{(i-1)!(j-1-i)!(n-j)!} f_X(u) f_X(v) \{F_X(u)\}^{i-1} \\ \times \{F_X(v) - F_X(u)\}^{j-1-i} \{1 - F_X(v)\}^{n-j} I(u < v).$$

$$f_{X_{(1)}, \dots, X_{(n)}}(x_1, \dots, x_n) = \begin{cases} n! f_X(x_1) \cdots f_X(x_n) & \text{if } -\infty < x_1 < \cdots < x_n < \infty, \\ 0 & \text{otherwise.} \end{cases}$$

- Sufficient statistics
 - Definition: A statistic $T(\mathbf{X})$ is sufficient for θ if the conditional distribution of the sample \mathbf{X} given the value of $T(\mathbf{X})$ does not depend on θ .
 - Factorization Theorem: $T(\mathbf{X})$ is sufficient for θ if and only if

$$f(\mathbf{x}|\theta) = g(T(\mathbf{x})|\theta)h(\mathbf{x}).$$

- Rao-Blackwell Theorem: Let X be a sample from a population indexed by $\theta \in \theta$ and statistic $T(\mathbf{X})$ is sufficient for θ . If $U(X)$ is a statistic used to estimate $\vartheta = \phi(\theta)$, and $E_\theta[U(X) - \vartheta]^2 < \infty$, then the statistics $h(T) = E[U(X)|T]$ satisfies

$$E_\theta[h(T) - \vartheta]^2 < E_\theta[U(X) - \vartheta]^2 \quad \theta \in \theta$$

and

$$E_\theta[h(T) - \vartheta]^2 = E_\theta[U(X) - \vartheta]^2 \quad \theta \in \theta$$

if and only if $P_\theta(U(X) = h(T)) = 1, \theta \in \theta$.

- Sufficiency make sure h is a statistics.

2 Questions

1. Let X_1, \dots, X_n be a random sample from a population with pdf

$$f_X(x) = \begin{cases} \frac{1}{\theta} & \text{if } 0 < x < \theta \\ 0 & \text{otherwise.} \end{cases}$$

Let $X_{(1)} < \dots < X_{(n)}$ be the order statistics. Show that $\frac{X_{(1)}}{X_{(n)}}$ and $X_{(n)}$ are independent.

2. Show that if T is sufficient statistic and $T = \phi(S)$, where ϕ is a real function and S is another statistic, then S is sufficient.

3. Let X_1, \dots, X_n be i.i.d. random variables from the exponential distribution $E(a, \theta)$ (i.e. $f_{a,\theta}(x) = \theta^{-1} e^{-(x-a)/\theta} I_{x>a}$). Find a two-dimensional sufficient statistics for (a, θ)

4. Let X and Y be two random variables such that Y has the binomial distribution with size N and probability π and, given $Y = y$, X has the binomial distribution with size y and probability p .

- (a) Suppose that π and N are known and $p \in (0, 1)$ is unknown. Show whether X is sufficient for p and whether Y is sufficient for p .