Hydrogen Storage Using Chemical Methods

Wenli Chen

Department of Chemical Engineering, Jiangsu University, Jiangsu, China

Abstract: Hydrogen storage is a key issue for the commercialization of electrical vehicles. Complex hydrides show energy-related functions such as chemically storing hydrogen atoms and reversible hydrogenation and dehydrogenation. In this paper, we will discuss the understanding of the mechanism of the process, using a ionic dynamics model.

Reference:

- 1. Motoaki Matsuo, Yuko Nakamori, Keitaro Yamada, and Shin-ichi Orimo. "Effects of microwave irradiation on the dehydriding reaction of the composites of lithium borohydride and microwave absorber." Applied physics letters 90, no. 23 (2007): 2907.
- 2. Motoaki Matsuo, Hitoshi Takamura, Hideki Maekawa, Hai-Wen Li, and Shin-ichi Orimo. "Stabilization of lithium superionic conduction phase and enhancement of conductivity of LiBH4 by LiCl addition." Applied Physics Letters 94, no. 8 (2009): 84103.
- 3. Motoaki Matsuo, Arndt Remhof, Pascal Martelli, Riccarda Caputo, Matthias Ernst, Yohei Miura, Toyoto Sato et al. "Complex hydrides with (BH4)— and (NH2)— anions as new lithium fast-ion conductors." Journal of the American Chemical Society 131, no. 45 (2009): 16389-16391.
- 4. Motoaki Matsuo, Toyoto Sato, Yohei Miura, Hiroyuki Oguchi, Yu Zhou, Hideki Maekawa, Hitoshi Takamura, and Shin-ichi Orimo. "Synthesis and lithium fastion conductivity of a new complex hydride Li3(NH2)2I with double-layered structure." Chemistry of Materials 22, no. 9 (2010): 2702-2704.
- 5. Yu Zhou, Motoaki Matsuo, Yohei Miura, Hitoshi Takamura, Hideki Maekawa, Arndt Remhof, Andreas Borgschulte, Andreas Züttel, Toshiya Otomo, and Shinichi Orimo. "Enhanced Electrical Conductivities of Complex Hydrides Li2(BH4)(NH2) and Li4(BH4)(NH2)3 by Melting." Materials transactions 52, no. 4 (2011): 654-657.