
CD8−like Progression: are cells later along the same lineage?
Focused comparison: B7H3_vs_41BB−L | Cell−level distributions and cached KS FDRs
L1: cytotoxic effector trajectory
FDR = 7.8e−03
Difference detected
L2: proliferating effector/dysfunction trajectory
FDR = 1.6e−03
Difference detected
L3: exhausted/dysfunctional trajectory
FDR = 0.283
No significant difference
Cumulative lineage−weighted fraction
100%
75%
50%
25%
0%
20
40
60
25
50
75
100
10
20
30
40
Shared pseudotime
mean: 26.8
mean: 27.6
mean: 37.2
mean: 29.4
mean: 16.8
mean: 18.0
CD28TM.CD28.mutζ
41BBL−CD28TMCD28.mutζ
Rightward = later shared pseudotime. Dashed lines and numbers: pooled cell−weighted means.
The KS test detects any distribution difference, not specifically 'later'.

CD8−like Progression−test overview

Cell−level test | Two mice per construct



All = global test; L1−L3 = shared lineages. FDR is adjusted within each comparison across its four tests.
L1: cytotoxic effector trajectory; L2: proliferating effector/dysfunction trajectory; L3: exhausted/dysfunctional trajectory



Each bar is one mouse's lineage−weighted mean pseudotime. Compare constructs within one panel; pseudotime scales differ between panels.