

Each cell's lineage weights sum to one; a segment is the mean weight on that lineage over all of the construct's cells.
Cells before the lineages split are divided between them. FDRs for each pair are on the next page.

CD8−like Fate−selection test overview

Cell−level test | Two mice per construct



All = global test; other columns = lineage pairs. FDR is adjusted within each comparison across its four tests.
L1: cytotoxic effector trajectory; L2: proliferating effector/dysfunction trajectory; L3: exhausted/dysfunctional trajectory

CD8−like Fate selection: how each construct's cells split between two lineages

Focused comparison: B7H3_vs_41BB−L | Cells on the first lineage only, on both (shared/early), or on the second only | Pooled cells



Shares are out of cells with weight on either lineage of the pair; cells only on the third lineage are not counted.
Relative weight w2 / (w1 + w2) < 0.25 = first lineage only, > 0.75 = second lineage only, otherwise shared. Numbers above bars are %.
Panel FDRs: pair−level MMD test for this comparison (detects a difference, not its direction).

CD8−like Fate selection: how each construct's cells split between two lineages

All CARs | Cells on the first lineage only, on both (shared/early), or on the second only | Pooled cells



Shares are out of cells with weight on either lineage of the pair; cells only on the third lineage are not counted.
Relative weight w2 / (w1 + w2) < 0.25 = first lineage only, > 0.75 = second lineage only, otherwise shared. Numbers above bars are %.
MMD FDRs for every pair are in the results TSV (adjusted within each comparison across its four tests).

CD8−like Lineage membership by mouse

Descriptive | Each bar is one mouse; n = cells



Segments are mean normalized lineage weights, not counts of cells assigned to one lineage.