CD4−like Fate selection: how each construct's cells split between two lineages

Focused comparison: B7H3_vs_41BB−L | Cells on the first lineage only, on both (shared/early), or on the second only | Pooled cells



Shares are out of cells with weight on either lineage of the pair; cells only on the third lineage are not counted.
Relative weight w2 / (w1 + w2) < 0.25 = first lineage only, > 0.75 = second lineage only, otherwise shared. Numbers above bars are %.
Panel FDRs: pair−level MMD test for this comparison (detects a difference, not its direction).

CD4−like Fate selection: how each construct's cells split between two lineages

All CARs | Cells on the first lineage only, on both (shared/early), or on the second only | Pooled cells



Shares are out of cells with weight on either lineage of the pair; cells only on the third lineage are not counted.
Relative weight w2 / (w1 + w2) < 0.25 = first lineage only, > 0.75 = second lineage only, otherwise shared. Numbers above bars are %.
MMD FDRs for every pair are in the results TSV (adjusted within each comparison across its four tests).

CD4−like Fate−selection test overview

Cell−level test | Two mice per construct



All = global test; other columns = lineage pairs. FDR is adjusted within each comparison across its four tests.
L1: regulatory/Treg−associated; L2: proliferating CD4; L3: activated helper/Th1−like


CD4−like Lineage membership by mouse
Descriptive | Each bar is one mouse; n = cells
CD28TM.CD28.mutζ
41BBL−CD28TMCD28.mutζ
CD8TM.41BB.mutζ
CD28TM.41BB.mutζ
Control CAR
n=191
n=365
n=125
n=213
n=651
n=201
n=49
n=223
n=91
n=311
100%
75%
50%
25%
0%
Mean lineage membership
D1
D2
Mouse
L1: regulatory/Treg−associated
L2: proliferating CD4
L3: activated helper/Th1−like
Segments are mean normalized lineage weights, not counts of cells assigned to one lineage.