CD8−like Fate selection: how each construct's cells split between two lineages

Focused comparison: B7H3_vs_41BB−L | Cells on the first lineage only, on both (shared/early), or on the second only | Pooled cells



Shares are out of cells with weight on either lineage of the pair; cells only on the third lineage are not counted.
Relative weight w2 / (w1 + w2) < 0.25 = first lineage only, > 0.75 = second lineage only, otherwise shared. Numbers above bars are %.
Panel FDRs: pair−level MMD test for this comparison (detects a difference, not its direction).

CD8−like Fate selection: how each construct's cells split between two lineages

All CARs | Cells on the first lineage only, on both (shared/early), or on the second only | Pooled cells



Shares are out of cells with weight on either lineage of the pair; cells only on the third lineage are not counted.
Relative weight w2 / (w1 + w2) < 0.25 = first lineage only, > 0.75 = second lineage only, otherwise shared. Numbers above bars are %.
MMD FDRs for every pair are in the results TSV (adjusted within each comparison across its four tests).

CD8−like Fate−selection test overview

Cell−level test | Two mice per construct



All = global test; other columns = lineage pairs. FDR is adjusted within each comparison across its four tests.
L1: cytotoxic effector trajectory; L2: proliferating effector/dysfunction trajectory; L3: exhausted/dysfunctional trajectory


CD8−like Lineage membership by mouse
Descriptive | Each bar is one mouse; n = cells
CD28TM.CD28.mutζ
41BBL−CD28TMCD28.mutζ
CD8TM.41BB.mutζ
CD28TM.41BB.mutζ
Control CAR
n=283
n=426
n=75
n=185
n=603
n=99
n=47
n=100
n=115
n=123
100%
75%
50%
25%
0%
Mean lineage membership
D1
D2
Mouse
L1: cytotoxic effector trajectory
L2: proliferating effector/dysfunction trajectory
L3: exhausted/dysfunctional trajectory
Segments are mean normalized lineage weights, not counts of cells assigned to one lineage.