

Review: Low-grade gliomas as neurodevelopmental disorders: insights from mouse models of neurofibromatosis-1

M. Thangarajh and D. H. Gutmann

Department of Neurology, School of Medicine, Washington University, Saint Louis, MO, USA

M. Thangarajh and D. H. Gutmann (2012) *Neuropathology and Applied Neurobiology* 38, 241–253

Low-grade gliomas as neurodevelopmental disorders: insights from mouse models of neurofibromatosis-1

Over the past few years, the traditional view of brain tumorigenesis has been revolutionized by advances in genomic medicine, molecular biology, stem cell biology and genetically engineered small-animal modelling. We now appreciate that paediatric brain tumours arise following specific genetic mutations in specialized groups of progenitor cells in concert with permissive changes in the local tumour microenvironment. This interplay between preneoplastic/neoplastic cells and non-neoplastic stromal cells is nicely illustrated by the neurofibromatosis type

1-inherited cancer syndrome, in which affected children develop low-grade astrocytic gliomas. In this review, we will use neurofibromatosis type 1 as a model system to highlight the critical role of growth control pathways, non-neoplastic cellular elements and brain region-specific properties in the development of childhood gliomas. The insights derived from examining each of these contributing factors will be instructive in the design of new therapies for gliomas in the paediatric population.

Keywords: astrocytoma, cyclic AMP, microglia, mTOR, neurofibromin, tumour microenvironment

Introduction

The most common solid tumours in children are primary brain tumours, contributing 22% of all childhood neoplasms. Of the primary brain tumours, gliomas represent the most frequent pathologic type and the leading cause of cancer-related death in children [1–3]. While the overall survival of children with low-grade glioma is good compared with other types, there are often significant cognitive and neuroendocrine adverse effects following successful adjuvant therapy [4]. In this regard, many children experience declines in their IQ and school performance, difficulties with executive function (attention

deficit), behaviour/social problems and hypothalamic/pituitary axis abnormalities [4–8].

Gliomas are a heterogeneous group of central nervous system (CNS) tumours and include astrocytomas, oligodendrogliomas and ependymomas. However, in children, the most common glioma is the WHO grade I pilocytic astrocytoma (PA) [9,10]. These tumours typically arise during the first two decades of life, without a clear sex predilection. PAs can occur sporadically or in the context of the neurofibromatosis type 1 (NF1)-inherited cancer syndrome. In either case, PAs usually affect midline structures, including the cerebellum, optic pathway and brainstem (Figure 1). On neuroimaging, PAs of the cerebellum appear as well-circumscribed and contrast-enhancing lesions, whereas those involving the optic pathway present as fusiform masses. Optic pathway gliomas (OPGs) can also exhibit prominent gadolinium contrast enhancement on magnetic resonance imaging.

Correspondence: David H. Gutmann, School of Medicine, Washington University, 660 South Euclid Avenue, Box 8111, St Louis, MO 63110, USA. Tel: +1 314 362 7379; Fax: +1 314 362 2388; E-mail: gutmannnd@neuro.wustl.edu




Figure 1. Location of sporadic and neurofibromatosis type 1 (NF1)-associated pilocytic astrocytoma (PA). Sagittal magnetic resonance images of the normal brain illustrate the most common locations for sporadic PA compared to NF1-associated PA. Whereas sporadic PAs tend to form in the cerebellum and optic pathway, NF1-associated PAs uncommonly arise in the cerebellum.

Pilocytic astrocytomas occasionally infiltrate adjacent CNS parenchyma, especially when they involve the optic pathways. Involvement of the subarachnoid space and leptomeningeal invasion are also not uncommon and do not necessarily indicate aggressive or malignant behaviour [10]. Malignant transformation is exceedingly rare, and some PAs have been reported to regress without medical intervention [11].

Histologically, PAs have a biphasic architecture with both cystic components and solid cellular areas (Figure 2A,B). Characteristically, they contain Rosenthal fibres and eosinophilic granular bodies, but typically do not harbour necrosis. Tumour cells express glial fibrillary acidic protein (GFAP), indicative of their glial (astrocytic) histogenesis (Figure 2C,D). Mitotic activity is usually low, and the growth fraction as estimated by Ki-67 immunolabelling is less than 1% [12]. While these tumours are low-grade neoplasms, they are vascular and can exhibit microvascular proliferation.

Genetic alterations in paediatric low-grade gliomas

Until recently, the most common genetic alteration associated with PAs was mutational inactivation of the *NF1* tumour suppressor gene. However, this mutation is only found in NF1-associated gliomas, and has not been reported as a causative molecular change in sporadic (non-NF1) paediatric low-grade gliomas [13–15]. The

NF1 gene product, neurofibromin, functions as a negative regulator of the p21-RAS proto-oncogene [16], such that loss of neurofibromin in NF1-associated gliomas results in increased RAS activation and cell growth [17] (Figure 3). The finding that RAS activation is observed in NF1-associated PA prompted several laboratories to determine whether mutational RAS activation accounted for sporadic PA tumorigenesis. However, this specific genetic alteration is a rare event in sporadic PA, observed in only 2–3% of cases [18,19].

Using converging genomic and gene expression strategies, recent work has culminated in the identification of signature mutations in the *BRAF* gene as a common mutation in sporadic PA [20–25]. *BRAF* is a brain RAF kinase molecule containing an amino terminal structure and a carboxyl terminal kinase domain. The amino terminal region is thought to regulate the activity of the kinase domain, such that the removal of this amino terminal domain results in unchecked *BRAF* kinase activity and increased mitogen-activated protein kinase (MAPK) mitogenic signalling. While *BRAF*-activating mutations, as seen in melanoma [26], are not typically observed in sporadic PA, genetic rearrangements result in the creation of a fusion molecule in which a portion of the *KIAA1549* gene is fused to the kinase domain of the *BRAF* gene [20–25]. Two-thirds of PAs are associated with this novel *KIAA1549:BRAF* gene fusion, which results in constitutive activation of the MAPK signalling pathway [27] (Figure 4). In addition, the presence of this




Figure 2. Neurofibromatosis type 1 (NF1)-associated pilocytic astrocytoma (PA). Histopathological analysis of a typical PA reveals both solid cellular areas and cystic regions (A) as well as areas with microvascular proliferation (B). The neoplastic glial cells are immunopositive for glial fibrillary acidic protein (GFAP) expression, a marker of astroglial lineage cells (C,D). Magnification = $\times 200$ (A–C) and $\times 600$ (D). Images were kindly provided by Dr Robert E. Schmidt, Neuropathology Division, School of Medicine, Washington University.

signature mutation is more commonly seen in cerebellar PAs [28], where it confers a threefold increase in progression-free survival [29].

Based on the discovery of the *KIAA1549:BRAF* mutational event in sporadic PAs, several reports have described other fusion rearrangements involving *BRAF*, including *FAM131B-BRAF* [30]. Similarly, oncogenic *RAF1* rearrangements have also been reported in some sporadic PAs [31].

Interestingly, *RAF* expression alone in the brains of genetically engineered mice does not result in glioma formation, unless coupled with protein kinase-B (Akt) activation or *Arf* tumour suppressor gene loss [32]. Similarly, activated *BRAF* expression alone is insufficient for gliomagenesis, but does result in glioma development when

combined with *Ink4a/Arf* loss [33]. In contrast, a recent report demonstrated that somatic retroviral delivery of mutationally activated *BRAF* into the brains of newborn mice results in the development of astrocytomas, with histological features matching the human disease [34]. Future mouse modelling will be required to reconcile these differences in experimental outcomes as well as to determine how deregulated *BRAF* function and deregulated MAPK signalling, as seen following *KIAA1549:BRAF* expression, result in glioma formation.

NF1-associated PA

NF1 is an autosomal dominant cancer predisposition syndrome affecting one in 3000 individuals worldwide. While




Figure 3. Neurofibromin is a negative regulator of cell proliferation. Neurofibromin functions as a glial cell tumour suppressor protein by accelerating the conversion of the RAS proto-oncogene from its active GTP-bound state to its inactive GDP-bound state. Loss of neurofibromin expression, as seen in neurofibromatosis type 1-associated gliomas, leads to increased RAS-GTP levels and hyperactivation of downstream Akt/mTOR signalling pathways to result in higher levels of cell proliferation. Akt, protein kinase-B; GDP, guanosine diphosphate; GRD, GTPase activating domain; GTP, guanosine triphosphate; MAPK, mitogen-activated protein kinase; MEK, mitogen-activated protein kinase kinase; mTOR, mammalian target of rapamycin.

pigmentary abnormalities (café-au-lait macules, skin-fold freckling and iris Lisch nodules) are common in affected individuals [35], 15–20% of children with NF1 develop low-grade gliomas (PAs) of the optic pathway and brain-stem (Figure 5). NF1-associated OPGs usually present at a younger age than sporadic OPGs. NF1-associated gliomas are typically detected by 4–5 years of age and rarely grow after age 10 years, whereas sporadic PAs are commonly first diagnosed in the second decade of life. In children with NF1, PAs form anywhere between the optic globe and the optic radiations, although the anterior optic pathway (optic nerves and chiasm) is most frequently involved. Infiltration of the post-chiasmal optic radiations is found in 10% of individuals with NF1-associated OPG, and is usually associated with more aggressive clinical behaviour [36].

Whereas sporadic OPG usually affects only one optic nerve, it is common to encounter bilateral optic nerve




Figure 4. BRAF fusion mutations predominate in sporadic pilocytic astrocytoma. A novel fusion of the KIAA1549 and BRAF genes results in increased BRAF kinase function. Deregulated BRAF kinase activity leads to increased mitogen-activated protein kinase (MAPK) pathway signalling, leading to higher levels of cell proliferation. KD, kinase domain; MEK, mitogen-activated protein kinase kinase.

involvement in children with NF1. NF1-associated OPGs can be large and exhibit prominent contrast enhancement on magnetic resonance imaging. However, neither the size nor degree of contrast enhancement correlates with the clinical behaviour of the tumour. In this regard, the clinical course of NF1-associated OPGs tends to be more indolent compared to their sporadic counterparts. While 15% of patients with NF1 develop OPGs [37,38], only one-third of children require treatment for clinical progression (visual loss, endocrinopathy). First-line treatment in the majority of cases is carboplatin/vincristine chemotherapy. Response rates range from 30% to 50%, and as many as 70% of children exhibit no further tumour growth while on treatment [39–42].

NF1 gene function

The *NF1* gene is a large tumour suppressor gene located on chromosome 17q11.2 and encodes a 220–250 kDa protein called neurofibromin [43–45]. Neurofibromin is expressed by numerous different cell types, including neurones, Schwann cells, oligodendrocytes, adrenal medulla, muscle and skin [46–48]. Based on its predicted protein sequence, neurofibromin was hypothesized to function primarily as a negative regulator of the p21-RAS proto-oncogene [49,50]. A small portion of the neurofibromin




Figure 5. Neurofibromatosis type 1-associated optic glioma. (A) A representative normal optic nerve (child without an optic glioma) is shown by axial magnetic resonance imaging (arrow). (B) Fusiform enlargement of the optic nerve is evident in a child with a large right optic pathway glioma (arrow). (C) The extent of the optic pathway glioma, which includes the optic chiasm, is shown on the sagittal magnetic resonance image (arrow).

protein shares sequence similarity with a family of molecules called GTPase-activating protein (GAPs) that function to inactivate RAS and reduce RAS-mediated growth signalling. Increased RAS activation leads to increased cell proliferation, and oncogenic transformation in some cell types, supporting the notion that neurofibromin negative growth regulation operates through RAS inhibition. In this regard, loss of neurofibromin in NF1-associated tumours is associated with increased levels of RAS activation [51–53].

Downstream RAS effectors relevant to neurofibromin growth regulation in neuroglial cells include Akt, mammalian target of rapamycin (mTOR) and MAPK. While BRAF regulates MAPK signalling, neurofibromin primarily regulates Akt/mTOR signalling in astrocytes *in vitro* and *in vivo* [54]. mTOR is a serine/threonine kinase involved in the phosphorylation of several downstream targets, and is important in ribosomal biogenesis [55]. The Akt/mTOR pathway is hyperactivated in *Nf1*^{−/−} astrocytes, as evidenced by increased activity (and phosphorylation) of the ribosomal S6 protein [54, 56]. In OPG mouse models and in human PA, phosphorylated S6 expression is detectable in astrocytes. *Nf1*^{−/−} astrocytes exhibit increased proliferation *in vitro*, which is blocked by rapamycin, a pharmacological inhibitor of mTOR [54]. The finding that mTOR inhibition blocks *Nf1*-deficient glial cell growth prompted preclinical genetically engineered mouse (GEM) studies aimed at determining whether rapamycin might be a suitable targeted therapy for NF1-associated glioma.

In addition to RAS regulation, neurofibromin positively controls intracellular cyclic adenosine monophosphate (cAMP) levels [57, 58]. While neurofibromin loss in NF1-associated neuroglial cells leads to increased RAS and RAS pathway activation, reduced neurofibromin expression results in attenuated intracellular cAMP levels. Consistent with this prediction, *Nf1*^{−/−} astrocytes have decreased basal levels of cAMP [58, 59]. Numerous cytokines function by binding G protein-coupled receptors and control cell growth and survival in a cAMP-dependent fashion [60]. One of these cytokines, CXCL12 (or stroma-derived factor-1- α) is highly expressed along the optic nerve in young children and mice [59]. CXCL12 normally increases cAMP levels in wild-type astrocytes to result in increased programmed cell death (apoptosis). In contrast, CXCL12 treatment of *Nf1*-deficient astrocytes leads to inappropriate cell survival [59]. This observation suggests that therapies targeting cAMP may have utility in the treatment of NF1-associated glioma.

Determinants of gliomagenesis

The unique predilection for NF1-associated gliomas to form in the optic pathway in young children raises the intriguing possibility that gliomagenesis requires more than *NF1* gene inactivation in glial progenitor cells. Several clinical observations support the requirement for other conditions for glioma development in children with NF1. First, while sporadic PAs commonly form in the cerebellum, this is an unusual location for NF1-associated PA. Second, NF1-associated PAs arise in younger children and are frequently less clinically aggressive. Third, the majority of children with NF1 do not develop gliomas, despite having a genetic susceptibility.

The underlying explanations for the pattern of gliomagenesis in NF1 are likely to reflect: (i) differences in susceptibilities of preneoplastic neuroglial cell types in the CNS; (ii) contributions from non-neoplastic cells and signals in the tumour microenvironment; and (iii) genomic modifiers that confer glioma risk.

Neuroglial populations respond differently to NF1 gene inactivation

The notion that neuroglial cells (astrocytes and astroglial progenitors) from different regions of the CNS are distinct has recently gained traction. Elegant studies by Richard Gilbertson and colleagues have demonstrated that another glial neoplasm (ependymoma) is comprised of separate tumour types based on their cell of origin and genetic background [61]. Based on these findings, it is possible that NF1-associated PAs are distinct from their sporadic counterparts, not only with respect to causative mutation (*BRAF* activation vs. *NF1* gene loss), but also cell of origin. In this regard, the molecular signature of NF1-associated PA was found to be distinct from that seen in sporadic PA [62].

The contribution of cellular heterogeneity to the pattern of gliomagenesis in NF1 is supported by recent studies demonstrating that mouse astrocytes and neural stem cells from different brain regions differ in their ability to proliferate in response to *Nf1* gene inactivation. While astrocytes from the mouse optic nerve and brainstem express neurofibromin, neocortical astrocytes have significantly lower levels of *Nf1* gene expression [63]. These differences in neurofibromin expression account for the response of these distinct astroglial cell populations to *Nf1* gene inactivation: astrocytes from the brainstem and optic

nerve increase their proliferation *in vitro* and *in vivo* following *Nf1* gene loss, while those from the neocortex fail to increase their growth [63].

In addition, neurofibromin dictates the proliferation of neural stem cells and gliogenesis in a brain region-specific manner [64]. Mouse neural stem cells from the brainstem increase their proliferation *in vitro* and their capacity to form differentiated glial cells *in vivo* following *Nf1* gene inactivation. In contrast, *Nf1*-deficient mouse neural stem cells from the cortex exhibit no significant increase in proliferation or glial differentiation relative to their wild-type counterparts. This cell autonomous difference in neural stem cell behaviour does not reflect neurofibromin expression levels, but rather the expression of one component of the mTOR complex, rictor, which specifies Akt activation.

Collectively, these results demonstrate that neuroglial cells from different brain regions respond uniquely to neurofibromin loss and that these differences may partly determine why gliomas form in the optic pathway and brainstem of children with NF1.

The contribution of the tumour microenvironment to glioma formation and growth

Studies in other cancers have revealed an important role for the tumour microenvironment in promoting tumour formation and growth [65–67]. The need for a permissive microenvironment in NF1-associated gliomagenesis is nicely illustrated by *Nf1* GEM modelling experiments. Mice lacking *Nf1* gene expression in astroglial progenitors alone do not form brain tumours [68], whereas *Nf1*+/- mice lacking *Nf1* gene expression in these same astroglial progenitors develop optic gliomas [69,70]. *Nf1*+/- mice have reduced neurofibromin expression in every cell of their body and brain, similar to children born with NF1. The requirement for reduced *Nf1* gene expression in the non-neoplastic cells of the brain supports a model in which gliomagenesis involves the productive interplay between *Nf1*+/- stromal cells and *Nf1*-/- preneoplastic/neoplastic cells (Figure 6).

Because *Nf1* loss in astroglial progenitors must be coupled with reduced *Nf1* gene expression in non-neoplastic (*Nf1*+/-) cells, *Nf1*+/- stromal cells must produce key growth/survival factors that facilitate *Nf1* OPG formation. In both human NF1-associated OPGs and *Nf1* GEM OPGs, there are numerous other cell types present, including non-neoplastic microglia, endothelial




Figure 6. Gliomagenesis requires the interplay between neoplastic and non-neoplastic cellular elements. Paediatric glioma formation ensues following cancer-causing genetic mutations in susceptible preneoplastic cells (glial cells or stem cells), leading to increased glial cell proliferation. These glial lineage cells elaborate molecules ('stromagens') that recruit non-neoplastic cells (microglia, blood vessels and reactive astrocytes) to form a microenvironment permissive for tumour evolution. The non-neoplastic cells in the tumour microenvironment in turn produce additional molecules ('gliomagens') that further promote neoplastic glial cell growth, neoplastic transformation and tumour progression.

cells and reactive astrocytes. Microglia appear early in the evolving *Nf1* mouse OPG [71], which suggests that they may be important stromal cells relevant to glioma formation and growth. These immune system-like cells comprise as many as 30% of the cells in human NF1-associated PAs [72] and produce numerous growth factors and cytokines [73,74].

The importance of microglia to NF1-associated optic glioma growth is underscored by several experimental observations. First, *Nf1*+/-, but not wild-type, microglia increase the proliferation of *Nf1*-/- astrocytes *in vitro* [75]. Second, soluble factors made by *Nf1*+/- microglia increase the proliferation of *Nf1*-deficient astrocytes *in vitro* [75]. Third, inhibition of microglia function attenuates *Nf1* mouse OPG proliferation *in vivo*. In these experiments, blocking microglial activation with the antibiotic minocycline reduces *Nf1* OPG growth [75]. Similarly, inhibiting the signalling pathway responsible for *Nf1*+/- microglial function results in decreased *Nf1* OPG proliferation *in vivo* [76]. Finally, genetic ablation of microglia leads to reduced optic nerve glioma proliferation *in vivo* [72]. Collectively, these observations establish microglia

as one of the key stromal cell types that drives glioma growth.

Based on *in vitro* experiments demonstrating that *Nf1*^{+/-} microglia produce soluble factors that increase *Nf1*-deficient astrocyte growth, we sought to identify these glioma-promoting molecules (gliomagens). Two such gliomagens have been studied, meningioma-expressed antigen-5 (MGEA5) and CXCL12. *Nf1*^{+/-} microglia express higher levels of MGEA5, a member of the hyaluronidase family of enzymes, than their wild-type counterparts. While the role of MGEA5 and related hyaluronidase molecules in NF1-associated gliomagenesis has not been fully elucidated, hyaluronidase treatment of *Nf1*-deficient astrocytes increases their proliferation, and inhibition of hyaluronidase function ameliorates the growth advantage conferred by *Nf1*^{+/-} microglia [75]. Additional studies will be required to clarify the role of this interesting molecule as a gliomagen in the formation and growth of NF1-associated gliomas.

In addition, *Nf1*^{+/-} microglia produce two to threefold more CXCL12 than wild-type microglia. The importance of CXCL12 to gliomagenesis is underscored by several observations. First, CXCL12 increases the survival of *Nf1*^{-/-} astrocytes in a cAMP-dependent fashion, but leads to increased apoptosis in wild-type astrocytes [59]. Second, pharmacological inhibition of cAMP degradation using a phosphodiesterase-4 (PDE4) inhibitor, Rolipram, attenuates *Nf1* mouse OPG growth *in vivo* [77]. Third, lentiviral expression of CXCL12 in the forebrain of *Nf1* OPG mice results in low-penetrance glioma formation [78]. To more effectively lower cyclic AMP levels and mimic the effect of CXCL12 on *Nf1*-deficient astroglial cells, lentiviral delivery of PDE4 into the forebrains of *Nf1* optic glioma mice results in a greater incidence of glioma formation [77].

Together, these findings suggest that gliomagenesis and continued tumour growth require interactions between non-neoplastic and neoplastic cell types. We envision a model in which *Nf1*-deficient preneoplastic glial progenitor cells produce molecules that attract microglia and other non-neoplastic cell types (stromagens). These stromagens create a microenvironment that supports glioma formation. In turn, the attracted *Nf1*^{+/-} stromal cells (microglia) elaborate gliomagens (growth factors and cytokines) that facilitate further *Nf1*-deficient astroglial cell proliferation/survival and permit neoplastic transformation. Each of these factors represents a potential target for stroma-directed therapies that aim to inhibit the

formation of a permissive tumour microenvironment or block the tumour-promoting activity of stroma-derived gliomagens.

Genomic determinants of gliomagenesis

As 80–85% of children with NF1 do not develop OPGs while the vast majority (90–95%) of *Nf1* genetically engineered mice harbouring identical genetic changes do, it is likely that subtle genomic alterations dictate which children with NF1 will develop a brain tumour. In experimental *Nf1* mouse models, the specific genetic background greatly influences glioma development. Mice heterozygous for mutations in the *Nf1* and *Trp53* (p53) tumour suppressor genes (*Nf1*^{+/-}; *Trp53*^{+/- cis}; NP^{cis} mice) develop brain gliomas at a much higher frequency on the B6/C57BL/6J (B6) genetic background compared to the 129S4/SvJae (129S4) background [79–81]. These findings support the existence of epigenetic or polymorphic differences that potentially confer resistance or susceptibility to gliomagenesis. These modifier loci could function by changing: (1) the expression or function of genes important for glial progenitor behaviour; (2) neurofibromin growth regulation; or (3) chemokine function or microglia activity, which separately or collectively impact on glioma formation or growth. While the modifier genes have yet to be elucidated, genomic polymorphisms in these loci might identify which children with NF1 are at highest risk for the development of OPG. The identification of single nucleotide polymorphisms predictive of glioma risk would positively impact on the management of children with NF1, and would allow clinicians to stratify children from an early age into clinically relevant subgroups for surveillance and potential treatments.

Implications for paediatric glioma-targeted therapies

The management approaches for PA in children currently involve surgery and genotoxic chemotherapy. Surgical resection is often a first-line therapy for cerebellar and hemispheric PAs, but is associated with significant morbidity when tumours are located in deeper structures (brainstem) or in the optic pathway. For these tumours, chemotherapy or radiotherapy is recommended. While radiotherapy is curative, its use for paediatric glioma is problematic, owing to the high incidence of cognitive and neuroendocrine adverse effects that result from

irradiating the developing brain. Moreover, radiation is contraindicated in children with NF1; following brain irradiation, there is an increased frequency of secondary high-grade glioma and moyamoya disease [82,83].

Chemotherapy with carboplatin and vincristine is usually well tolerated and achieves good disease control [40,42,84]. However, antineoplastic drugs cause DNA damage not only in cancer cells, but may also impact other dividing cell populations in the developing brain. For this reason, more targeted therapies are actively being sought for this patient population.

With the availability of robust preclinical mouse models of *Nf1* OPG, it becomes possible to identify and evaluate alternative therapeutic strategies. Using *Nf1* OPG mice, previous studies have shown that treatment with a genotoxic alkylating agent, temozolomide, results in decreased OPG volumes, reduced glioma proliferation and increased glioma cell death [85]. These GEM strains have subsequently been used to evaluate both neoplastic and non-neoplastic cell-directed therapies. In this regard, treatments that target the deregulated RAS/mTOR [85] and cyclic AMP [77] signalling pathways in the *Nf1*-deficient astroglial cells reduce *Nf1* OPG proliferation. Similarly, therapies that inhibit non-neoplastic microglia function also attenuate OPG growth in these *Nf1*-mutant mice *in vivo* [72,75,76].

Another use of *Nf1* genetically engineered mice is the development of therapies that restore visual function. Previous studies have revealed that improvement in visual function infrequently accompanies the successful chemotherapeutic treatment of NF1-associated glioma [41]. Insights into the cellular and molecular basis for optic nerve dysfunction have been partly revealed by the analysis of *Nf1* OPG mice. In these studies, reduced visual evoked potentials are seen early during tumour development and are associated with axonal damage and retinal ganglion cell death [86,87]. *Nf1*+/- neurones exhibit increased apoptosis, which reflects reduced levels of intracellular cAMP [88]. Raising cAMP levels with the PDE4 inhibitor (Rolipram) during tumour formation resulted in reduced retinal ganglion cell apoptosis [88], suggesting a potential neuroprotective adjuvant strategy for managing OPGs in children with NF1.

Conclusion

Advances in genomic technologies, small-animal modelling and molecular biology discovery methods offer

unprecedented opportunities to develop personalized treatments for children with low-grade gliomas. It is possible that future therapies will incorporate genetic/genomic diagnostics to identify patients at high risk for glioma development and progression, as well as specific glioma subtypes most likely to respond to targeted therapies. Similarly, the use of robust low-grade glioma GEM strains facilitates the identification of treatments with efficacy against particular molecular subgroups of paediatric low-grade glioma. These next-generation treatments might target specific cell types or gliomagens in the tumour microenvironment in combination with standard or biologically based antineoplastic therapies. This ecosystems approach could reduce the toxicity of current treatments on the developing brains of children. Finally, the use of adjuvant neuroprotective strategies has the potential to decrease secondary neurocognitive declines and partially restore vision in affected children.

Acknowledgements

We thank Dr Robert Schmidt (Division of Neuropathology, School of Medicine, Washington University) for the photomicrographs shown in Figure 2.

Conflict of interest

The authors have no conflicts of interest to disclose.

References

- 1 Surawicz TS, McCarthy BJ, Kupelian V, Jukich PJ, Bruner JM, Davis FG. Descriptive epidemiology of primary brain and CNS tumors: results from the Central Brain Tumor Registry of the United States, 1990–1994. *Neuro-Oncol* 1999; **1**: 14–25
- 2 Jukich PJ, McCarthy BJ, Surawicz TS, Freels S, Davis FG. Trends in incidence of primary brain tumors in the United States, 1985–1994. *Neuro-Oncol* 2001; **3**: 141–51
- 3 Surawicz TS, Davis F, Freels S, Laws ER Jr, Menck HR. Brain tumor survival: results from the National Cancer Data Base. *J Neurooncol* 1998; **40**: 151–60
- 4 Armstrong GT, Conklin HM, Huang S, Srivastava D, Sanford R, Ellison DW, Merchant TE, Hudson MM, Hoehn ME, Robison LL, Gajjar A, Morris EB. Survival and long-term health and cognitive outcomes after low-grade glioma. *Neuro-Oncol* 2011; **13**: 223–34
- 5 Mulhern RK, Reddick WE, Palmer SL, Glass JO, Elkin TD, Kun LE, Taylor J, Langston J, Gajjar A. Neurocognitive

- deficits in medulloblastoma survivors and white matter loss. *Ann Neurol* 1999; **46**: 834–41
- 6 Palmer SL, Goloubeva O, Reddick WE, Glass JO, Gajjar A, Kun L, Merchant TE, Mulhern RK. Patterns of intellectual development among survivors of pediatric medulloblastoma: a longitudinal analysis. *J Clin Oncol* 2001; **19**: 2302–8
 - 7 Briere ME, Scott JG, McNall-Knapp RY, Adams RL. Cognitive outcome in pediatric brain tumor survivors: delayed attention deficit at long-term follow-up. *Pediatr Blood Cancer* 2008; **50**: 337–40
 - 8 Sands SA, Oberg JA, Gardner SL, Whiteley JA, Glade-Bender JL, Finlay JL. Neuropsychological functioning of children treated with intensive chemotherapy followed by myeloablative consolidation chemotherapy and autologous hematopoietic cell rescue for newly diagnosed CNS tumors: an analysis of the Head Start II survivors. *Pediatr Blood Cancer* 2010; **54**: 429–36
 - 9 Sievert AJ, Fisher MJ. Pediatric low-grade gliomas. *J Child Neurol* 2009; **24**: 1397–408
 - 10 Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, Burger PC, Jouvet A, Scheithauer BW, Kleihues P. The 2007 WHO classification of tumours of the central nervous system. *Acta Neuropathol* 2007; **114**: 97–109
 - 11 Parsa CF, Hoyt CS, Lesser RL, Weinstein JM, Strother CM, Muci-Mendoza R, Ramella M, Manor RS, Fletcher WA, Repka MX, Garrity JA, Ebner RN, Monteiro ML, McFadzean RM, Rubtsova IV, Hoyt WF. Spontaneous regression of optic gliomas: thirteen cases documented by serial neuroimaging. *Arch Ophthalmol* 2001; **119**: 516–29
 - 12 Giannini C, Scheithauer BW, Burger PC, Christensen MR, Wollan PC, Sebo TJ, Forsyth PA, Hayostek CJ. Cellular proliferation in pilocytic and diffuse astrocytomas. *J Neuropathol Exp Neurol* 1999; **58**: 46–53
 - 13 Gutmann DH, Donahoe J, Brown T, James CD, Perry A. Loss of neurofibromatosis 1 (NF1) gene expression in NF1-associated pilocytic astrocytomas. *Neuropathol Appl Neurobiol* 2000; **26**: 361–7
 - 14 Kluwe L, Hagel C, Tatagiba M, Thomas S, Stavrou D, Oster-tag H, von Deimling A, Mautner VF. Loss of NF1 alleles distinguish sporadic from NF1-associated pilocytic astrocytomas. *J Neuropathol Exp Neurol* 2001; **60**: 917–20
 - 15 Wimmer K, Eckart M, Meyer-Puttlitz B, Fonatsch C, Pietsch T. Mutational and expression analysis of the NF1 gene argues against a role as tumor suppressor in sporadic pilocytic astrocytomas. *J Neuropathol Exp Neurol* 2002; **61**: 896–902
 - 16 Martin GA, Viskochil D, Bollag G, McCabe PC, Crosier WJ, Haubruck H, Conroy L, Clark R, O'Connell P, Cawthon RM, Innis MA, McCormick F. The GAP-related domain of the neurofibromatosis type 1 gene product interacts with ras p21. *Cell* 1990; **63**: 843–9
 - 17 Lau N, Feldkamp MM, Roncari L, Loehr AH, Shannon P, Gutmann DH, Guha A. Loss of neurofibromin is associated with activation of RAS/MAPK and PI3-K/AKT signaling in a neurofibromatosis 1 astrocytoma. *J Neuropathol Exp Neurol* 2000; **59**: 759–67
 - 18 Sharma MK, Zehnbauser BA, Watson MA, Gutmann DH. RAS pathway activation and an oncogenic RAS mutation in sporadic pilocytic astrocytoma. *Neurology* 2005; **65**: 1335–6
 - 19 Janzarik WG, Kratz CP, Loges NT, Olbrich H, Klein C, Schafer T, Scheurlen W, Roggendorf W, Weiller C, Niemeyer C, Korinthenberg R, Pfister S, Omran H. Further evidence for a somatic KRAS mutation in a pilocytic astrocytoma. *Neuropediatrics* 2007; **38**: 61–3
 - 20 Pfister S, Janzarik WG, Remke M, Ernst A, Werft W, Becker N, Toedt G, Wittman A, Kratz C, Olbrich H, Amadi R, Thieme B, Joss S, Radlwinner B, Kulozik A, Pietsch T, Herold-Mende C, Gnekow A, Reifenberger G, Korshunov A, Scheurlen W, Omran H, Lichter P. BRAF gene duplication constitutes a mechanism of MAPK pathway activation in low-grade astrocytomas. *J Clin Invest* 2008; **118**: 1739–49
 - 21 Bar EE, Lin A, Tihan T, Burger PC, Eberhart CG. Frequent gains at chromosome 7q34 involving BRAF in pilocytic astrocytoma. *J Neuropathol Exp Neurol* 2008; **67**: 878–87
 - 22 Jones DT, Kocialkowski S, Liu L, Pearson DM, Backlund LM, Ichimura K, Collins VP. Tandem duplication producing a novel oncogenic BRAF fusion gene defines the majority of pilocytic astrocytomas. *Cancer Res* 2008; **68**: 8673–7
 - 23 Sievert AJ, Jackson EM, Gai X, Hakonarson H, Judkins AR, Resnick AC, Sutton LN, Storm PB, Shikh TH, Biegel JA. Duplication of 7q34 in pediatric low-grade astrocytomas detected by high-density single-nucleotide polymorphism-based genotype arrays results in a novel BRAF fusion gene. *Brain Pathol* 2009; **19**: 449–58
 - 24 Wan PT, Garnett MJ, Roe SM, Lee S, Niculescu-Duvaz D, Good VM, Jones CM, Marshall CJ, Springer CJ, Barford D, Marais R, Cancer Genome Project. Mechanism of activation of the RAF-ERK signaling pathway by oncogenic mutations of B-RAF. *Cell* 2004; **116**: 855–67
 - 25 Yu J, Deshmukh H, Gutmann RJ, Emnett RJ, Rodriguez FJ, Watson MA, Nagarajan R, Gutmann DH. Alterations of BRAF and HIPK2 loci predominate in sporadic pilocytic astrocytoma. *Neurology* 2009; **73**: 1526–31
 - 26 Davies H, Bignell GR, Cox C, Stephens P, Edkins S, Clegg S, Teague J, Woffendin H, Garnett MJ, Bottomley W, Davis N, Dicks E, Ewing R, Floyd Y, Gray K, Hall S, Hawes R, Hughes J, Mosmidou V, Menzies A, Mould C, Parker A, Stevens C, Watt S, Hooper S, Wilson R, Jayatilake H, Gusterson BA, Cooper C, Shipley J, Hargrave D, Pritchard-Jones K, Maitland N, Chenevix-Trench G, Riggins GJ, Bigner DD, Palmieri G, Cossu A, Flanagan H, Nicholson A, Ho JW, Leung SY, Yuen ST, Weber BL, Siegler HE, D'Arrow TL, Paterson H, Marais R, Marshall CJ, Wooster R, Stratton MR, Futreal PA. Mutations of the BRAF gene in human cancer. *Nature* 2002; **417**: 949–54

- 27 Forshe T, Tatevossian RG, Lawson AR, Ma J, Neale G, Ogunkolade BW, Jones TA, Aarum J, Dalton J, Bailey S, Chaplin T, Carter RL, Gajjar A, Broniscer A, Young BD, Ellison DW, Sheer D. Activation of the ERK/MAPK pathway: a signature genetic defect in posterior fossa pilocytic astrocytomas. *J Pathol* 2009; **218**: 172–81
- 28 Jacob K, Albrecht S, Sollier C, Faury D, Sader E, Montpetit A, Serre D, Hauser P, Garami M, Bognar L, Hanzely Z, Montes JL, Atkinson J, Farmer JP, Bouffet E, Hawkins C, Tabori U, Jabado N. Duplication of 7q34 is specific to juvenile pilocytic astrocytomas and a hallmark of cerebellar and optic pathway tumours. *Br J Cancer* 2009; **101**: 722–33
- 29 Hawkins C, Walker E, Mohamed N, Zhang C, Jacob K, Shirinian M, Alon N, Kahn D, Fried I, Scheinemann K, Tsangaris E, Dirks P, Tressler R, Bouffet E, Jabado N, Tabori U. BRAF-KIAA1549 Fusion Predicts Better Clinical Outcome in Pediatric Low-Grade Astrocytoma. *Clin Cancer Res* 2011; **17**: 4790–8
- 30 Cin H, Meyer C, Herr R, Janzarik WG, Lambert S, Jones DT, Jacob K, Benner A, Witt H, Remke M, Bender S, Falkenstein F, Can Ahn TN, Olbrich H, von Deimling A, Pekrun A, Kulozik AE, Gnekow A, Scheurlen W, Witt O, Omran H, Jabado N, Collins VP, Brummer T, Marschalek R, Lichter P, Korshunov A, Pfister SM. Oncogenic FAM131B-BRAF fusion resulting from 7q34 deletion comprises an alternative mechanism of MAPK pathway activation in pilocytic astrocytoma. *Acta Neuropathol* 2011; **121**: 763–74
- 31 Jones DT, Kocialkowski S, Liu L, Pearson DM, Ichimura K, Collins VP. Oncogenic RAF1 rearrangement and a novel BRAF mutation as alternatives to KIAA1549:BRAF fusion in activating the MAPK pathway in pilocytic astrocytoma. *Oncogene* 2009; **28**: 2119–23
- 32 Lyustikman Y, Momota H, Pao W, Holland EC. Constitutive activation of Raf-1 induces glioma formation in mice. *Neoplasia* 2008; **10**: 501–10
- 33 Robinson JP, VanBrocklin MW, Guilbeault AR, Signorelli DL, Brandner S, Holmen SL. Activated BRAF induces gliomas in mice when combined with Ink4a/Arf loss or Akt activation. *Oncogene* 2010; **29**: 335–44
- 34 Gronych J, Korshunov A, Bageritz J, Milde T, Jugold M, Hambardzumyan D, Remke M, Hartmann C, Witt H, Jones DT, Witt O, Heiland S, Bendszus M, Holland EC, Pfister S, Lichter P. An activated mutant BRAF kinase domain is sufficient to induce pilocytic astrocytoma in mice. *J Clin Invest* 2011; **121**: 1344–48
- 35 Gutmann DH, Aylsworth A, Carey JC, Korf B, Marks J, Pyeritz RE, Rubenstein A, Viskochil D. The diagnostic evaluation and multidisciplinary management of neurofibromatosis 1 and neurofibromatosis 2. *JAMA* 1997; **278**: 51–7
- 36 Balcer LJ, Liu GT, Heller G, Bilaniuk L, Volpe NJ, Galetta SL, Molloy PT, Phillips PC, Janss AJ, Vaughn S, Maguire MG. Visual loss in children with neurofibromatosis type 1 and optic pathway gliomas: relation to tumor location by magnetic resonance imaging. *Am J Ophthalmol* 2001; **131**: 442–5
- 37 Listernick R, Charrow J, Greenwald M, Mets M. Natural history of optic pathway tumors in children with neurofibromatosis type 1: a longitudinal study. *J Pediatr* 1994; **125**: 63–6
- 38 Listernick R, Louis DN, Packer RJ, Gutmann DH. Optic pathway gliomas in children with neurofibromatosis 1: consensus statement from the NF1 Optic Glioma Task Force. *Ann Neurol* 1997; **41**: 143–49
- 39 Listernick R, Ferner RE, Liu GT, Gutmann DH. Optic pathway gliomas in neurofibromatosis-1: controversies and recommendations. *Ann Neurol* 2007; **61**: 189–98
- 40 Packer RJ, Lange B, Ater J, Nicholson HS, Allen J, Walker R, Prados M, Jakacki R, Reaman G, Needles MN, Phillips PC, Ryan J, Boyett JM, Geyer R, Finlay J. Carboplatin and vincristine for recurrent and newly diagnosed low-grade gliomas of childhood. *J Clin Oncol* 1993; **11**: 850–6
- 41 Dalla Via P, Opocher E, Pinello ML, Calderone M, Viscardi E, Clementi M, Battistella PA, Laverda AM, Da Dalt L, Perilongo G. Visual outcome of a cohort of children with neurofibromatosis type 1 and optic pathway glioma followed by a pediatric neuro-oncology program. *Neuro-Oncol* 2007; **9**: 430–7
- 42 Mahoney DH Jr, Cohen ME, Friedman HS, Kepner JL, Gerner L, Langston JW, James HE, Duffner PK, Kun LE. Carboplatin is effective therapy for young children with progressive optic pathway tumors: a Pediatric Oncology Group phase II study. *Neuro-Oncol* 2000; **2**: 213–20
- 43 Wallace MR, Marchuk DA, Andersen LB, Letcher R, Odeh HM, Saulino AM, Fountain JW, Brereton A, Nicholson J, Mitchell AL, Collins FS. Type 1 neurofibromatosis 1 gene: identification of a large transcript disrupted in three NF1 patients. *Science* 1990; **249**: 181–86
- 44 Viskochil D, Buchberg AM, Xu G, Cawthon RM, Stevens J, Wolff RK, Culver M, Carey JC, Copeland NG, Jenkins NA, White R. Deletions and a translocation interrupt a cloned gene at the neurofibromatosis type 1 locus. *Cell* 1990; **62**: 187–92
- 45 Xu GF, O'Connell P, Viskochil D, Cawthon R, Robertson M, Culver M, Stevens J, Robertson M, Dunn D, Gesteland R, O'Connell, White R. The neurofibromatosis type 1 gene encodes a protein related to GAP. *Cell* 1990; **62**: 599–608
- 46 Daston MM, Scrabble H, Nordlund M, Sturbaum AK, Nissen LM, Ratner N. The protein product of the neurofibromatosis type 1 gene is expressed at highest abundance in neurons, Schwann cells, and oligodendrocytes. *Neuron* 1992; **8**: 415–28
- 47 Gutmann DH, Wood DL, Collins FS. Identification of the neurofibromatosis type 1 gene product. *Proc Natl Acad Sci U S A* 1991; **88**: 9658–62
- 48 DeClue JE, Cohen BD, Lowy DR. Identification and characterization of the neurofibromatosis type 1 gene product. *Proc Natl Acad Sci U S A* 1991; **88**: 9914–18
- 49 Xu GF, Lin B, Tanaka K, Dunn D, Wood D, Gesteland R, White R, Weiss R, Tamanoi F. The catalytic domain of the

- neurofibromatosis type 1 gene product stimulates ras GTPase and complements ira mutants of *S. cerevisiae*. *Cell* 1990; **63**: 835–41
- 50 Ballester R, Marchuk D, Boguski M, Saulino A, Letcher R, Wigler M, Collins FS. The NF1 locus encodes a protein functionally related to mammalian GAP and yeast IRA proteins. *Cell* 1990; **63**: 851–9
 - 51 Basu TN, Gutmann DH, Fletcher JA, Glover TW, Collins FS, Downward J. Aberrant regulation of ras proteins in malignant tumour cells from type 1 neurofibromatosis patients. *Nature* 1992; **356**: 713–15
 - 52 Bollag G, Clapp DW, Shih S, Adler F, Zhang YY, Thompson P, Lange BJ, Freedman MH, McCormick F, Jacks T, Shannon K. Loss of NF1 results in activation of the Ras signaling pathway and leads to aberrant growth in haematopoietic cells. *Nat Genet* 1996; **12**: 144–8
 - 53 DeClue JE, Papageorge AG, Fletcher JA, Diehl SR, Ratner N, Vass WC, Lowy DR. Abnormal regulation of mammalian p21ras contributes to malignant tumor growth in von Recklinghausen (type 1) neurofibromatosis. *Cell* 1992; **69**: 265–73
 - 54 Dasgupta B, Yi Y, Chen DY, Weber JD, Gutmann DH. Proteomic analysis reveals hyperactivation of the mammalian target of rapamycin pathway in neurofibromatosis 1-associated human and mouse brain tumors. *Cancer Res* 2005; **65**: 2755–60
 - 55 Guertin DA, Sabatini DM. Defining the role of mTOR in cancer. *Cancer Cell* 2007; **12**: 9–12
 - 56 Sandsmark DK, Zhang H, Hegedus B, Pelletier CL, Weber JD, Gutmann DH. Nucleophosmin mediates mammalian target of rapamycin-dependent actin cytoskeleton dynamics and proliferation in neurofibromin-deficient astrocytes. *Cancer Res* 2007; **67**: 4790–9
 - 57 Tong J, Hannan F, Zhu Y, Bernards A, Zhong Y. Neurofibromin regulates G protein-stimulated adenylyl cyclase activity. *Nat Neurosci* 2002; **5**: 95–6
 - 58 Dasgupta B, Dugan LL, Gutmann DH. The neurofibromatosis 1 gene product neurofibromin regulates pituitary adenylyl cyclase-activating polypeptide-mediated signaling in astrocytes. *J Neurosci* 2003; **23**: 8949–54
 - 59 Warrington NM, Woerner BM, Dagainakatte GC, Dasgupta B, Perry A, Gutmann DH, Rubin JB. Spatiotemporal differences in CXCL12 expression and cyclic AMP underlie the unique pattern of optic glioma growth in neurofibromatosis type 1. *Cancer Res* 2007; **67**: 8588–95
 - 60 Miller RJ, Rostene W, Apartis E, Banisadr G, Biber K, Milligan ED, White FA, Zhang J. Chemokine action in the nervous system. *J Neurosci* 2008; **28**: 11792–95
 - 61 Johnson RA, Wright KD, Poppleton H, Mohankumar KM, Finkelstein D, Pounds SB, Rand V, Leary SE, White E, Eden C, Hogg T, Northcott P, Mack S, Neale G, Wang YD, Coyle B, Atkinson J, DeWire M, Kranenburg TA, Gillespie Y, Allen JC, Merchant T, Boop FA, Sanford RA, Gajjar A, Ellison DW, Taylor MD, Grundy RG, Gilbertson RJ. Cross-species genomics matches drive mutations and cell compartments to model ependymoma. *Nature* 2010; **466**: 632–36
 - 62 Sharma MK, Mansur DB, Reifengerger G, Perry A, Leonard JR, Aldape KD, Albin MG, Emnett RJ, Loeser S, Watson MA, Nagarajan R, Gutmann DH. Distinct genetic signatures among pilocytic astrocytomas relate to their brain region origin. *Cancer Res* 2007; **67**: 890–900
 - 63 Yeh TH, Lee DY, Gianino SM, Gutmann DH. Microarray analyses reveal regional astrocyte heterogeneity with implications for neurofibromatosis type 1 (NF1)-regulated glial proliferation. *Glia* 2009; **57**: 1239–49
 - 64 Lee DY, Yeh TH, Emnett RJ, White CR, Gutmann DH. Neurofibromatosis-1 regulates neuroglial progenitor proliferation and glial differentiation in a brain region-specific manner. *Genes Dev* 2010; **24**: 2317–29
 - 65 Bhowmick NA, Chytil A, Plith D, Gorska AE, Dumont N, Shappell S, Washington MK, Neilson EG, Moses HL. TGF-beta signaling in fibroblasts modulates the oncogenic potential of adjacent epithelia. *Science* 2004; **303**: 848–51
 - 66 Kenny HA, Kaur S, Coussens LM, Lengyel E. The initial steps of ovarian cancer metastasis are mediated by MMP-2 cleavage of vitronectin and fibronectin. *J Clin Invest* 2008; **118**: 1367–79
 - 67 Kessenbrock K, Plaks V, Werb Z. Matrix metalloproteases: regulators of the tumor microenvironment. *Cell* 2010; **141**: 52–67
 - 68 Bajenaru ML, Zhu Y, Hedrick NM, Donahoe J, Parada LF, Gutmann DH. Astrocyte-specific inactivation of the neurofibromatosis 1 gene (NF1) is insufficient for astrocytoma formation. *Mol Cell Biol* 2002; **22**: 5100–13
 - 69 Bajenaru ML, Hernandez MR, Perry A, Zhu Y, Parada LF, Garbow JR, Gutmann DH. Optic nerve glioma in mice requires astrocyte Nf1 gene inactivation and Nf1 brain heterozygosity. *Cancer Res* 2003; **63**: 8573–7
 - 70 Zhu Y, Harada T, Liu L, Lush ME, Guignard F, Harada C, Burns DK, Bajenaru ML, Gutmann DH, Parada LF. Inactivation of NF1 in CNS causes increased glial progenitor proliferation and optic glioma formation. *Development* 2005; **132**: 5577–88
 - 71 Bajenaru ML, Garbow JR, Perry A, Hernandez MR, Gutmann DH. Natural history of neurofibromatosis 1-associated optic nerve glioma in mice. *Ann Neurol* 2005; **57**: 119–27
 - 72 Simmons GW, Pong WW, Emnett RJ, White CR, Gianino SM, Rodriguez FJ, Gutmann DH. Neurofibromatosis-1 heterozygosity increases microglia in a spatially and temporally restricted pattern relevant to mouse optic glioma formation and growth. *J Neuropathol Exp Neurol* 2011; **70**: 51–62
 - 73 Graeber MB, Scheithauer BW, Kreutzberg GW. Microglia in brain tumors. *Glia* 2002; **40**: 252–59
 - 74 Hanisch UK, Kettenmann H. Microglia: active sensor and versatile effector cells in the normal and pathologic brain. *Nat Neurosci* 2007; **10**: 1387–94

- 75 Dagniakatte GC, Gutmann DH. Neurofibromatosis-1 (Nf1) heterozygous brain microglia elaborate paracrine factors that promote Nf1-deficient astrocyte and glioma growth. *Hum Mol Genet* 2007; **16**: 1098–112
- 76 Dagniakatte GC, Gianino SM, Zhao NW, Parsadanian AS, Gutmann DH. Increased c-Jun-NH2-kinase signaling in neurofibromatosis-1 heterozygous microglia drives microglia activation and promotes optic glioma proliferation. *Cancer Res* 2008; **68**: 10358–66
- 77 Warrington NM, Gianino SM, Jackson E, Goldhoff P, Garbow JR, Piwnica-Worms D, Gutmann DH, Rubin JB. Cyclic AMP suppression is sufficient to induce gliomagenesis in a mouse model of neurofibromatosis-1. *Cancer* 2010; **70**: 5717–27
- 78 Sun T, Gianino SM, Jackson E, Piwnica-Worms D, Gutmann DH, Rubin JB. CXCL12 alone is insufficient for gliomagenesis in Nf1 mutant mice. *J Neuroimmunol* 2010; **224**: 108–13
- 79 Reilly KM, Tuskan RG, Christy E, Loisel DA, Ledger J, Bronson RT, Smith CD, Tsang S, Munroe DJ, Jacks T. Susceptibility to astrocytoma in mice mutant for Nf1 and Trp53 is linked to chromosome 11 and subject to epigenetic effects. *Proc Natl Acad Sci U S A* 2004; **101**: 13008–13
- 80 Reilly KM, Loisel DA, Bronson RT, McLaughlin ME, Jacks T. Nf1:Trp53 mutant mice develop glioblastoma with evidence of strain-specific effects. *Nat Genet* 2000; **26**: 109–13
- 81 Hawes JJ, Tuskan RG, Reilly KM. Nf1 expression is dependent on strain background: implications for tumor suppressor haploinsufficiency studies. *Neurogenetics* 2007; **8**: 121–30
- 82 Sharif S, Ferner R, Birch JM, Gillespie JE, Gattamaneni HR, Baser ME, Evans DG. Second primary tumors in neurofibromatosis 1 patients treated for optic glioma: substantial risks after radiotherapy. *J Clin Oncol* 2006; **24**: 2570–5
- 83 Grill J, Couanet D, Cappelli C, Habrand JL, Rodriguez D, Sainte-Rose C, Kalifa C. Radiation-induced cerebral vasculopathy in children with neurofibromatosis and optic pathway glioma. *Ann Neurol* 1999; **45**: 393–6
- 84 Gnekow AK, Kortmann RD, Pietsch T, Emser A. Low grade chiasmatic-hypothalamic glioma-carboplatin and vincristin chemotherapy effectively defers radiotherapy within a comprehensive treatment strategy – report from the multicenter treatment study for children and adolescents with a low grade glioma – HIT-LGG 1996 – of the Society of Pediatric Oncology and Hematology (GPOH). *Klin Padiatr* 2004; **216**: 331–42
- 85 Hegedus B, Banerjee D, Yeh TH, Rothermich S, Perry A, Rubin JB, Gutmann DH. Preclinical cancer therapy in a mouse model of neurofibromatosis-1 optic glioma. *Cancer Res* 2008; **68**: 1520–8
- 86 Hegedus B, Hughes FW, Garbow JR, Gianino S, Banerjee D, Kim K, Ellisman ME, Gutmann DH. Optic nerve dysfunction in a mouse model of neurofibromatosis-1 optic glioma. *J Neuropathol Exp Neurol* 2009; **68**: 542–51
- 87 Kim KY, Ju WK, Hegedus B, Gutmann DH, Ellisman MH. Ultrastructural characterization of the optic pathway in a mouse model of neurofibromatosis-1 optic glioma. *Neuroscience* 2010; **170**: 178–88
- 88 Brown JA, Gianino SM, Gutmann DH. Defective cAMP generation underlies the sensitivity of CNS neurons to neurofibromatosis-1 heterozygosity. *J Neurosci* 2010; **30**: 5579–89

Received 22 August 2011

Accepted after revision 9 October 2011

Published online Article Accepted on 28 October 2011