

Molecular Therapies for Tuberous Sclerosis and Neurofibromatosis

David Neal Franz · Brian D. Weiss

Published online: 29 April 2012
© Springer Science+Business Media, LLC 2012

Abstract Neurofibromatosis type 1 (NF1) and tuberous sclerosis complex (TSC) are autosomal-dominant genetic disorders that result from dysregulation of the PI3K/AKT/mammalian target of rapamycin (mTOR) pathway. NF1 is caused by mutations in the NF1 gene on chromosome 17q11.2. Its protein product, neurofibromin, functions as a tumor suppressor and ultimately produces constitutive upregulation of mTOR. TSC is caused by mutations in either the *TSC1* (chromosome 9q34) or *TSC2* (chromosome 16p.13.3) genes. Their protein products, hamartin and tuberin, respectively, form a dimer that acts via the GAP protein Rheb (Ras homolog enhanced in brain) to directly inhibit mTOR, again resulting in upregulation. Specific inhibitors of mTOR are in clinical use, including sirolimus, everolimus, temsirolimus, and deforolimus. Everolimus has been shown to reduce the volume and appearance of subependymal giant cell astrocytomas (SEGA), facial angiofibromas, and renal angiomyolipomas associated with TSC, with a recent FDA approval for SEGA not suitable for surgical resection. This article reviews the use of mTOR inhibitors in these diseases, which have the potential to be a disease-modifying therapy in these and other conditions.

Keywords Neurofibromatosis · Tuberous sclerosis complex · mTOR inhibition · Rapamycin · Everolimus
Molecular therapies

Introduction

Cure rates for most pediatric cancers have been increasing over the past several decades [1–4]; however, recently, survival rates have not continued to rise, possibly reflecting that current chemotherapeutic doses have been maximized. For patients with refractory acute lymphoblastic leukemia (ALL) or solid tumors, the development of novel, biologically targeted treatment strategies is crucial. These rather different diagnoses have a common pathway that is dysregulated or aberrantly expressed through the PI3K/AKT/mammalian target of rapamycin (mTOR) pathway [5–10].

mTOR is a serine/threonine kinase that acts as a central cellular regulator of many functions, including proliferation, cell size/growth, translation, metabolism, autophagy, angiogenesis, and survival by responding to the availability of nutrients (glucose, amino acids), as illustrated in Fig. 1 [11]. mTOR exerts control on these processes by regulating the protein translation machinery, which increases the synthesis of nutrient and amino acids transporters as well as key pro-growth and survival molecules (eg, HIF-1 α , cyclin D1, and myc) [12]. mTOR combines with several other cellular components to form two distinct complexes, termed mTORC1 and mTORC2 [13]. mTORC1 is thought to primarily regulate cell growth, translation, and autophagy in response to nutrient and energy availability via phosphorylation of its downstream targets, including S6K1, p34cdc2, and 4E-BP1. mTORC2 plays a key role in cytoskeletal rearrangements and cell survival. Phosphorylation by mTORC2 in conjunction with

D. N. Franz (✉)
Departments of Pediatrics and Neurology, Tuberous Sclerosis
Clinic, Cincinnati Children's Hospital Medical Center,
Cincinnati, OH 45229, USA
e-mail: Franz@cchmc.org

B. D. Weiss
Department of Pediatrics, Division of Hematology/Oncology,
Neuroblastoma Program, Cincinnati Children's Hospital Medical
Center,
Cincinnati, OH 45229, USA




Fig. 1 Major components of the mTOR pathway and associated diseases. The mammalian target of rapamycin (mTOR) exists as two complexes (mTORC1 and mTORC2), of which only mTORC1 is inhibited by rapamycin and related compounds. mTOR is a central regulator that senses when a cell is in an appropriate bioenergetic state before committing to cell growth and division. Insulin, hormones, growth factors as well as the availability of carbohydrates (CHO) and amino acids stimulate mTORC1 through activation of PI3 kinase (PI3K) and Akt (also known as protein kinase B) via IRS-1. The energy status (AMP/ATP ratio driven by mitochondrial signals) of the cell modulates mTORC1 via AMP protein kinase (AMPK) and the TSC1/2 complex. Upregulation of mTORC1 activates S6 kinase1 (S6K1) and ribosomal S6 kinase (rS6), and inhibits eukaryotic translation initiation factor 4E-binding protein 1 (4EBP1) and eukaryotic initiation factor 4E (eIF4E), which results in cell proliferation, increased transcription of genes, which regulate cell metabolism and proliferation, an increase in ribosomal protein synthesis, cap-dependent translation, eEF2-mediated elongation, and promotes cytoskeletal rearrangements. Conversely, activated mTORC1 blocks apoptosis and autophagy. HIF—hypoxia-inducible factors; LKB1—liver kinase B1; PTEN—phosphatase on chromosome 10; Rheb—Ras homolog enhanced in brain

phosphoinositide-dependent kinase 1 is necessary to fully activate AKT [14]. This review looks at the genetics and clinical implications of mutations disrupting mTOR signaling in two autosomal-dominant genetic disorders, neurofibromatosis type 1 (NF1) and tuberous sclerosis complex (TSC), and evidence for the utility of pharmaceutical agents that target the mTOR pathway for treating them.

Neurofibromatosis Type 1

NF1 is a progressive disorder with an incidence of 1:3500 [15] that is caused by a mutation in the *NF1* tumor suppressor gene, located on chromosome 17q11.2 and comprised of 60 exons spanning 350 kb of genomic DNA. Mutation analysis of the *NF1* gene allows identification of 95% of mutations with a wide spectrum of mutations [16]. To date, no phenotype-genotype correlations have been made with the exception of complete loss of the *NF1* gene (megabase deletions), which is associated with severe mental retardation.

The diagnosis of NF1 is clinical and based on criteria developed at the National Institutes of Health consensus conference (Table 1) [17]. NF1 is characterized by diverse, progressive cutaneous, neurological, skeletal, and neoplastic manifestations, with no standard drug treatment options available. Patients with NF1 have an increased risk of developing both benign and malignant tumors of the central and peripheral nervous system including plexiform neurofibromas (27%), optic gliomas (15–20%), pheochromocytomas (1%), and malignant peripheral nerve sheath tumors (MPNSTs) (10%) [15, 18]. MPNSTs can arise in patients with or without NF1, but in patients with NF1 they tend to occur earlier (2nd and 3rd decade of life) and have a poorer response to therapy and, therefore, a worse prognosis [19]. The lifetime risk of developing MPNSTs in patients with NF1 is 8–13% [20]. MPNSTs can arise spontaneously or by malignant transformation of plexiform neurofibroma.

There are multiple types of neurofibromas including localized cutaneous, localized intraneuronal, diffuse, massive soft tissue, and plexiform neurofibromas. Plexiform neurofibromas are benign nerve sheath tumors that grow along the length of nerves and involve multiple branches of a nerve. These tumors are usually diagnosed early in life, may be multiple, and can develop throughout life. Early childhood, puberty, and childbearing age in females are considered to be the periods of greatest risk for disease progression [21]. Between 20 and 44% of individuals with NF1 develop plexiform neurofibromas [22]. While considered “benign,” these tumors may cause significant disfigurement, as well as compression of vital structures. For example, plexiform neurofibromas may infiltrate the orbit, displace the globe, and compromise vision; paraspinal tumors (also referred to as dumbbell lesions) can compress the spinal cord and cause paralysis; tumors in mediastinum may compress the trachea or great vessels; and tumors of the extremities can cause local nerve infiltration, progressive neurologic deficit, and often cause unremitting pain [21].

Plexiform neurofibromas, including paraspinal neurofibromas, cause major morbidity and mortality in NF1 [18, 23, 24]. The rate of growth of this histologically benign neoplasm has been described as unpredictable and often episodic [15], and the only known cure of these tumors is complete surgical excision. However, complete excision is often not possible due to the infiltrating nature of these tumors. In addition, up to 44% of tumors progress after the first surgery, most commonly in patients younger than 10 years of age with head and neck tumors that could not be completely resected [21]. Also, the complex, infiltrating nature of plexiform neurofibromas can make tumor growth difficult to detect without the use of magnetic resonance imaging with three-dimensional volumetric analysis, a technology not widely available [25].

Patients with NF1 offer unique challenges to the design of clinical trials to test new medications for plexiform

Table 1 NIH diagnostic criteria for neurofibromatosis

Clinical characteristics: At least 2 of the following

- Six or more café-au-lait spots (≥ 0.5 cm in prepubertal subjects or ≥ 1.5 cm in postpubertal subjects)
- Two or more cutaneous or subcutaneous neurofibromas or one plexiform neurofibroma
- Freckling in the axilla or groin
- Optic glioma
- Two or more Lisch nodules
- A distinctive bony lesion (dysplasia of the sphenoid bone or dysplasia or thinning of long bone cortex)
- A first-degree relative with NF1

NIH National Institutes of Health; NF1 neurofibromatosis type 1

Data from [17]

neurofibromas. First, plexiform neurofibromas may not change appreciably for 12–18 months without any intervention, which means that trials must treat patients for long periods before benefit can be detected. In addition, treating a patient with a benign condition on new drugs with which we have minimal experience is a risk most are reasonably unwilling to take. A number of medical treatments including thalidomide, cis-retinoic acid, interferon α -2b, methotrexate and vinblastine, the farnesyltransferase inhibitor tipifarnib (R115777), and the antifibrotic agent pirfenidone have been evaluated in NF1 patients with plexiform neurofibromas with the goal of reducing the size or growth of these tumors [26–28]. However, to date, no medical treatment has demonstrated a clear benefit for patients with NF1 and plexiform neurofibromas.

Tuberous Sclerosis Complex

TSC is a multisystem genetic disorder characterized by growth of hamartomas in multiple organs, including the lung, kidney, and skin [29–31]. About 70–85% of individuals who have a definitive diagnosis of TSC have an identifiable *TSC1* or *TSC2* mutation [29, 32, 33]. *TSC1*, located on chromosome 9q34, encodes the protein hamartin; *TSC2*, located on chromosome 16p13.3, encodes tuberlin. *TSC2* mutations are present in 70–80% of affected individuals and are consistently associated with more severe clinical disease, whereas *TSC1* mutations are only present in 20–30% of individuals.

Affecting both adults and children, TSC is inherited in an autosomal-dominant fashion, but it occurs sporadically in two thirds of cases [29]. The true prevalence of TSC is unknown; however, it is estimated at 1:6000–10,000 and it currently affects 1.5 million people worldwide [32, 34, 35]. A wide variation exists in the level of disease severity and, in fact, some patients with TSC remain asymptomatic. As such, the diagnostic criteria for TSC consist of a set of major

and minor features that guide the level of probability of disease into definite, probable, or possible categories (Table 2) [36]. Virtually any organ system can be affected in patients with TSC, but the disease is most commonly characterized by central nervous system (CNS) manifestations and by the presence of hamartomatous lesions in other organs. Although historically thought to be benign, recent evidence suggests that hamartomas may have an underlying neoplastic nature [37]. In particular, a variant of angiomyolipoma (AML), called epithelioid AML, is an extremely aggressive tumor [38].

Anatomic and functional abnormalities of the CNS that result in neurologic dysfunction contribute significantly to morbidity and mortality in patients with TSC. Brain lesions include cortical tubers, subependymal nodules (SENs), and subependymal giant cell astrocytomas (SEGAs). Cortical tubers are found in 80–90% of patients [29, 32] and are associated with seizures and developmental delays [39–41], and consequently represent an area of high interest for further research. Tubers and the surrounding cortex may have to be surgically removed, often successfully, to treat patients with intractable epilepsy [41, 42]. SENs occur in about 90% of patients with TSC, and are asymptomatic nodular lesions that extend into the ventricles of the CNS. SENs usually remain dormant but have the potential to increase in size and develop into SEGAs [43]. SEGAs are slow-growing, glioneuronal tumors that develop in approximately 5–20% of patients [29, 44, 45]. SEGAs generally develop adjacent to the foramen of Monro and may remain clinically benign until they reach sufficient size to cause ventricular obstruction. Once this occurs, the potential for sudden death secondary to acute hydrocephalus is increased [43, 44]. SEGAs may also produce visual impairment, endocrinopathies, and focal neurologic deficits. SEGAs are often large and difficult to resect once they produce clinical symptoms.

A hallmark of TSC is epilepsy, which can be seen in up to 90% of patients. Epilepsy tends to increase in severity with

Table 2 Common clinical manifestations and diagnostic criteria for TSC

| Clinical characteristic | Diagnosis |
|--|--|
| Major features | Definite TSC: Either 2 major features or 1 major feature plus 2 minor features |
| Facial angiofibromas or forehead plaque | |
| Nontraumatic ungula or periungual fibroma | |
| Hypomelanotic macules (three or more) | |
| Shagreen patch migration lines | |
| Multiple retinal nodular hamartomas | |
| Cortical tuber ^a | |
| Subependymal nodule | |
| Subependymal giant cell astrocytoma | |
| Cardiac rhabdomyoma, single or multiple | |
| Lymphangiomyomatosis, renal angiomyolipoma, or both ^b | Probable TSC: 1 major plus 1 minor feature |
| Minor features | |
| Multiple, randomly distributed pits in dental enamel | |
| Hamartomatous rectal polyps ^c | |
| Bone cysts ^d | |
| Cerebral white matter radial migration lines ^{a,d} | |
| Gingival fibromas | |
| Nonrenal hamartoma ^c | |
| Retinal achromic patch | |
| Confetti-like skin lesions | |
| Multiple renal cysts ^c | Possible TSC: Either 1 major feature or 2 or more minor features |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |
| | |

^a When cerebral cortical dysplasia and cerebral white matter migration tracts occur together, they should be counted as one rather than two features of tuberous sclerosis

^b When both lymphangiomyomatosis and renal angiomyolipomas are present, other features of tuberous sclerosis should be present before a definite diagnosis is assigned

^c Histologic confirmation is suggested

^d Radiographic confirmation is sufficient

TSC tuberous sclerosis complex

Data from Anlauf et al. [34]

age and usually is difficult to control [32, 46]. Although complex partial seizures are most commonly observed, most seizure types can occur [35, 47]. A particularly devastating type of seizure, infantile spasms (also known as West syndrome), is seen in approximately one third of very young children with TSC with usual onset during the first year of life [47, 48]. The seizures generally stop by about age 5 years, but may be replaced by other seizure types. If not controlled, infantile spasms are associated with severe developmental delay and cognitive impairment. Developmental

and behavioral difficulties, including mental retardation, learning disabilities, psychiatric disorders, cognitive impairment, autism, or attention deficit/hyperactivity disorder, are also common in patients with TSC [32, 43, 46, 49–52].

The majority (90%) of patients with TSC have skin involvement [29, 30]. The hallmark lesion is the ash leaf macule, an ovoid hypopigmented lesion most noticeable in dark-skinned individuals or those with sun exposure. Affected individuals also develop cutaneous fibromas, which have a predilection for the face (angiofibromas, previously adenoma sebaceum), periungual tissue, lower back, and gingivae. Nonpharmaceutical therapies such as laser ablation or surgical resection are the standard therapies for TSC-associated skin lesions [53].

Renal AMLs are the second leading cause of death in patients with TSC, occur in approximately 75% of patients, and usually present with multiple lesions in both kidneys [29, 54]. Tumors greater than 4 cm in diameter are more likely to become symptomatic with potentially life-threatening hemorrhage, and are usually slated for treatment with nephron-sparing prophylactic selective arterial embolization or radio-frequency ablation [54, 55]. Nephrectomy is reserved for multiple large tumors that involve most of the kidney and are not amenable to embolization.

Pulmonary lymphangiomyomatosis (LAM) is the third leading cause of death in patients with TSC and occurs almost exclusively in premenopausal females [29, 32]. Although 40% of females with TSC show radiographic evidence of LAM, only about 5% of patients are functionally affected, but in these patients the disorder is almost invariably fatal [52, 54]. Currently, there is no effective treatment for LAM; in addition to symptomatic treatment, progesterone has been administered (due to the presumed involvement of hormones in the pathogenesis of LAM) with variable success [29, 53, 56]. Lung transplantation is reserved for patients with respiratory failure.

Other common lesions include cardiac rhabdomyomas and retinal lesions [30, 57, 58]. TSC-related cardiac rhabdomyomas generally regress with increasing age; therefore, treatment of this lesion is rarely required [53]. For symptomatic treatment of visual impairment, standard treatment options are photocoagulation and photodynamic therapy.

Genetics of TSC and NF1: Role of the mTOR Pathway

The *NF1* gene encodes a protein, termed neurofibromin, which functions partly as a Ras-GTPase activating protein (RasGAP). Neurofibromin loss in tumor cells leads to hyperactivation of Ras and its downstream signaling intermediates, which are critical for transmitting the Ras growth signal and thus the development of neoplasia in patients with NF1. One of these downstream proteins is the mTOR

molecule, which was found to be activated in both NF1-deficient primary human and mouse cells as well as in human and genetically engineered *Nf1* mouse tumor models. This aberrant activation was dependent on Ras and PI3 kinase/AKT signaling [6, 7]. The mTOR inhibitor sirolimus was shown to dramatically reduce the increased proliferation associated with loss of neurofibromin expression in human MPNST cell lines; and in a genetic mouse model of NF1-deficient MPNST development, sirolimus completely inhibited the growth of these tumors in vivo [6].

Mice carrying compound mutations in the NF-1 and p53 tumor suppressors on the same chromosome (NPcis animals) develop aggressive MPNSTs that are histologically indistinguishable from human tumors with an average latency of 5 months. These lesions develop as a result of somatic loss of the wild-type *Nf1* and p53 alleles, and therefore are also genetically similar to human MPNSTs. NPcis mice with palpable tumors (approximately 300 mm³) that were treated with sirolimus had significantly improved survival and tumor shrinkage compared to control mice [59].

These findings identify the *NF-1* tumor suppressor as a negative regulator of mTOR, and demonstrate that sirolimus blocks the growth of several *NF1*-deficient cells and tumor cell types in vitro and in vivo. The combination of cell culture and preclinical mouse modeling data provides a strong rationale for the use of sirolimus in treating human NF1-associated tumors.

In the case of TSC, the normal TSC1 and TSC2 proteins form a complex in cells to function as a GTPase-activating protein toward Ras homolog enriched in brain (Rheb) [60, 61]. Through Rheb, the TSC1/TSC2 protein (hamartin-tuberin) complex limits the activation of mTOR. Mutation of the *TSC1* and/or *TSC2* genes leads to a deficient hamartin-tuberin complex, and results in constitutive activation of mTOR leading to hyperactive mTOR signaling and abnormal cell division.

Evidence from in vitro, in murine models, and in humans indicates that a wide spectrum of *TSC1/TSC2* mutations resulting in mTOR hyperactivity is responsible for the developmental abnormalities in the brain that occur in patients with TSC. Developing neurons acquire an abnormal morphology, with large cell bodies, multiple axons, and few, large, dendritic spines, decreased myelin, and abnormal synaptic architecture [40, 62–68]. Increased neuronal proliferation and aberrant neuronal migration lead to abnormal cortical architecture [29, 65–67]. Developing glial cells have also exhibited an abnormal morphology and increased growth and proliferation [40, 66, 69, 70].

Evidence from murine models suggests that mTOR inhibition may be able to at least partly correct abnormal synaptic transmission, synaptic plasticity, and neuronal excitability, and thereby has the potential to prevent progression or development of neurologic dysfunction, including epilepsy and

cognitive/behavioral dysfunction in patients with TSC [49, 66, 71–74].

Clinical Trials Evaluating mTOR Inhibition in NF1 and TSC

There are agents currently available and others under preclinical and clinical development that target signaling nodes within the PI3K/AKT/mTOR pathway [10, 23, 75, 76]. Rapamycin and its rapalogs (eg, everolimus, deforolimus, and temsirolimus) are allosteric inhibitors of mTORC1, whereas mTORC2 is generally resistant to rapalogs, but this resistance may vary by cell type [77–79]. Rapamycin (sirolimus) was isolated from the bacterial strain *Streptomyces hygroscopicus* found in the soil on Easter Island (“Rapa Nui” in the island’s native language) [80]. While its anti-neoplastic properties were identified early on, rapamycin was not further developed for this indication because of its variable bioavailability and its aqueous insolubility [80]. Rapamycin is also an effective immunosuppressant and is currently US Food and Drug Administration (FDA) approved as an anti-rejection medication in solid organ and bone marrow transplants [81].

There are several ongoing clinical trials involving pharmacologic inhibition of mTOR in patients with NF1. The US Department of Defense (DOD) funded Neurofibroma (NF) Consortium is currently conducting a trial of the oral mTOR inhibitor sirolimus for children and adults with NF1-associated plexiform neurofibromas. The DOD NF Consortium is also conducting a trial of the oral mTOR inhibitor everolimus for children with NF1 and chemotherapy-refractory progressive low-grade gliomas (optic pathway gliomas). Everolimus is one of the second-generation, aqueous rapalogs with much improved bioavailability compared to sirolimus. These trials are ongoing and thus no published results are available.

A clinical series has demonstrated responses to single-agent sirolimus in SEGAs and a low-grade glioma in five pediatric patients with TSC [82]. Everolimus has demonstrated significant reductions in tumor volume in SEGAs associated with TSC, which led to the approval of the drug by the FDA for this indication [83, 84]. Primary SEGA volume, as assessed on independent central review ($P < 0.001$ for baseline vs 6 months), was reduced by $\geq 30\%$ in 21 patients (75%) and at least 50% in 9 patients (32%) [83]. There were no new lesions, worsening hydrocephalus, evidence of increased intracranial pressure, or necessity for surgical resection or other therapy for SEGAs. In addition, a marked reduction in seizure frequency was seen [83].

Some early clinical trials suggest that mTOR inhibitors effectively and safely treat TSC-associated renal lesions by causing tumor regression when on active treatment [30, 72, 85–88]. Although no data exist for the use of mTOR inhibitors

in the treatment of cardiac rhabdomyomas, abnormalities in the mTOR signaling pathway have been confirmed [89, 90]. Therefore, it is plausible that an mTOR inhibitor may be used to treat cardiac rhabdomyoma. Some suggest that mTOR inhibitors also may have a role in the treatment of angiofibroma, as small and early-phase clinical trials have demonstrated some regression in these lesions [87].

Conclusions

To date, treatment options for patients with NF1 or TSC have been limited and have demonstrated variable rates of success. In both NF1 and TSC, neurofibromin and the hamartin/tuberin complex, respectively, act as “molecular brakes” limiting activation of downstream signaling intermediates. mTOR, a common downstream intermediate to both neurofibromin and hamartin/tuberin, appears to play an integral role in the development of NF1 and TSC and their complications. Pre-clinical data and early clinical trials have demonstrated a beneficial effect of mTOR inhibitors in the treatment of several clinical manifestations of TSC, including SEGA and renal AML volume, epilepsy, and skin lesions. Ongoing clinical trials evaluating sirolimus and everolimus in NF1 will evaluate the potential role of these agents for treating this condition. Inhibition of mTOR is a rational target and has the potential to change the paradigm of how patients are treated.

Disclosure Conflicts of interest: D.N. Franz: has received grant support for study drug and costs for conducting the research described in this paper, Novartis; has received a consulting fee or honorarium for conduct and design of clinical trials, Novartis; has received support for travel costs to meet with regulatory agencies, present research results, design protocols, Novartis; has received support for study medication, administrative support for study conduct, Novartis; has given expert testimony for various plaintiff's and defendant's attorneys; has received payment of travel and lodging expenses for invited lectures, not as part of a speaker's bureau; is on the speaker's bureau for UCB Pharma; B.D. Weiss: none.

References

- Pui CH, Evans WE. Treatment of acute lymphoblastic leukemia. *N Engl J Med*. 2006;354(2):166–78.
- Gaynon PS, Trigg ME, Heerema NA, Sather HN, Hammond GD, et al. Children's Cancer Group trials in childhood acute lymphoblastic leukemia: 1983–1995. *Leukemia*. 2000;14(12):2223–33.
- Rechnitzer C. Increased survival of children with solid tumours: how did we get there and how to keep the success going? *Canc Imag*. 2011;11:865–9.
- Howlader N, Noone AM, Krapcho M, Neyman N, Aminou R, Waldron W, et al. SEER Cancer Statistics Review, 1975–2008. National Cancer Institute, Bethesda, MD; 2010.
- Stoddart A, Fleming HE, Paige CJ. The role of the preBCR, the interleukin-7 receptor, and homotypic interactions during B-cell development. *Immunol Rev*. 2000;175:47–58.
- Johannessen CM, Reczek EE, James MF, Brems H, Legius E, Cichowski K. The NF1 tumor suppressor critically regulates TSC2 and mTOR. *Proc Natl Acad Sci U S A*. 2005;102(24):8573–8.
- Dasgupta B, Yi Y, Chen DY, Weber JD, Gutmann DH. Proteomic analysis reveals hyperactivation of the mammalian target of rapamycin pathway in neurofibromatosis 1-associated human and mouse brain tumors. *Cancer Res*. 2005;65(7):2755–60.
- Perry B, Banyard J, McLaughlin ER, Watnick R, Sohn A, Brindley DN, et al. AKT1 overexpression in endothelial cells leads to the development of cutaneous vascular malformations in vivo. *Arch Dermatol*. 2007;143(4):504–6.
- Vivanco I, Sawyers CL. The phosphatidylinositol 3-Kinase AKT pathway in human cancer. *Nat Rev Cancer*. 2002;2(7):489–501.
- Chiarini F, Fala F, Tazzari PL, Ricci F, Astolfi A, Pession A, et al. Dual inhibition of class IA phosphatidylinositol 3-kinase and mammalian target of rapamycin as a new therapeutic option for T-cell acute lymphoblastic leukemia. *Cancer Res*. 2009;69(8):3520–8.
- Houghton PJ, Huang S. mTOR as a target for cancer therapy. *Curr Top Microbiol Immunol*. 2004;279:339–59.
- Bjornsti MA, Houghton PJ. The TOR pathway: a target for cancer therapy. *Nat Rev Cancer*. 2004;4(5):335–48.
- Bhaskar PT, Hay N. The two TORCs and Akt. *Developmental Cell*. 2007;12(4):487–502.
- Sarbassov DD, Guertin DA, Ali SM, Sabatini DM. Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science*. 2005;307(5712):1098–101.
- Korf BR. Malignancy in neurofibromatosis type 1. *Oncologist*. 2000;5(6):477–85.
- Messiaen LM, Callens T, Mortier G, Beysen D, Vandenbroucke I, Van RN, et al. Exhaustive mutation analysis of the NF1 gene allows identification of 95% of mutations and reveals a high frequency of unusual splicing defects. *Hum Mutat*. 2000;15(6):541–55.
- US Department of Health and Human Services, National Institutes of Health. Neurofibromatosis. NIH Consens Statement Online 1987 Jul 13–15;6(12):1–19.
- Korf BR. Plexiform neurofibromas. *Am J Med Genet*. 1999;89(1):31–7.
- Carli M, Ferrari A, Mattke A, et al. Pediatric malignant peripheral nerve sheath tumor: The Italian and German soft tissue sarcoma cooperative group. *J Clin Oncol*. 2005;23:8422–30.
- Evans DG, Baser ME, McGaughran J, Sharif S, Howard E, Moran A. Malignant peripheral nerve sheath tumours in neurofibromatosis. *J Mol Genet*. 2002;39:311–4.
- Needle MN, Cnaan A, Dattilo J, Chatten J, Phillips PC, Shochat S, et al. Prognostic signs in the surgical management of plexiform neurofibroma: the Children's Hospital of Philadelphia experience, 1974–1994. *J Pediatr*. 1997;131(5):678–82.
- Huson SM, Harper PS, Compston DA. Von Recklinghausen neurofibromatosis. A clinical and population study in south-east Wales. *Brain*. 1988;111(Pt 6):1355–81.
- Ricardi VM, Eichner JE. Neurofibromatosis: phenotype, natural history and pathogenesis. Baltimore: Johns Hopkins University Press; 1986.
- Rasmussen SA, Yang Q, Friedman JM. Mortality in neurofibromatosis 1: an analysis using U.S. death certificates. *Am J Hum Genet*. 2001;68(5):1110–8.
- Solomon J, Warren K, Dombi E, et al. Automated detection and volume measurement of plexiform neurofibromas in neurofibromatosis 1 using magnetic resonance imaging. *Comput Med Imag Graph*. 2004;28:257–65.

26. Gupta A, Cohen BH, Ruggieri P, Packer RJ, Phillips PC. Phase I study of thalidomide for the treatment of plexiform neurofibroma in neurofibromatosis 1. *Neurology*. 2003;60(1):130–2.
27. Widemann BC, Salzer WL, Arceci RJ, Blaney SM, Fox E, End D, et al. Phase I trial and pharmacokinetic study of the farnesyltransferase inhibitor tipifarnib in children with refractory solid tumors or neurofibromatosis type I and plexiform neurofibromas. *J Clin Oncol*. 2006;24(3):507–16.
28. Babovic-Vuksanovic D, Widemann BC, Dombi E, Gillespie A, Wolters PL, Toledo-Tamula MA, et al. Phase I trial of pirfenidone in children with neurofibromatosis 1 and plexiform neurofibromas. *Pediatr Neurol*. 2007;36(5):293–300.
29. Crino PB, Nathanson KL, Henske EP. The tuberous sclerosis complex. *N Engl J Med*. 2006;355(13):1345–56.
30. Curatolo P, Bombardieri R, Jozwiak S. Tuberous sclerosis. *Lancet*. 2008;372(9639):657–68.
31. Franz DN, Bissler JJ, McCormack FX. Tuberous sclerosis complex: neurological, renal and pulmonary manifestations. *Neuropediatrics*. 2010;41(5):199–208.
32. Baskin Jr HJ. The pathogenesis and imaging of the tuberous sclerosis complex. *Pediatr Radiol*. 2008;38(9):936–52.
33. Sancak O, Nellist M, Goedbloed M, Elfferich P, Wouters C, Maat-Kievit A, et al. Mutational analysis of the TSC1 and TSC2 genes in a diagnostic setting: genotype: phenotype correlations and comparison of diagnostic DNA techniques in tuberous sclerosis complex. *Eur J Hum Genet*. 2005;13(6):731–41.
34. Anlauf M, Garbrecht N, Bauersfeld J, Schmitt A, Henopp T, Komminoth P, et al. Hereditary neuroendocrine tumors of the gastroenteropancreatic system. *Virchows Arch*. 2007;451 Suppl 1:S29–38.
35. Holmes GL, Stafstrom CE. Tuberous sclerosis complex and epilepsy: recent developments and future challenges. *Epilepsia*. 2007;48(4):617–30.
36. Roach ES, Gomez MR, Northrup H. Tuberous sclerosis complex consensus conference: revised clinical diagnostic criteria. *J Child Neurol*. 1998;13(12):624–8.
37. Mete O, van der Kwast TH. Epithelioid angiomyolipoma: a morphologically distinct variant that mimics a variety of intra-abdominal neoplasms. *Arch Pathol Lab Med*. 2011;135(5):665–70.
38. Martignoni G, Pea M, Reghellin D, Zamboni G, Bonetti F. PEComas: the past, the present and the future. *Virchows Archive*. 2008;452(2):119–32.
39. Kwiatkowski DJ, Manning BD. Tuberous sclerosis: a GAP at the crossroads of multiple signaling pathways. *Hum Mol Genet*. 2005;14(2):R251–8.
40. Wong M. Mechanisms of epileptogenesis in tuberous sclerosis complex and related malformations of cortical development with abnormal glioneuronal proliferation. *Epilepsia*. 2008;49(1):8–21.
41. Major P, Rakowski S, Simon MV, Cheng ML, Eskandar E, Baron J, et al. Are cortical tubers epileptogenic? Evidence from electrocorticography. *Epilepsia*. 2009;50(1):147–54.
42. Weiner HL, Ferraris N, LaJoie J, Miles D, Devinsky O. Epilepsy surgery for children with tuberous sclerosis complex. *J Child Neurol*. 2004;19(9):687–9.
43. Kopp CM, Muzykewicz DA, Staley BA, Thiele EA, Pulsifer MB. Behavior problems in children with tuberous sclerosis complex and parental stress. *Epilepsy Behavior*. 2008;13(3):505–10.
44. Goh S, Butler W, Thiele EA. Subependymal giant cell tumors in tuberous sclerosis complex. *Neurology*. 2004;63(8):1457–61.
45. Adriaansen ME, Schaefer-Prokop CM, Stijnen T, Duynham DA, Zonnenberg BA, Prokop M. Prevalence of subependymal giant cell tumors in patients with tuberous sclerosis and a review of the literature. *Eur J Neurol*. 2009;16(6):691–6.
46. Prather P, de Vries PJ. Behavioral and cognitive aspects of tuberous sclerosis complex. *J Child Neurol*. 2004;19(9):666–74.
47. Thiele EA. Managing epilepsy in tuberous sclerosis complex. *J Child Neurol*. 2004;19(9):680–6.
48. Tsao CY. Current trends in the treatment of infantile spasms. *Neuropsychiatric Diseases and Treatment*. 2009;5:289–99.
49. Ehninger D, de Vries PJ, Silva AJ. From mTOR to cognition: molecular and cellular mechanisms of cognitive impairments in tuberous sclerosis. *J Intellect Disabil Res*. 2009;53(10):838–51.
50. D'Agati E, Moavero R, Cerminara C, Curatolo P. Attention-deficit hyperactivity disorder (ADHD) and tuberous sclerosis complex. *J Child Neurol*. 2009;24(10):1282–7.
51. Winterkorn EB, Pulsifer MB, Thiele EA. Cognitive prognosis of patients with tuberous sclerosis complex. *Neurology*. 2007;68(1):62–4.
52. Johnson SR. Lymphangioleiomyomatosis. *Eur Respir J*. 2006;27(5):1056–65.
53. Yates JR. Tuberous sclerosis. *Eur J Hum Genet*. 2006;14(10):1065–73.
54. Franz DN. Non-neurologic manifestations of tuberous sclerosis complex. *J Child Neurol*. 2004;19(9):690–8.
55. Sooriakumaran P, Gibbs P, Coughlin G, Attard V, Elmslie F, Kingswood C, et al. Angiomyolipomata: challenges, solutions, and future prospects based on over 100 cases treated. *BJU Int*. 2010;105(1):101–6.
56. McCormack FX. Lymphangioleiomyomatosis: a clinical update. *Chest*. 2008;133(2):507–16.
57. Jozwiak S, Jozwiak S, Oldak M. Molecular activity of sirolimus and its possible application in tuberous sclerosis treatment. *Med Res Rev*. 2006;26(2):160–80.
58. Isaacs H. Perinatal (fetal and neonatal) tuberous sclerosis: a review. *Am J Perinatol*. 2009;26(10):755–60.
59. Johannessen CM, Johnson BW, Williams SM, Chan AW, Reczek EE, Lynch RC, et al. TORC1 is essential for NF1-associated malignancies. *Curr Biol*. 2008;18(1):56–62.
60. Napolioni V, Curatolo P. Genetics and molecular biology of tuberous sclerosis complex. *Curr Genom*. 2008;9(7):475–87.
61. Dowling RJ, Topisirovic I, Fonseca BD, Sonenberg N. Dissecting the role of mTOR: lessons from mTOR inhibitors. *Biochim Biophys Acta*. 2010;1804(3):433–9.
62. Choi YJ, Di Nardo A, Kramvis I, Meikle L, Kwiatkowski DJ, Sahin M, et al. Tuberous sclerosis complex proteins control axon formation. *Genes Dev*. 2008;22(18):2485–95.
63. de Vries PJ, Howe CJ. The tuberous sclerosis complex proteins: a GRIPP on cognition and neurodevelopment. *Trends Mol Med*. 2007;13(8):319–26.
64. Krishnan ML, Commowick O, Jeste SS, Weisenfeld N, Hans A, Gregas MC, et al. Diffusion features of white matter in tuberous sclerosis with tractography. *Pediatr Neurol*. 2010;42(2):101–6.
65. Meikle L, Talos DM, Onda H, Polizzi K, Rotenberg A, Sahin M, et al. A mouse model of tuberous sclerosis: neuronal loss of Tsc1 causes dysplastic and ectopic neurons, reduced myelination, seizure activity, and limited survival. *J Neurosci*. 2007;27(21):5546–58.
66. Napolioni V, Moavero R, Curatolo P. Recent advances in neurobiology of tuberous sclerosis complex. *Brain Dev*. 2009;31(2):104–13.
67. Swiech L, Perycz M, Malik A, Jaworski J. Role of mTOR in physiology and pathology of the nervous system. *Biochim Biophys Acta*. 2008;1784(1):116–32.
68. Tavazoie SF, Alvarez VA, Ridenour DA, Kwiatkowski DJ, Sabatini BL. Regulation of neuronal morphology and function by the tumor suppressors Tsc1 and Tsc2. *Nat Neurosci*. 2005;8(12):1727–34.
69. Crino PB. Molecular pathogenesis of tuber formation in tuberous sclerosis complex. *J Child Neurol*. 2004;19(9):716–25.
70. Moavero R, Cerminara C, Curatolo P. Epilepsy secondary to tuberous sclerosis: lessons learned and current challenges. *Childs Nerv Syst*. 2010;26(11):1495–504.
71. Ehninger D, Han S, Shilyansky C, Zhou Y, Li W, Kwiatkowski DJ, et al. Reversal of learning deficits in a Tsc2^{+/-} mouse model of tuberous sclerosis. *Nat Med*. 2008;14(8):843–8.

72. Sampson JR. Therapeutic targeting of mTOR in tuberous sclerosis. *Biochem Soc Trans.* 2009;37(Pt 1):259–64.
73. Wong M. Mammalian target of rapamycin (mTOR) inhibition as a potential antiepileptogenic therapy: from tuberous sclerosis to common acquired epilepsies. *Epilepsia.* 2010;51(1):27–36.
74. Zeng LH, Xu L, Gutmann DH, Wong M. Rapamycin prevents epilepsy in a mouse model of tuberous sclerosis complex. *Ann Neurol.* 2008;63(4):444–53.
75. Guertin DA, Sabatini DM. The pharmacology of mTOR inhibition. *Sci Signal.* 2009;2(67):e24.
76. Brown VI, Seif AE, Reid GS, Teachey DT, Grupp SA. Novel molecular and cellular therapeutic targets in acute lymphoblastic leukemia and lymphoproliferative disease. *Immunol Res.* 2008;42(1–3):84–105.
77. Akcakanat A, Singh G, Hung MC, Meric-Bernstam F. Rapamycin regulates the phosphorylation of rictor. *Biochem Biophys Res Commun.* 2007;362(2):330–3.
78. Sarbassov DD, Ali SM, Sengupta S, Sheen JH, Hsu PP, Bagley AF, et al. Prolonged rapamycin treatment inhibits mTORC2 assembly and Akt/PKB. *Mol Cell.* 2006;22(2):159–68.
79. Zeng Z, Sarbassov DD, Samudio IJ, Yee KW, Munsell MF, Ellen JC, et al. Rapamycin derivatives reduce mTORC2 signaling and inhibit AKT activation in AML. *Blood.* 2007;109(8):3509–12.
80. Sehgal SN, Baker H, Vezina C. Rapamycin (AY-22,989), a new antifungal antibiotic. II. Fermentation, isolation and characterization. *J Antibiot (Tokyo).* 1975;28(10):727–32.
81. Saunders RN, Metcalfe MS, Nicholson ML. Rapamycin in transplantation: a review of the evidence. *Kidney Int.* 2001;59(1):3–16.
82. Franz DN, Leonard J, Tudor C, Chuck G, Care M, Sethuraman G, et al. Rapamycin causes regression of astrocytomas in tuberous sclerosis complex. *Ann Neurol.* 2006;59(3):490–8.
83. Krueger DA, Care MM, Holland K, Agricola A, Tudor C, Mangeshkar P, et al. Everolimus for subependymal giant-cell astrocytomas in tuberous sclerosis. *N Engl J Med.* 2010;363(19):1801–11.
84. Afinitor® (everolimus) tablets for oral administration [prescribing information]. East Hanover, NJ: Novartis Pharmaceuticals Corp.; 2010.
85. Bissler JJ, McCormack FX, Young LR, Elwing JM, Chuck G, Leonard JM, et al. Sirolimus for angiomyolipoma in tuberous sclerosis complex or lymphangioleiomyomatosis. *N Engl J Med.* 2008;358(2):140–51.
86. Curatolo P, D'Argenzio L, Cerminara C, Bombardieri R. Management of epilepsy in tuberous sclerosis complex. *Expert Rev Neurother.* 2008;8(3):457–67.
87. Krueger DA, Franz DN. Current management of tuberous sclerosis complex. *Paediatric Drugs.* 2008;10(5):299–313.
88. Ess KC. Tuberous sclerosis complex: a brave new world? *Curr Opin Neurol.* 2010;23(2):189–93.
89. Jozwiak J, Sahin M, Jozwiak S, Kotulska K, Ploski R, Szperl M, et al. Cardiac rhabdomyoma in tuberous sclerosis: hyperactive Erk signaling. *Int J Cardiol.* 2009;132(1):145–7.
90. Kotulska K, Larysz-Brysz M, Grajkowska W, Jozwiak J, Wlodarski P, Sahin M, et al. Cardiac rhabdomyomas in tuberous sclerosis complex show apoptosis regulation and mTOR pathway abnormalities. *Pediatr Dev Pathol.* 2009;12(2):89–95.