

# Climate Change Project

Christian Chimezie

ATM 320

---

## Introduction

The purpose of this project is to determine whether climate-change is happening on a global scale by analyzing atmospheric data recorded over the last 40 year period. Global temperatures are represented in terms of anomalies, which are departures from a long-term average. The Southern Oscillation Index (SOI) is an index based on observed sea-level pressure differences between Tahiti and Australia; this index might be useful in modeling atmospheric temperature trends. We will analyze trends in global temperature anomaly, SOI, and also local temperatures in Albany, New York. We hope that analyzing historical atmospheric data will give insight into the phenomenon of climate change.

## Temperature and SOI Statistics

### 1980 - 1999

Local temperature:

$T_{avg}$ (°C)	$\sigma T$ (°C)	$T_{max}$ (°C)	$T_{min}$ (°C)
8.58	10.48	22.4	-10.0

Global temperature anomaly:

$T_{avg}$ (°C)	$\sigma T$ (°C)	$T_{max}$ (°C)	$T_{min}$ (°C)
0.3	0.1	0.52	0.13

Southern Oscillation Index:

avg	$\sigma$	max	min
0.12	0.93	2.0	-2.1

### 2000 - 2019

Local temperature:

$T_{avg}$ (°C)	$\sigma T$ (°C)	$T_{max}$ (°C)	$T_{min}$ (°C)
9.24	9.6	24.7	-10.7

Global temperature anomaly:

$T_{avg}$ (°C)	$\sigma T$ (°C)	$T_{max}$ (°C)	$T_{min}$ (°C)
0.52	0.26	1.37	-0.04

Southern Oscillation Index:

avg	$\sigma$	max	min
0.03	1.62	4.8	-6.0

## 1980 - 2019

Local temperature:

$T_{avg}$ ( $^{\circ}\text{C}$ )	$\sigma T$ ( $^{\circ}\text{C}$ )	$T_{max}$ ( $^{\circ}\text{C}$ )	$T_{min}$ ( $^{\circ}\text{C}$ )
9.21	9.64	24.7	-10.7


Global temperature anomaly:

$T_{avg}$ ( $^{\circ}\text{C}$ )	$\sigma T$ ( $^{\circ}\text{C}$ )	$T_{max}$ ( $^{\circ}\text{C}$ )	$T_{min}$ ( $^{\circ}\text{C}$ )
0.51	0.26	1.37	-0.04


Southern Oscillation Index:

avg	$\sigma$	max	min
0.04	1.6	4.8	-6.0


## Visualizing Temperature and SOI Trends




**Figure 1.** Degree 1 polynomial fit of local temperature as a function of time.




**Figure 2.** Degree 1 polynomial fit of global temperature anomaly as a function of time.




**Figure 3.** Degree 1 polynomial fit of SOI as a function of time.




**Figure 4.** Degree 1 polynomial fit of local temperature compared to global temperature anomaly.




**Figure 5.** Degree 1 polynomial fit of local temperature compared to SOI.



**Figure 6.** Histogram of monthly local temperature values.



**Figure 7.** Histogram of monthly global temperature anomaly values.



**Figure 8.** Histogram of SOI values.

## Correlation Coefficient

The correlation coefficient is a quantification of the strength of the linear relationship between two variables. Correlations of -1 or +1 imply an exact linear relationship, while 0 implies no correlation. The Pearson correlation coefficient is calculated using the equation 1.

$$r = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sqrt{\sum (x - \bar{x})^2 \sum (y - \bar{y})^2}} \quad (1)$$

The statistical significance of the correlation coefficient can be interpreted using the p-value method. The p-value  $P$  is calculated using a t-distribution with  $n - 2$  degrees of freedom as shown in equation 2.

$$t = \frac{r\sqrt{n-2}}{\sqrt{1-r^2}} \quad (2)$$

We use  $\alpha = 0.05$  as the threshold significance level.

- If  $P < \alpha$ , the variables *are* linear because  $r$  *is* sufficiently different from zero.
- If  $P > \alpha$ , the variables *are not* linear because  $r$  *is not* sufficiently different from zero.