

3.3 Relation to subgroups — Part 2

November 5, 2021

Last time

Main results for today

Relating $H^\bullet(H, A)$, $H^\bullet(G/H, A^H)$, $H^\bullet(G, A)$

Main results for today

Relating $H^\bullet(H, A)$, $H^\bullet(G/H, A^H)$, $H^\bullet(G, A)$

Let G be a group, A a G -module, and H a subgroup of G . Then there exist group homomorphisms, called **restriction maps**,

$$\text{Res} : H^i(G, A) \rightarrow H^i(H, A) \quad \text{for all } i \geq 0.$$

Main results for today

Relating $H^\bullet(H, A)$, $H^\bullet(G/H, A^H)$, $H^\bullet(G, A)$

Let G be a group, A a G -module, and H a subgroup of G . Then there exist group homomorphisms, called **restriction maps**,

$$\text{Res} : H^i(G, A) \rightarrow H^i(H, A) \quad \text{for all } i \geq 0.$$

If $[G : H] = n$ is finite, there exist group homomorphisms, called **corestriction maps**,

$$\text{Cor} : H^i(H, A) \rightarrow H^i(G, A) \quad \text{for all } i \geq 0.$$

Main results for today

Relating $H^\bullet(H, A)$, $H^\bullet(G/H, A^H)$, $H^\bullet(G, A)$

Let G be a group, A a G -module, and H a subgroup of G . Then there exist group homomorphisms, called **restriction maps**,

$$\text{Res} : H^i(G, A) \rightarrow H^i(H, A) \quad \text{for all } i \geq 0.$$

If $[G : H] = n$ is finite, there exist group homomorphisms, called **corestriction maps**,

$$\text{Cor} : H^i(H, A) \rightarrow H^i(G, A) \quad \text{for all } i \geq 0.$$

If instead H is a normal subgroup of G , there exist group homomorphisms, called **inflation maps**,

$$\text{Inf} : H^i(G/H, A) \rightarrow H^i(G, A) \quad \text{for all } i \geq 0.$$

Main results for today

Relating Res, Cor, Inf

Main results for today

Relating Res, Cor, Inf

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

The tool to help define Res, Cor, Inf:
Shapiro's lemma

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A)$$

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A)$$

with G -action given by taking

$$\phi : \mathbb{Z}[G] \rightarrow A$$

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A)$$

with G -action given by taking

$$\phi : \mathbb{Z}[G] \rightarrow A$$

to

$$g' \cdot \phi : \mathbb{Z}[G] \rightarrow A : g \mapsto \phi(gg')$$

for any $g' \in G$.¹

¹Check the map $g' \cdot \phi$ is again H -equivariant.

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A).$$

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \mathrm{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A).$$

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \mathrm{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Proof.

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A).$$

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \mathrm{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Proof. Fix a projective $\mathbb{Z}[G]$ -resolution P_\bullet of \mathbb{Z} .

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A).$$

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \mathrm{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Proof. Fix a projective $\mathbb{Z}[G]$ -resolution P_\bullet of \mathbb{Z} . Since $\mathbb{Z}[G]$ is free as a $\mathbb{Z}[H]$ -module,

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A).$$

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \mathrm{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Proof. Fix a projective $\mathbb{Z}[G]$ -resolution P_\bullet of \mathbb{Z} . Since $\mathbb{Z}[G]$ is free as a $\mathbb{Z}[H]$ -module, P_\bullet is a projective $\mathbb{Z}[H]$ -resolution of \mathbb{Z} .

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A).$$

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \mathrm{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Proof. Fix a projective $\mathbb{Z}[G]$ -resolution P_\bullet of \mathbb{Z} . Since $\mathbb{Z}[G]$ is free as a $\mathbb{Z}[H]$ -module, P_\bullet is a projective $\mathbb{Z}[H]$ -resolution of \mathbb{Z} . The book gives isomorphisms

$$\mathrm{Hom}_{\mathbb{Z}[G]}(P_i, \mathrm{CoInd}_H^G(A)) \cong \mathrm{Hom}_{\mathbb{Z}[H]}(P_i, A).$$

Shapiro's lemma

Let $H < G$ and A be a H -module. Form the **co-induced module**

$$\mathrm{CoInd}_H^G(A) := \mathrm{Hom}_H(\mathbb{Z}[G], A).$$

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \mathrm{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Proof. Fix a projective $\mathbb{Z}[G]$ -resolution P_\bullet of \mathbb{Z} . Since $\mathbb{Z}[G]$ is free as a $\mathbb{Z}[H]$ -module, P_\bullet is a projective $\mathbb{Z}[H]$ -resolution of \mathbb{Z} . The book gives isomorphisms

$$\mathrm{Hom}_{\mathbb{Z}[G]}(P_i, \mathrm{CoInd}_H^G(A)) \cong \mathrm{Hom}_{\mathbb{Z}[H]}(P_i, A).$$

Taking cohomology yields the lemma.

Shapiro's lemma

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \operatorname{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Shapiro's lemma

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \operatorname{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Let $\operatorname{CoInd}^G(A) := \operatorname{CoInd}_H^G(A)$ if $H = *$.

Shapiro's lemma

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \operatorname{CoInd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Let $\operatorname{CoInd}^G(A) := \operatorname{CoInd}_H^G(A)$ if $H = *$.

Since $H^i(*, A) = 0$ for $i > 0$, we have:

Shapiro's lemma

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \operatorname{Colnd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Let $\operatorname{Colnd}^G(A) := \operatorname{Colnd}_H^G(A)$ if $H = *$.

Since $H^i(*, A) = 0$ for $i > 0$, we have:

Corollary. $H^i(G, \operatorname{Colnd}^G(A)) = 0$ for $i > 0$.

Shapiro's lemma

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \operatorname{Colnd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Let $\operatorname{Colnd}^G(A) := \operatorname{Colnd}_H^G(A)$ if $H = *$.

Since $H^i(*, A) = 0$ for $i > 0$, we have:

Corollary. $H^i(G, \operatorname{Colnd}^G(A)) = 0$ for $i > 0$.

If $H < G$, $\operatorname{Colnd}^G(A) = \bigoplus \operatorname{Colnd}^H(A)$

Shapiro's lemma

Lemma. (Shapiro) Given subgroup H of a group G and an H -module A , there are canonical isomorphisms

$$H^i(G, \operatorname{Colnd}_H^G(A)) \cong H^i(H, A) \quad \text{for all } i \geq 0.$$

Let $\operatorname{Colnd}^G(A) := \operatorname{Colnd}_H^G(A)$ if $H = *$.

Since $H^i(*, A) = 0$ for $i > 0$, we have:

Corollary. $H^i(G, \operatorname{Colnd}^G(A)) = 0$ for $i > 0$.

If $H < G$, $\operatorname{Colnd}^G(A) = \bigoplus \operatorname{Colnd}^H(A)$ and so

$$H^i(H, \operatorname{Colnd}^G(A)) = \bigoplus H^i(H, \operatorname{Colnd}^H(A)) = 0.$$

The maps Res, Cor, Inf

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map is the inclusion $A^G \hookrightarrow A^H$.

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map is the inclusion $A^G \hookrightarrow A^H$.

$i \Rightarrow i + 1$. We do the case $i \geq 1$. The case $i = 0$ is similar.

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map is the inclusion $A^G \hookrightarrow A^H$.

$i \Rightarrow i + 1$. We do the case $i \geq 1$. The case $i = 0$ is similar.

The SES

$$0 \rightarrow A \rightarrow \text{CoInd}^G(A) \rightarrow A' \rightarrow 0$$

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map is the inclusion $A^G \hookrightarrow A^H$.

$i \Rightarrow i + 1$. We do the case $i \geq 1$. The case $i = 0$ is similar.

The SES

$$0 \rightarrow A \rightarrow \text{Colnd}^G(A) \rightarrow A' \rightarrow 0$$

induces the LES

$$H^i(G, \text{Colnd}^G(A)) \rightarrow H^i(G, A') \rightarrow H^{i+1}(G, A) \rightarrow H^{i+1}(G, \text{Colnd}^G(A)).$$

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map is the inclusion $A^G \hookrightarrow A^H$.

$i \Rightarrow i + 1$. We do the case $i \geq 1$. The case $i = 0$ is similar.

The SES

$$0 \rightarrow A \rightarrow \text{CoInd}^G(A) \rightarrow A' \rightarrow 0$$

induces the LES

$$H^i(G, \text{CoInd}^G(A)) \rightarrow H^i(G, A') \rightarrow H^{i+1}(G, A) \rightarrow H^{i+1}(G, \text{CoInd}^G(A)).$$

Since $i > 0$, we can use Shapiro's lemma.

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map is the inclusion $A^G \hookrightarrow A^H$.

$i \Rightarrow i + 1$. We do the case $i \geq 1$. The case $i = 0$ is similar.

The SES

$$0 \rightarrow A \rightarrow \text{CoInd}^G(A) \rightarrow A' \rightarrow 0$$

induces the LES

$$0 \rightarrow H^i(G, A') \rightarrow H^{i+1}(G, A) \rightarrow 0.$$

Since $i > 0$, we can use Shapiro's lemma.

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map is the inclusion $A^G \hookrightarrow A^H$.

$i \Rightarrow i + 1$. We do the case $i \geq 1$. The case $i = 0$ is similar.

The SES

$$0 \rightarrow A \rightarrow \text{CoInd}^G(A) \rightarrow A' \rightarrow 0$$

induces

$$H^i(G, A') \cong H^{i+1}(G, A).$$

Since $i > 0$, we can use Shapiro's lemma.

Restriction maps

G group, $H < G$, A a G -module.

We define maps $\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map is the inclusion $A^G \hookrightarrow A^H$.

$i \Rightarrow i + 1$. We do the case $i \geq 1$. The case $i = 0$ is similar.

The SES

$$0 \rightarrow A \rightarrow \text{CoInd}^G(A) \rightarrow A' \rightarrow 0$$

induces

$$H^i(G, A') \cong H^{i+1}(G, A).$$

Define $\text{Res} : H^{i+1}(G, A) \rightarrow H^{i+1}(H, A)$ as

$\text{Res} : H^i(G, A') \rightarrow H^i(H, A')$.

Restriction maps

Explicitly,

$$\text{Res} : H^i(G, A) \rightarrow H^i(H, A)$$

takes a cochain

$$f : G^i \rightarrow A$$

to the the same function with its domain restricted to H^i :

$$\text{Res}(f) : H^i \rightarrow A.$$

Corestriction maps

G group, H subgroup of finite index in G , A a G -module.

We define maps $\text{Cor} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

Corestriction maps

G group, H subgroup of finite index in G , A a G -module.

We define maps $\text{Cor} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map “averages over the coset representatives g_1, \dots, g_n of G/H ”:

$$A^H \rightarrow A^G : a \mapsto \sum_{j=1}^n g_j \cdot a$$

Corestriction maps

G group, H subgroup of finite index in G , A a G -module.

We define maps $\text{Cor} : H^i(G, A) \rightarrow H^i(H, A)$ inductively on i .

$i = 0$. The map “averages over the coset representatives g_1, \dots, g_n of G/H ”:

$$A^H \rightarrow A^G : a \mapsto \sum_{j=1}^n g_j \cdot a$$

$i \Rightarrow i + 1$. Do the same “dimension shifting” argument using Shapiro’s lemma.

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Proof.

$$H^i(G, A) \xrightarrow{\text{Res}} H^i(H, A) \xrightarrow{\text{Cor}} H^i(G, A)$$

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Proof. By dimension shifting, it suffices to prove the case $i = 0$.

$$H^i(G, A) \xrightarrow{\text{Res}} H^i(H, A) \xrightarrow{\text{Cor}} H^i(G, A)$$

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Proof. By dimension shifting, it suffices to prove the case $i = 0$.

$$H^0(G, A) \xrightarrow{\text{Res}} H^0(H, A) \xrightarrow{\text{Cor}} H^0(G, A)$$

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Proof. By dimension shifting, it suffices to prove the case $i = 0$.

$$A^G \xrightarrow{\text{Res}} A^H \xrightarrow{\text{Cor}} A^G$$

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Proof. By dimension shifting, it suffices to prove the case $i = 0$.

$$A^G \xrightarrow{\text{Res}} A^H \xrightarrow{\text{Cor}} A^G$$

$$a \longmapsto a$$

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Proof. By dimension shifting, it suffices to prove the case $i = 0$.

$$A^G \xrightarrow{\text{Res}} A^H \xrightarrow{\text{Cor}} A^G$$

$$a \longmapsto a \longmapsto \sum_{j=1}^n g_j \cdot a$$

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Proof. By dimension shifting, it suffices to prove the case $i = 0$.

$$A^G \xrightarrow{\text{Res}} A^H \xrightarrow{\text{Cor}} A^G$$

$$a \longmapsto a \longmapsto \sum_{j=1}^n a$$

Restriction-Corestriction

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Proof. By dimension shifting, it suffices to prove the case $i = 0$.

$$A^G \xrightarrow{\text{Res}} A^H \xrightarrow{\text{Cor}} A^G$$

$$a \longmapsto a \longmapsto n \cdot a$$

Inflation maps

For a group G , normal subgroup H , and G -module A , we define maps

$$\text{Inf} : H^i(G/H, A^H) \rightarrow H^i(G, A)$$

on degree zero and use dimension shifting to extend the definition to all $i \geq 0$.

Inflation maps

For a group G , normal subgroup H , and G -module A , we define maps

$$\text{Inf} : H^i(G/H, A^H) \rightarrow H^i(G, A)$$

on degree zero and use dimension shifting to extend the definition to all $i \geq 0$.

Explicitly, Inf takes a cochain

$$f : (G/H)^i \rightarrow A^H$$

to the cochain

$$G^i \rightarrow (G/H)^i \xrightarrow{f} A^H \rightarrow A$$

Inflation-Restriction

Theorem B. Let G be a group, A a G -module, and H a normal subgroup of G . There exists a map

$$\tau : H^1(H, A)^{G/H} \rightarrow H^2(G/H, A^H),$$

called the **transgression map**, fitting into an exact sequence

$$\begin{aligned} 0 \longrightarrow H^1(G/H, A^H) &\xrightarrow{\text{Inf}} H^1(G, A) \xrightarrow{\text{Res}} H^1(G/H, A)^{G/H} \\ &\xrightarrow{\tau} H^2(G/H, A^H) \xrightarrow{\text{Inf}} H^2(G, A) \end{aligned}$$

Proof. Consequence of the Hochschild-Serre spectral sequence.

Inflation-Restriction

Proposition 3.3.17. In Theorem B, let $i > 1$ and assume $H^j(H, A) = 0$ for $1 \leq j \leq i - 1$. Then there is a natural map

$$\tau_{i,A} : H^i(H, A)^{G/H} \rightarrow H^{i+1}(G/H, A^H)$$

fitting into an exact sequence

$$\begin{aligned} 0 \longrightarrow H^i(G/H, A^H) &\xrightarrow{\text{Inf}} H^i(G, A) \xrightarrow{\text{Res}} H^i(G/H, A)^{G/H} \\ &\xrightarrow{\tau} H^{i+1}(G/H, A^H) \xrightarrow{\text{Inf}} H^{i+1}(G, A) \end{aligned}$$

Proof. Follows from Theorem B and dimension shifting.

References

Section 3.3 from Gille-Szamuely, Central Simple Algebras and Galois Cohomology.

Sharifi's notes on group and Galois cohomology,
<https://www.math.ucla.edu/~sharifi/groupcoh.pdf>

For spectral sequences, see e.g. Vakil's notes on algebraic geometry, section 1.7, <http://math.stanford.edu/~vakil/216blog/FOAGnov1817public.pdf>