

3.3 Relation to subgroups

November 1, 2021

Introduction

Why use cohomology groups?

They are invariants that tell things apart.

Why use cohomology groups?

They are invariants that tell things apart.

De Rham cohomology

$H_{dR}^i(X)$ is a real vector space.

↖ Smooth manifold

Why use cohomology groups?

They are invariants that tell things apart.

De Rham cohomology

$H_{dR}^i(X)$ is a real vector space.

This invariant distinguishes spheres up to homotopy equivalence:

Why use cohomology groups?

They are invariants that tell things apart.

De Rham cohomology

$H_{dR}^i(X)$ is a real vector space.

This invariant distinguishes spheres up to homotopy equivalence:

For $n > m \geq 0$, we have $S^n \not\cong S^m$ since

$$H^n(S^n) \cong \mathbb{Z} \quad \text{but} \quad H^n(S^m) \cong 0.$$

Why use cohomology groups?

They also help classify things.

Nonabelian group cohomology (chapter 2)

$H^0(\overset{\text{group}}{\downarrow} G, A) = A^G$ is a group. (nonabelian) group with compatible G-action

$H^1(G, A)$ is a pointed set.

H^1 classifies twisted forms.

Why use cohomology groups?

They also help classify things.

Nonabelian group cohomology (chapter 2)

$H^0(G, A) = A^G$ is a group.

$H^1(G, A)$ is a pointed set.

H^1 classifies twisted forms.

Group cohomology (chapter 3)

group ↘ ↙ abelian group with compatible G -action

$H^i(G, A)$ for $i \geq 0$.

$H^2(G, A)$ classifies group extensions of G by A

0-class corresponds to split extension (the semidirect product $A \rtimes G$ with the given G -action on A)

How to compute cohomology groups?

De Rham cohomology $H^\bullet(X)$

How to compute cohomology groups?

De Rham cohomology $H^\bullet(X)$

Break the manifold X up into subspaces U and V

How to compute cohomology groups?

De Rham cohomology $H^\bullet(X)$

Break the manifold X up into subspaces U and V and relate $H^\bullet(U)$ and $H^\bullet(V)$ to $H^\bullet(X)$. (Mayer-Vietoris sequence)

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

► $H < G$

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

► $H < G \Rightarrow H$ acts on A

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

► $H < G \Rightarrow H$ acts on $A \Rightarrow$ get $H^\bullet(H, A)$.

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

- ▶ $H < G \Rightarrow H$ acts on $A \Rightarrow$ get $H^\bullet(H, A)$.
- ▶ $H \triangleleft G$

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

- ▶ $H < G \Rightarrow H$ acts on $A \Rightarrow$ get $H^\bullet(H, A)$.
- ▶ $H \triangleleft G \not\Rightarrow G/H$ acts on A (since H -action on A may be nontrivial)

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

- ▶ $H < G \Rightarrow H$ acts on $A \Rightarrow$ get $H^\bullet(H, A)$.
- ▶ $H \triangleleft G \not\Rightarrow G/H$ acts on A (since H -action on A may be nontrivial)
- ▶ $H \triangleleft G \Rightarrow G/H$ acts on A^H

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

- ▶ $H < G \Rightarrow H$ acts on $A \Rightarrow$ get $H^\bullet(H, A)$.
- ▶ $H \triangleleft G \not\Rightarrow G/H$ acts on A (since H -action on A may be nontrivial)
- ▶ $H \triangleleft G \Rightarrow G/H$ acts on $A^H \Rightarrow$ get $H^\bullet(G/H, A^H)$.

How to compute cohomology groups?

Group cohomology $H^i(G, A)$

Break G -module A up into a short exact sequence

$$0 \rightarrow A' \rightarrow A \rightarrow A'' \rightarrow 0.$$

This induces a long exact sequence in cohomology.

Or break G up.

- ▶ $H < G \Rightarrow H$ acts on $A \Rightarrow$ get $H^\bullet(H, A)$.
- ▶ $H \triangleleft G \not\Rightarrow G/H$ acts on A (since H -action on A may be nontrivial)
- ▶ $H \triangleleft G \Rightarrow G/H$ acts on $A^H \Rightarrow$ get $H^\bullet(G/H, A^H)$.

How can we relate $H^\bullet(H, A)$ and $H^\bullet(G/H, A^H)$ to $H^\bullet(G, A)$?

Main results for section

Relating $H^\bullet(H, A)$, $H^\bullet(G/H, A^H)$, $H^\bullet(G, A)$

Main results for section

Relating $H^\bullet(H, A)$, $H^\bullet(G/H, A^H)$, $H^\bullet(G, A)$

Let G be a group, A a G -module, and H a subgroup of G . Then there exist group homomorphisms, called **restriction maps**,

$$\text{Res} : H^i(G, A) \rightarrow H^i(H, A) \quad \text{for all } i \geq 0.$$

Main results for section

Relating $H^\bullet(H, A)$, $H^\bullet(G/H, A^H)$, $H^\bullet(G, A)$

Let G be a group, A a G -module, and H a subgroup of G . Then there exist group homomorphisms, called **restriction maps**,

$$\text{Res} : H^i(G, A) \rightarrow H^i(H, A) \quad \text{for all } i \geq 0.$$

If $[G : H] = n$ is finite, there exist group homomorphisms, called **corestriction maps**,

$$\text{Cor} : H^i(H, A) \rightarrow H^i(G, A) \quad \text{for all } i \geq 0.$$

Main results for section

Relating $H^\bullet(H, A)$, $H^\bullet(G/H, A^H)$, $H^\bullet(G, A)$

Let G be a group, A a G -module, and H a subgroup of G . Then there exist group homomorphisms, called **restriction maps**,

$$\text{Res} : H^i(G, A) \rightarrow H^i(H, A) \quad \text{for all } i \geq 0.$$

If $[G : H] = n$ is finite, there exist group homomorphisms, called **corestriction maps**,

$$\text{Cor} : H^i(H, A) \rightarrow H^i(G, A) \quad \text{for all } i \geq 0.$$

If instead H is a normal subgroup of G , there exist group homomorphisms, called **inflation maps**,

$$\text{Inf} : H^i(G/H, A) \rightarrow H^i(G, A) \quad \text{for all } i \geq 0.$$

Main results for section

Relating Res, Cor, Inf

Main results for section

Relating Res, Cor, Inf

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Main results for section

Theorem B. Let G be a group, A a G -module, and H a normal subgroup of G . There exists a map

$$\tau : H^1(H, A)^{G/H} \rightarrow H^2(G/H, A^H),$$

called the **transgression map**, fitting into an exact sequence

$$\begin{aligned} 0 \longrightarrow H^1(G/H, A^H) &\xrightarrow{\text{Inf}} H^1(G, A) \xrightarrow{\text{Res}} H^1(G/H, A)^{G/H} \\ &\xrightarrow{\tau} H^2(G/H, A^H) \xrightarrow{\text{Inf}} H^2(G, A) \end{aligned}$$

Application

Assume Theorem A holds:

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Assume Theorem A holds:

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Assume Theorem A holds:

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Proof.

$$\begin{array}{ccccc} H^i(G, A) & \xrightarrow{\text{Res}} & H^i(H, A) & \xrightarrow{\text{Cor}} & H^i(G, A). \\ & & \searrow & \nearrow & \\ & & \times [G:H] & & \end{array}$$

Assume Theorem A holds:

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Proof. Let $H = *$ in Theorem A and note $H^i(*, A) = 0$ for $i > 0$:

$$\begin{array}{ccccc} H^i(G, A) & \xrightarrow{\text{Res}} & H^i(H, A) & \xrightarrow{\text{Cor}} & H^i(G, A). \\ & & \searrow & \nearrow & \\ & & \times [G:H] & & \end{array}$$

Assume Theorem A holds:

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Proof. Let $H = *$ in Theorem A and note $H^i(*, A) = 0$ for $i > 0$:

$$\begin{array}{ccccc} H^i(G, A) & \xrightarrow{\text{Res}} & H^i(*, A) & \xrightarrow{\text{Cor}} & H^i(G, A) \\ & & \searrow \text{ } \times n \nearrow & & \end{array}$$

Assume Theorem A holds:

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Proof. Let $H = *$ in Theorem A and note $H^i(*, A) = 0$ for $i > 0$:

$$H^i(G, A) \xrightarrow{\text{Res}} 0 \xrightarrow{\text{Cor}} H^i(G, A).$$

$\searrow \times n \nearrow$

Assume Theorem A holds:

Theorem A. Let G be a group, A a G -module, and H a subgroup of finite index n in G . Then

$$\text{Cor} \circ \text{Res} : H^i(G, A) \rightarrow H^i(G, A)$$

is given by multiplication by n for all $i \geq 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Proof. Let $H = *$ in Theorem A and note $H^i(*, A) = 0$ for $i > 0$:

$$H^i(G, A) \xrightarrow{\text{Res}} 0 \xrightarrow{\text{Cor}} H^i(G, A).$$

$\searrow \times n$

So $\times n$ map is the $\times 0$ map.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Corollary. (Schur-Zassenhaus theorem, special case)

Let G be a group and A an abelian group. If $|G|$ and $|A|$ are finite and coprime, then any group extension of G by A is the semidirect product $A \rtimes G$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Corollary. (Schur-Zassenhaus theorem, special case)

Let G be a group and A an abelian group. If $|G|$ and $|A|$ are finite and coprime, then any group extension of G by A is the semidirect product $A \rtimes G$.

Proof. An extension of G by A gives a G -module structure on A and an element of $H^2(G, A)$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Corollary. (Schur-Zassenhaus theorem, special case)

Let G be a group and A an abelian group. If $|G|$ and $|A|$ are finite and coprime, then any group extension of G by A is the semidirect product $A \rtimes G$.

Proof. An extension of G by A gives a G -module structure on A and an element of $H^2(G, A)$. By earlier discussion, it suffices to show that $H^2(G, A) = 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Corollary. (Schur-Zassenhaus theorem, special case)

Let G be a group and A an abelian group. If $|G|$ and $|A|$ are finite and coprime, then any group extension of G by A is the semidirect product $A \rtimes G$.

Proof. An extension of G by A gives a G -module structure on A and an element of $H^2(G, A)$. By earlier discussion, it suffices to show that $H^2(G, A) = 0$.

By previous corollary, $|G| \cdot H^2(G, A) = 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Corollary. (Schur-Zassenhaus theorem, special case)

Let G be a group and A an abelian group. If $|G|$ and $|A|$ are finite and coprime, then any group extension of G by A is the semidirect product $A \rtimes G$.

Proof. An extension of G by A gives a G -module structure on A and an element of $H^2(G, A)$. By earlier discussion, it suffices to show that $H^2(G, A) = 0$.

By previous corollary, $|G| \cdot H^2(G, A) = 0$.

Also $|A| \cdot H^2(G, A) = 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Corollary. (Schur-Zassenhaus theorem, special case)

Let G be a group and A an abelian group. If $|G|$ and $|A|$ are finite and coprime, then any group extension of G by A is the semidirect product $A \rtimes G$.

Proof. An extension of G by A gives a G -module structure on A and an element of $H^2(G, A)$. By earlier discussion, it suffices to show that $H^2(G, A) = 0$.

By previous corollary, $|G| \cdot H^2(G, A) = 0$.

Also $|A| \cdot H^2(G, A) = 0$.

So $\gcd(|G|, |A|) \cdot H^2(G, A) = 0$.

Corollary. If G is a finite group of order n , then $n \cdot H^i(G, A) = 0$ for any G -module A and $i > 0$.

Corollary. (Schur-Zassenhaus theorem, special case)

Let G be a group and A an abelian group. If $|G|$ and $|A|$ are finite and coprime, then any group extension of G by A is the semidirect product $A \rtimes G$.

Proof. An extension of G by A gives a G -module structure on A and an element of $H^2(G, A)$. By earlier discussion, it suffices to show that $H^2(G, A) = 0$.

By previous corollary, $|G| \cdot H^2(G, A) = 0$.

Also $|A| \cdot H^2(G, A) = 0$.

So $1 \cdot H^2(G, A) = 0$.