
⚠ 1. HEART ATTACK (Myocardial Infarction)

Instructions & First Aid

- Make the person **sit upright** and keep them calm.
 - **Call emergency services immediately.**
 - Give **Aspirin 300 mg** to chew (ONLY if the person is conscious & not allergic).
 - **Loosen tight clothes**; ensure fresh air.
 - If the person becomes unresponsive → **start CPR** if trained.
 - **Do NOT** allow walking, running, or food/water.
-

⚠ 2. STROKE (Use FAST)

FAST Check

- **F – Face:** Smile → look for drooping.
- **A – Arms:** Raise both arms → one drops.
- **S – Speech:** Slurred/strange words.
- **T – Time:** Reach hospital **immediately**.

First Aid

- Lay them on their **side** (recovery position) if vomiting.
 - Keep the head raised slightly.
 - **Do NOT** give food, water, or medicine.
 - Note the **time symptoms started**.
-

⚠ 3. SEVERE BREATHING DIFFICULTY

First Aid

- Sit the person upright; **do NOT lie them down**.
- Loosen tight clothes.
- Provide **fresh air** / open windows.
- If they have an inhaler → help them use it.

- Call **emergency help immediately**.
 - If lips turn blue → **critical emergency**.
-

4. LOSS OF CONSCIOUSNESS / SUDDEN CONFUSION

First Aid

- Check **response** by calling their name.
 - If unresponsive but breathing → place in **recovery position**.
 - If not breathing → **start CPR**.
 - Keep airway open; do NOT give food/water.
 - Call for help immediately.
-

5. SERIOUS FALLS & FRACTURES

First Aid

- Do **not** move the injured area.
 - Support the limb with cloth/soft padding.
 - Apply **cold pack** for swelling.
 - If hip fracture suspected → keep person still.
 - For head injury with vomiting/confusion → **hospital immediately**.
-

6. SEVERE BLEEDING

First Aid

- Apply **firm direct pressure** with clean cloth.
 - Raise the bleeding part above heart level (if possible).
 - Do NOT remove soaked cloth → add another on top.
 - If bleeding does not stop → emergency care.
-

7. SEPSIS / SEVERE INFECTION

First Aid

- Check for fever, chills, confusion.
 - Keep the person hydrated with sips (if conscious).
 - Lay them down; keep warm.
 - Seek emergency care **immediately** (life-threatening).
-

8. SEVERE ABDOMINAL PAIN

First Aid

- Do NOT give painkillers unless advised (may hide symptoms).
 - No food or drink until evaluated.
 - Make the person lie comfortably with knees bent.
 - Seek urgent medical help if pain lasts >30 min or worsens.
-

9. UNCONTROLLED DIABETES

Low Blood Sugar (Hypoglycemia)

- If conscious → give **sugar**, glucose water, juice.
- If unconscious → **do NOT give anything by mouth**.
- Seek help urgently.

High Blood Sugar (Hyperglycemia)

- Encourage water (if conscious).
 - Check sugar if possible.
 - Seek immediate medical care if confused or breathing fast.
-

10. ADVERSE DRUG REACTION / ALLERGY

First Aid

- Look for swelling of face, lips, throat, breathing trouble.
- Stop the medicine immediately.
- If breathing difficulty → **medical emergency**.
- Use **EpiPen** if available (rare in rural homes).

11. SEVERE DEHYDRATION

First Aid

- Give **ORS** or water slowly (if conscious).
- Lay flat and raise legs slightly.
- Check urine colour (dark = severe).
- Seek help if dizziness/confusion persists.