

**Revêtements Étales and the Fundamental Group
(SGA I)**

Alexander Grothendieck

Contents

Translation Notes	5
Chapter 1. Étale morphisms	7
1. Basics of differential calculus	7
2. Quasi-finite morphisms	7
3. Unramified morphisms	8

Translation Notes

The original can be found on the arXiv. I don't know French, I just used Google translate and some knowledge of the subject matter.

CHAPTER 1

Étale morphisms

To simplify the exposition we assume that all preschemes in the following are locally Noetherian (at least, starting from section 2).

1. Basics of differential calculus

Let X be a prescheme on Y , and $\Delta_{X/Y}$ the diagonal morphism $X \rightarrow X \times_Y X$. This is an immersion, and thus a closed immersion of X into an open subset V of $X \times_Y X$. Let \mathcal{I}_X be the ideal of the closed sub-prescheme corresponding to the diagonal in V (N.B. if one really wishes to do things intrinsically, without assuming that X is separated over Y — a **misleading** hypothesis — then one should consider the set-theoretic inverse image of $\mathcal{O}_{X \times X}$ in X and denote by \mathcal{I}_X the augmentation ideal in the above...). The sheaf $\mathcal{I}_X/\mathcal{I}_X^2$ can be thought of as a quasi-coherent sheaf on X , which we denote by $\Omega_{X/Y}^1$. It is of finite type if $X \rightarrow Y$ is of finite type. It behaves well with respect to a change of base $Y' \rightarrow Y$. We also introduce the sheaves $\mathcal{O}_{X \times_Y X}/\mathcal{I}_X^{n+1} = \mathcal{P}_{X/Y}^n$, which are sheaves of *rings* on X , giving us preschemes denoted by $\Delta_{X/Y}^n$ and called the *n-th infinitesimal neighbourhood of X/Y* . The polysyllogism is entirely trivial, even if rather long¹; it seems wise to not discuss it until we use it for something helpful, with smooth morphisms.

2. Quasi-finite morphisms

PROPOSITION 2.1. *Let $A \rightarrow B$ be a local homomorphism (N.B. all rings are now Noetherian) and \mathfrak{m} the maximal ideal of A . Then the following conditions are equivalent:*

- (i) $B/\mathfrak{m}B$ is of finite dimension over $k = A/\mathfrak{m}$.
- (ii) $\mathfrak{m}B$ is **[...]** and $B/\mathfrak{r}(B) = k(B)$ is an extension of $k = k(A)$.
- (iii) The completion \hat{B} of B is finite over the completion \hat{A} of A .

We then say that B is *quasi-finite* over A . A morphism $f: X \rightarrow Y$ is said to be quasi-finite in x (or the Y -prescheme f is said to be quasi-finite in x) if \mathcal{O}_x is quasi-finite over $\mathcal{O}_{f(x)}$. **[...]** A morphism is said to be quasi-finite if it is quasi-finite in each point².

COROLLARY 2.2. *If A is complete then quasi-finiteness is equivalent to finiteness.*

We could give the usual polysyllogism (i), (ii), (iii), (iv), (v) for quasi-finite morphisms, but that doesn't seem necessary here.

¹cf. EGA IV 16.3.

²In EGA II 6.2.3 we further suppose that f is of finite type.

3. Unramified morphisms

PROPOSITION 3.1. *Let $f: X \rightarrow Y$ be a morphism of finite type, $x \in X$, and $y = f(x)$. Then the following conditions are equivalent:*

- (i) $\mathcal{O}_x/\mathfrak{m}_y\mathcal{O}_x$ is a finite separable extension of $k(y)$.
- (ii) $\Omega_{X/Y}^1$ is **null** in x .
- (iii) The diagonal morphism $\Delta_{X/Y}$ is an open immersion on a neighbourhood of x .

PROOF. For the implication (i) \implies (ii), we are brought by Nakayama to the case where $Y = \operatorname{Spec}(k)$ and $X = \operatorname{Spec}(k')$, where it is well known and otherwise trivial by the definition of separable; (ii) \implies (iii) comes from a nice and easy characterisation of open immersions, using Krull; (iii) \implies (i) follows as well from reducing to the case where $Y = \operatorname{Spec}(k)$ and the diagonal morphism is everywhere an open immersion. One must then prove that X is finite **[...]** and this leads us to consider the case where k is algebraically closed. But then every closed point of X is isolated (since it is identical to the inverse image of the diagonal by the morphism $X \rightarrow X \times_k X$ defined by x), whence X is finite. We can thus suppose that X reduces to a point, of the ring A , and so $A \otimes_k A \rightarrow A$ is an isomorphism, hence $A = k$. \square

DEFINITION 3.2. (a) *We then say that f is unramified in x , or that X is unramified in x on Y .*

- (b) *Let $A \rightarrow B$ be a local homomorphism. We say that it is unramified, or that B is a local unramified algebra on A , if $B/\mathfrak{m}_A B$ is a finite separable extension of A/\mathfrak{m}_A , i.e. if $\mathfrak{m}_A B = \mathfrak{m}_B$ and $k(B)$ is a separable extension of $k(A)$ ³.*

REMARK. The fact that B is unramified over A **[...]**. Unramified implies quasi-finite.

COROLLARY 3.3. *The set of points where f is unramified is open.*

³Cf. regrets in III 1.2.