

Please stick the barcode label here.

Candidate Number

HONG KONG DIPLOMA OF SECONDARY
EDUCATION EXAMINATION MOCK

MATHEMATICS Extended Part

Module 2 (Algebra and Calculus)

Question-Answer Book

Time allowed : 2 $\frac{1}{2}$ hours

This paper must be answered in English

INSTRUCTIONS

- (1) After the announcement of the start of the examination, you should first write your Candidate Number in the space provided on Page 1 and stick barcode labels in the spaces provided on Pages 1.
- (2) This paper consists of TWO sections, A and B.
- (3) This paper carried 100 marks.
- (4) Attempt ALL questions in this paper. Write your answers in the spaces provided in this Question-Answer Book. Do not write in the margins. Answers written in the margins will not be marked.
- (5) Graph paper and supplementary answer sheets will be supplied on request. Write your Candidate Number, mark the question number box and stick a barcode label on each sheet, and fasten them with string INSIDE this book.
- (6) Unless otherwise specified, all working must be clearly shown.
- (7) Unless otherwise specified, numerical answers must be exact.
- (8) The diagrams in this paper are not necessarily drawn to scale.
- (9) No extra time will be given to candidates for sticking on the barcode labels or filling in the question number boxes after the 'Time is up' announcement.

FORMULAS FOR REFERENCE

$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$	$\sin A + \sin B = 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2}$
$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$	$\sin A - \sin B = 2 \cos \frac{A+B}{2} \sin \frac{A-B}{2}$
$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$	$\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2}$
$2 \sin A \cos B = \sin(A+B) + \sin(A-B)$	$\cos A - \cos B = -2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}$
$2 \cos A \cos B = \cos(A+B) + \cos(A-B)$	
$2 \sin A \sin B = \cos(A-B) - \cos(A+B)$	

SECTION A (50 marks)

1. Let $f(x) = \sin^2 x + x \cos x$ for all real numbers x .

(a) Express $f\left(\frac{\pi}{2} + h\right) - f\left(\frac{\pi}{2}\right)$ in terms of h . (2 marks)

(b) Define $g(x) = e^{f(x)}$ for all real numbers x . Using first principles, find $g'\left(\frac{\pi}{2}\right)$. (4 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

2. (a) Prove that $\frac{\cos x + \cos 2x - \cos 3x - \cos 4x}{\sin x + \sin 2x + \sin 3x + \sin 4x} = \tan x$, where $\sin x + \sin 2x + \sin 3x + \sin 4x \neq 0$. (2 marks)
- (b) Prove that $\frac{\sin x + \sin 2x + \sin 3x + \sin 4x}{\cos x + \cos 2x + \cos 3x + \cos 4x} = \tan \frac{5x}{2}$, where $\cos x + \cos 2x + \cos 3x + \cos 4x \neq 0$. (1 mark)
- (c) Solve $\frac{\cos \theta + \cos 2\theta - \cos 3\theta - \cos 4\theta}{\sin \theta + \sin 2\theta + \sin 3\theta + \sin 4\theta} = \frac{\sin \theta + \sin 2\theta + \sin 3\theta + \sin 4\theta}{\cos \theta + \cos 2\theta + \cos 3\theta + \cos 4\theta}$ for $-\pi \leq \theta \leq \pi$. (3 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

3. (a) Using integration by parts, prove that

$$\int e^{-x}[f(x) - f'(x)] dx = -e^{-x}f(x) + C,$$

where C is a constant of integration.

(2 marks)

(b) Prove that $\frac{x^n + nx^{n-1} + x + 1}{(x + x^n)^2} = \frac{1}{x + x^n} - \frac{d}{dx} \left(\frac{1}{x + x^n} \right)$ for any positive integer $n \geq 2$.

(2 marks)

(c) Hence, evaluate $\int_5^8 e^{-x} \cdot \frac{x^3 + 3x^2 + x + 1}{(x + x^3)^2} dx$.

(2 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

(b) If $\|\vec{v}_2\| = \frac{1}{5}$ and $\theta = \frac{\pi}{6}$, find the distance between $\vec{v}_2$ and $\vec{v}_3$. (3 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

- DSE-MOCK-MATH-EP(M2)-9

- 9

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

(b) Hence, find the value of $\sum_{k=10}^{90} (5 + 4k) \cdot 3^{2k+1}$. (2 marks)

Answers written in the margins will not be marked.


Answers written in the margins will not be marked.

7. An inner bowl B_1 and an outer bowl B_2 are placed concentrically to form a double-walled container with a cavity in between.

The inner surface of B_1 and the outer surface of B_2 are generated by revolving the curves C_1 and C_2 about the y -axis respectively, where:

$$C_1 : y = x^2$$

$$C_2 : x^2 + (y - 1)^2 = \frac{3}{4} \quad \text{for } y \leq 1$$



It is given that C_1 and C_2 touch each other tangentially at $y = \frac{1}{2}$, sealing the top edge of the cavity.

- Find the total capacity of the cavity between the two bowls. (3 marks)
- Water is poured into the empty cavity between the bowls at a constant rate of $\frac{\pi}{1280}$ cubic units per second. Let h be the depth of water in the cavity at time t seconds (where $0 < h \leq \frac{1}{2}$).
 - Express the volume of water $V(h)$ in the cavity in terms of h . (2 marks)
 - Find the rate of change of the water depth $\frac{dh}{dt}$ when the water depth is $h = \frac{1}{4}$. (2 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

8. (a) Given that $\begin{vmatrix} -2 & 2 & 2 - \lambda_2 \\ -\lambda_1 & 1 & 1 + \lambda_1 \\ 3 & -1 & \lambda_2 - 1 \end{vmatrix} = 0$, where λ_1 and λ_2 are non-negative integers. Find the value of $\lambda_1 + \lambda_2$. (3 marks)

- (b) Consider the system of linear equations:

$$\begin{cases} 3x + 2y + (2 - \lambda_2)z = k \\ \lambda_1 x + y + (\lambda_1 + 1)z = k^2 \\ -2x - y + (\lambda_2 - 1)z = 1 \end{cases}$$

Find the values of k such that the system of linear equations has infinitely many solutions. (2 marks)

- (c) Hence, solve the system of linear equations for each value of k obtained in (b). (2 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There is a faint, repeating watermark diagonally across the page that reads "chinwasiu".

This image shows a blank sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There is a faint, repeating watermark diagonally across the page that reads "chinwasiu".

SECTION B (50 marks)

9. Let $f(x) = \frac{x^3+x^2-5x+3}{x^2+3}$ for all real numbers x . Let C be the curve $y = f(x)$.

- (a) Find the asymptote(s) of C . (2 marks)
- (b) Find the point(s) of inflexion of C . (3 marks)
- (c) Sketch C . (3 marks)
- (d) (i) Find the equation of the tangent L to C at $x = -\sqrt{3}$. (2 marks)
- (ii) Prove that $L \geq f(x)$ for all real numbers x . (3 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

10. (a) (i) Using integration by parts, find $\int \ln x \, dx$. (2 marks)

(ii) Let $f(x)$ be a continuous function satisfying $f(-x) = f(x)$ for all real numbers x .
Prove that

$$\int_{-a}^a f(x) \, dx = 2 \int_0^a f(x) \, dx \quad (2 \text{ marks})$$

(b) Let $t > 1$. Define $I(t) = \int_{1/t}^t \frac{\ln\left(\frac{1+x^2}{x^2}\right)}{x(1+x^2)} \, dx$.

(i) Prove that $I(t) = \int_{1/t}^t \frac{x \ln(1+x^2)}{1+x^2} \, dx$. (2 marks)

(ii) Show that

$$I(t) = \int_0^{\ln t} \left(\ln(1+e^{2x}) - \frac{2x}{1+e^{2x}} \right) \, dx \quad (3 \text{ marks})$$

(iii) Using the results of (a)(i) and (b)(ii), evaluate $\int_{1/e}^e \frac{\ln\left(\frac{1+x^2}{x^2}\right)}{x(1+x^2)} \, dx$ (2 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

12. Let $\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3, \mathbf{v}_4$ be four position vectors in 3D space such that their lengths are all equal to 1, i.e., $|\mathbf{v}_1| = |\mathbf{v}_2| = |\mathbf{v}_3| = |\mathbf{v}_4| = 1$. The terminal points of these four vectors are V_1, V_2, V_3, V_4 respectively, forming a tetrahedron.

- (a) (i) Prove that $(\mathbf{v}_1 - \mathbf{v}_2) \times (\mathbf{v}_1 - \mathbf{v}_3) = \mathbf{v}_1 \times \mathbf{v}_2 + \mathbf{v}_2 \times \mathbf{v}_3 + \mathbf{v}_3 \times \mathbf{v}_1$. (2 marks)
- (ii) If $(\mathbf{v}_1 - \mathbf{v}_2) \times (\mathbf{v}_1 - \mathbf{v}_3) \neq (\mathbf{v}_1 - \mathbf{v}_2) \times (\mathbf{v}_1 - \mathbf{v}_4)$, explain why the four points V_1, V_2, V_3, V_4 are not coplanar. (1 mark)
- (b) Now assume $\mathbf{v}_1 = \mathbf{k}$, $\mathbf{v}_2 = \cos \theta \mathbf{i} + \sin \theta \mathbf{j}$, $\mathbf{v}_3 = -\cos \theta \mathbf{i} + \sin \theta \mathbf{j}$, and $\mathbf{v}_4 = -\mathbf{k}$, where $0 < \theta < \frac{\pi}{2}$.
- (i) Calculate $(\mathbf{v}_1 - \mathbf{v}_2) \times (\mathbf{v}_1 - \mathbf{v}_3)$ and $(\mathbf{v}_1 - \mathbf{v}_2) \times (\mathbf{v}_1 - \mathbf{v}_4)$, and verify that they are not equal. (2 marks)
- (ii) Find the volume V of the tetrahedron $V_1V_2V_3V_4$ in terms of θ . (2 marks)
- (c) As θ varies,
- (i) find the maximum volume of the tetrahedron $V_1V_2V_3V_4$. (2 marks)
- (ii) when the volume attains its maximum value, find the dihedral angle between the face $V_1V_2V_3$ and the base $V_2V_3V_4$. (3 marks)

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

Answers written in the margins will not be marked.

END OF PAPER