# Results


# **Descriptives**


Descriptives

	Dose	Happiness
N	Control 15 mins 30 mins	5 5 5
Missing	Control 15 mins 30 mins	0 0 0
Mean	Control 15 mins 30 mins	2.20 3.20 5.00
Median	Control 15 mins 30 mins	2.00 3.00 5.00
Standard deviation	Control 15 mins 30 mins	1.30 1.30 1.58
Minimum	Control 15 mins 30 mins	1.00 2.00 3.00
Maximum	Control 15 mins 30 mins	4.00 5.00 7.00
Skewness	Control 15 mins 30 mins	0.541 0.541 0.00
Std. error skewness	Control 15 mins 30 mins	0.913 0.913 0.913
Kurtosis	Control 15 mins 30 mins	-1.49 -1.49 -1.20
Std. error kurtosis	Control 15 mins 30 mins	2.00 2.00 2.00
Shapiro-Wilk W	Control 15 mins 30 mins	0.902 0.902 0.987
Shapiro-Wilk p	Control 15 mins 30 mins	0.421 0.421 0.967

### **Plots**

**Happiness** 






# Relationships, Prediction, and Group Comparisons

You have entered a numeric variable for Variable 1 / Dependent Variable and a nominal variable for Variable 2 / Independent Variables. Hence, a <u>one way ANOVA</u>, which is is a test for the difference between several population means, seems to be a good option for you! In order to run this analysis in jamovi, go to: ANOVA > ANOVA

• Drop your dependent (numeric) variable in the box below Dependent Variable and your independent (grouping) variable in the box below Fixed Factors

If the normality or homoscedasticity assumption is violated, you could use the non-parametric Kruskal-Wallis test. Click on the links to learn more about these tests!

Scatter Plots of Bivariate Relationships - Dependent/Independent Variables



## **One-Way ANOVA**

#### One-Way ANOVA

		F	df1	df2	р
Happiness	Welch's	4.32	2	7.94	0.054
	Fisher's	5.12	2	12	0.025

#### **Group Descriptives**

	Dose	N	Mean	SD	SE
Happiness	Control	5	2.20	1.30	0.583
	15 mins	5	3.20	1.30	0.583
	30 mins	5	5.00	1.58	0.707

### **Assumption Checks**

Normality Test (Shapiro-Wilk)

	W	р
Happiness	0.917	0.171

Note. A low p-value suggests a violation of the assumption of normality


#### Homogeneity of Variances Test (Levene's)

	F	df1	df2	р
Happiness	0.0917	2	12	0.913

[3]

#### **Plots**

## Mean (95% CI)



#### **Post Hoc Tests**

Tukey Post-Hoc Test - Happiness

		Control	15 mins	30 mins
Control	Mean difference p-value	_	-1.00 0.516	-2.80 0.021
15 mins	Mean difference p-value		_ _	-1.80 0.147
30 mins	Mean difference p-value			_ _

## **ANOVA**

ANOVA - Happiness

	Sum of Squares	df	Mean Square	F	р	$\omega^2$
Dose	20.1	2	10.07	5.12	0.025	0.354
Residuals	23.6	12	1.97			

[3]

## **Assumption Checks**


Homogeneity of Variances Test (Levene's)

F	df1	df2	р
0.0917	2	12	0.913

#### Normality Test (Shapiro-Wilk)


Statistic	р
0.917	0.171

### Q-Q Plot



# **Estimated Marginal Means**

#### Dose



#### Robust ANOVA

	F	р
Dose	3.00	0.160

Note. Method of trimmed means, trim level 0.2

#### **Post Hoc Tests**

Post Hoc Tests - Dose

				95% Confidence interva	
		psi-hat	р	Lower	Upper
Control	15 mins	-1.00	0.435	-5.32	3.32
Control	30 mins	-3.00	0.181	-7.32	1.32
15 mins	30 mins	-2.00	0.317	-6.32	2.32

### References

[1] The jamovi project (2022). jamovi. (Version 2.3) [Computer Software]. Retrieved from <a href="https://www.jamovi.org">https://www.jamovi.org</a>.

[2] R Core Team (2021). R: A Language and environment for statistical computing. (Version 4.1) [Computer software]. Retrieved from <a href="https://cran.r-project.org">https://cran.r-project.org</a>. (R packages retrieved from MRAN snapshot 2022-01-01).

[3] Fox, J., & Weisberg, S. (2020). *car: Companion to Applied Regression*. [R package]. Retrieved from <a href="https://cran.r-project.org/package=car">https://cran.r-project.org/package=car</a>.

[4] Lenth, R. (2020). *emmeans: Estimated Marginal Means, aka Least-Squares Means*. [R package]. Retrieved from <a href="https://cran.r-project.org/package=emmeans">https://cran.r-project.org/package=emmeans</a>.