

Dark Matter Halo Shape After Major Merger

Chip Fowler,¹★

¹*Department of Astronomy, University of Arizona, 933 N Cherry Ave, 85719, United States of America*

9 May 2025

ABSTRACT

Dark matter halos are the exclusive sites of galaxy formation; mergers between galaxies can impact the properties of the halo, such as shape, spin, and concentration. Studying dark matter halos is one of the ways we understand how galaxies change due to mergers. To investigate this topic, I will use the N-body simulation of the evolution of the Milky Way-M31-M33 system described in [van der Marel et al. \(2012\)](#) to study the merger between the Milky Way and M31. Using this simulation, I will determine the final shape of the combined dark matter halo. After analyzing its triaxiality, the halo was determined to be prolate. This result provides further evidence for the expectations for post-merger halo shape predicted in papers like [Abadi et al. \(2010\)](#) and [Drakos et al. \(2019\)](#); this result allows astronomers to differentiate how a halo changes shape by accreting over time versus changes due to a merger, which has implications for the mass distribution of the galaxy overall.

Key words: Local Group – Major Merger – Dark Matter Halo – Halo Shape – Hernquist Profile – Isodensity Contours

1 INTRODUCTION

Every galaxy is theorized to be surrounded by a **dark matter halo** – a virialized distribution of dark matter that is more densely populated than the surrounding region. Dark matter halos are decoupled from the expansion of the universe, and only grown by accreting in asymmetric "filaments" or "sheets" in space ([Chua et al. 2019](#)). Dark matter halos also attract baryons, and therefore, are the site of galaxy formation. When galaxies merge, their dark matter halos also merge with each other, changing their intrinsic properties. Quantifying how a merger affects the structure of a dark matter halo is one of the ways we can draw conclusions about how galaxies change over time.

A **galaxy**, as defined by [Willman & Strader \(2012\)](#), is "a gravitationally bound collection of stars whose properties cannot be explained by a combination of baryons and Newton's laws of gravity." In this definition, dark matter is necessary to explain the formation of galaxies; dark matter clusters exert the gravitational force needed to condense gas and dust into galaxies. Since the dark matter **halo shape** is unique to the galaxy it hosts, it is closely entwined with the galaxy's growth and merger history ([Drakos et al. 2019](#)). The shape of a halo can be approximated as an ellipsoid with 3 principle axes; halos are generally not spherical. Dark matter clusters are the only site of galaxy formation; therefore, understanding how a halo can be changed has implications for **galaxy evolution** overall. Galaxy evolution studies how galaxies in the early universe were assembled, and how they have been formed into the diverse structures we see today. Since dark matter halos are exclusive to the galaxy they surround, discovering patterns in dark matter halo properties could shed light on greater trends in galaxy evolution.

Galaxy mergers and dark matter halos are modeled using N-body simulations. An N-body simulation models interactions between particles in a system under the effects of some force; in the case of

galaxy mergers, an N-body simulation models collisions between two or more galaxies under the influence of each other's gravity. The baryon components of a galaxy such as gas, stars, and dust are difficult to simulate, so many of these simulations are done with only dark matter particles, in what is referred to as a DMO simulation ([Chua et al. 2019](#)). While these simulations are good approximations for dark matter-dominant regions of a galaxy, they are not accurate for the bright, baryon-dominant components of a galaxy such as the galactic center, which can alter the shape of nearby dark matter distribution (see Figure 1) ([Abadi et al. 2010](#)). Dark matter halo studies are primarily interested in studying the shape, spin, concentration, and mass profile of these structures.

There are many uncertainties when using N-body simulations to examine dark matter halos. How a halo is reshaped as a result of merger is, as previously mentioned, one of the primary focuses of dark matter halo studies. One of the biggest questions in regard to halo shape is the effect baryons have, especially on the "inner" halo where the galaxy is located ([Chua et al. 2019](#)). Papers such as [Abadi et al. \(2010\)](#) outline one way to tackle the inclusion of baryons in a simulation of this nature. Active galactic nuclei can also pose issues in determining the properties of dark matter halos, since they eject baryons and dark matter from the galactic center ([Chua et al. 2019](#)). In many simulations, such as [Drakos et al. \(2019\)](#), two equal-size galaxies are used to examine how a **major merger** influences the dark matter halo. Mergers between galaxies of comparable size are considered major mergers, while collisions between a larger galaxy and a smaller galaxy are considered minor mergers. In actuality, major mergers are rare; it is much more common for galaxies of different sizes to merge.

★ E-mail: nataliefowler@arizona.edu




Figure 1. Figure 4 from [Abadi et al. \(2010\)](#). In this paper, the effects the baryon component of a galaxy has on dark matter halo shape was examined. These graphs represent the distribution of particles included in an N-body simulation, plotted so that the galaxy is viewed face-on – position in the x direction versus the position in the z direction. The top panels represent simulations where particles were binned according to their gravitational potential, while in the bottom panels, they are binned by their density. On the left, only the particles that make up the dark matter halo were used, while on the right, the baryon components that make up the bright parts of the galaxy were included. In the right-hand panes, the shape and position of the central galaxy is depicted in black. In the dark matter + baryons simulation, the halo becomes more spherical due to the presence of the central galaxy.

2 THIS PROJECT

In this paper, I will examine and quantify the shape of the dark matter halo remnant resulting from a major merger between two galaxies in the **Local Group**: our Milky Way Galaxy and the Andromeda Galaxy. The Local Group describes the collection of galaxies near and including the Milky Way; of major interest is the study of the eventual merger of the Milky Way and the Andromeda Galaxy (M31), which are expected to collide in 4-5 billion years.

This paper is designed to address how the Milky Way-M31 merger will influence the shape of the remnant halo. From previous work such as [Drakos et al. \(2019\)](#), it appears that halo remnants are stretched along the axis where they collide. I will examine if predictions such as these are true for the halo remnant of this specific major merger.

Dark matter halos are unique to each galaxy; therefore, studying their evolution can lend some insight into the history of the galaxy itself. Since a majority of galaxies are thought to have gone through a merger, studying the dark matter halo can reveal trends in galaxy evolution that apply to many galaxies.

3 METHODOLOGY

In order to examine this merger, I will use a modified version of the N-body simulation described in [van der Marel et al. \(2012\)](#). This simulation does not include collisions between individual particles. Gas makes up only a small portion of the total galaxy mass, so it was excluded from the simulation as well. The galaxies in the simulation have dark matter halos represented by a **Hernquist Profile**. The Hernquist Profile, as defined in [Hernquist \(1990\)](#), is the distribution of mass within a given radius in a spherical object. Each galaxy has 500,000 dark matter particles that make up its halo. Each galaxy's disk mass was chosen to best suit its observed rotation curve and is




Figure 2. Figure 6 from [Drakos et al. \(2019\)](#). The results of a simulation of a major merger between two equal-sized galaxies. Plotted on top of a diagram of the halo particle distribution are iso-density contours (white) and the shape ratio (red). The shape ratio generally agrees with the isodensity contours. This diagram is very similar to what I intend to produce with my project.

represented by an exponential profile. Each galaxy's bulge mass was determined from literature values, and set up with an $R^{1/4}$ profile. Initially, the Milky Way is at rest and is set at the origin of the Galactocentric frame.

I will modify the simulation to only include the dark matter halo particles. Although the smaller Triangulum Galaxy (M33) is involved in this event and included in the simulation, only the Milky Way and M31 merge, therefore only the Milky Way and M31 particles will be considered. I will examine the combined halo remnant after the merger has occurred, and the dark matter halo has settled – about 9 billion years in the future, or around snap number 630 in the files for both galaxies. In order to examine the combined dark matter halo, I will concatenate the arrays for the positions and masses of each dark matter particle in each galaxy for the snapnumber 630 file. I will be using the high-res version of the file, since I am only examining one snapnumber instead of a sequence of files. Once I have the concatenated array of the combined halo particles, I will fit the Hernquist Profile to the combined Milky Way-M31 dark matter halo data. I will then use the Python package *photutils* to plot elliptical **isodensity contours** on 2-D histograms of the spatial projections. An isodensity contour represents points on the graph that have the same density (see Figure 2). Photutils will return the ellipticity of a given isodensity contour; I will use the ellipticity of the isodensity contour closest to the scale radius determined by fitting the Hernquist Profile to obtain the axial ratio of the projection. Once the axial ratios are obtained, they can be compared to determine the overall shape of the halo.

The Hernquist mass profile is defined as:

$$M(r) = M_{\text{halo}} r^2 / (a + r)^2 \quad (1)$$

Where M_{halo} is the mass of the dark matter halo, r is the distance from the Galactic center, and a is the scale radius. The scale radius of the Hernquist Profile represents the radius that encloses the majority of the mass in a galaxy. The scale radius for the merger remnant will be determined by trying multiple values until the profile closely resembles the halo mass distribution. After the projections of the galaxy in x, y, and z are plotted and the isodensity contours are fitted using photutils, the shape of the halo can be determined. Halos are ellipsoids, which are triaxial; their shapes are described by the axial ratios q and s . These ratios are written as

$$q = b/a \quad (2)$$

$$s = c/a \quad (3)$$

where a is the length of the major axis, b is the length of the intermediate axis, and c is the length of the minor axis. The axial ratios are related to the ellipticity by the generalized equation

$$e = 1 - m/w \quad (4)$$

Where e is the ellipticity, m is the length of the major axis of the ellipse, and w is the length of the minor axis of the ellipse. Chua et al. (2019) defines the "Triaxiality Parameter" as

$$T = (1 - q^2)/(1 - s^2) \quad (5)$$

where T is the triaxiality of the distribution, and q and s are the axial ratios described above. The value of T determines the shape of the halo; if $T > 0.67$, the halo is considered prolate, but if $T < 0.33$, the halo is considered oblate (Chua et al. 2019). If $0.33 < T < 0.67$, the halo is considered triaxial (Chua et al. 2019).

I will produce three plots in total. I will first plot and fit the Hernquist Profile to determine the scale radius. After the scale radius has been determined, I will create two plots of the particle position projections: one of x versus y , and one of z versus y . Using photutils, I will overplot isodensity contours on these histograms. Photutils will return the ellipticity of a given isodensity contour, which can be used to determine the axial ratio. I will compare the axial ratios taken from the elliptical contours closest to the scale radius; the scale radius encloses a majority of the remnant's mass, making it an appropriate choice to use to determine the triaxiality.

Following from the results of Drakos et al. (2019) and Abadi et al. (2010), I expect to find that the combined dark matter halo of the merged galaxies will become more axisymmetric. Galaxies on radial orbits will generally form prolate halo remnants, while galaxies on tangential orbits will form oblate remnants (Drakos et al. 2019). M31 is on a mostly radial orbit with respect to the Milky Way, so I expect the halo remnant will be prolate.

4 RESULTS

Figure 3 shows the fitting of the Hernquist Profile for the dark matter halo remnant. The remnant halo mass profile was created by computing the mass enclosed at about every 6 kpc, starting at 5 kpc from the center out to a radius of 200 kpc. The Hernquist Profile was computed via equation (1), using the same radius bins that were used to create the remnant mass profile. The best fit for the Hernquist Profile is achieved with a scale radius of 96 kpc; with this scale radius, the fractional difference between profiles is within 7% at all radii.

Figure 4 represents the particle projections in the x and y directions. The projection in x is plotted on the x axis and the projection in y is plotted on the y axis; in order to use photutils, the axes must be converted from kpc to pixels. The colormap represents the density of the particles; yellow areas are more densely packed, while dark purple areas are less dense. Elliptical isodensity contours were overplotted on the distribution using photutils. For determining the axial ratio q , I have chosen to define x as the intermediate axis b , and y as the major axis a . The ellipticity of the 96 kpc isodensity contour determined from this graph is 0.155. Using this result in equation 4, the axial ratio q is 0.845.

Figure 5 represents the particle projections in the z and y directions, with z plotted on the x axis and y plotted on the corresponding y axis. Like in Figure 4, the axes were converted from kpc to pixels in order to use photutils; also like Figure 4, the colormap represents densely populated areas in yellow, and sparsely populated areas in purple. To determine the axial ratio s , I have chosen to define z as the minor axis c , while the y axis represents the major axis a . The ellipticity of the




Figure 3. The fitting of the Hernquist Profile for the halo remnant. The distance from the center of the halo remnant is plotted on the x axis, while the log of the mass in $10^{10} M_{\text{sun}}$ is plotted on the y axis. The red line represents the mass of the combined dark matter halo at every radius, starting from 5 kiloparsecs from the center of the merged galaxy, up to 200 kiloparsecs. The dashed line represents the Hernquist Profile. The best fit for the Hernquist Profile is obtained using a scale radius of 96 kpc.




Figure 4. The dark matter particle projections in the x and y directions of the halo remnant. A colormap was used to represent the particle count density. Axes are plotted in pixel widths. Elliptical isodensity contours are plotted overtop in pink. In the inner halo, the distribution is more circular, while at larger radii, it becomes more elliptical.

96 kpc isodensity contour determined from this graph is 0.2. Using this result in equation 4, the axial ratio s is 0.802.

5 DISCUSSION

With the axial ratios q and s determined from Figures 4 and 5 respectively, the Triaxiality Parameter (equation 5) was found to be 0.8. Following the definition in Chua et al. (2019), this result suggests




Figure 5. The dark matter particle projections in the z and y directions of the halo remnant. A colormap was used to represent the particle count density. Axes are plotted in pixel widths. Elliptical isodensity contours are plotted overtop in pink. As observed in the XY projection histogram, the ellipticity of the halo appears to increase based on distance from the center.

that the remnant is prolate in shape. This measurement agrees with my hypothesis for the final shape.

The remnant's prolateness agrees with the predictions for halo shape based in papers such as [Drakos et al. \(2019\)](#) and [Abadi et al. \(2010\)](#); galaxies on radial orbits with respect to each other, such as the Milky Way and M31, will produce prolate halo remnants. This result's consistency with the literature suggests that dark matter halos undergo quantifiable and predictable changes based on the properties of the merger. These results can be applied to study the halos of galaxies astronomers observe today. The halo shape changes the potential well of the galaxy, altering how the galaxy accretes over time. The shape of the potential well affects the orbit of satellites; an object moving around a galaxy with a triaxial halo will orbit differently than it would around a galaxy with an oblate halo. Defining the halo shape of a host galaxy is of primary interest in describing the orbital history of a satellite object.

The triaxiality of the halo is dependent on the radius at which the axial ratios are measured. The choice of the scale radius of the Hernquist Profile is appropriate, but since it is only one measurement, it neglects how the triaxiality changes with radius. This measurement also excludes how the internal energy available to the remnant alters its shape; [Drakos et al. \(2019\)](#) determined that the axial ratio scaled roughly linearly with energy. This was a DMO simulation, which also ignores how the baryonic components of the galaxy influence the shape of the halo.

6 CONCLUSION

Dark matter halos are the exclusive sites of galaxy formation; mergers between galaxies can impact the properties of the halo, such as shape, spin, and concentration. Studying dark matter halos is one of the ways we understand how galaxies change due to mergers. To investigate this topic, I used the N-body simulation of the evolution of the Milky Way-M31-M33 system described in [van der Marel et al. \(2012\)](#) to study the merger between the Milky Way and M31. Using this simulation, I examined the final shape of the combined dark matter halo.

From this simulation, I determined that the dark matter halo remnant was prolate. This result agrees with my hypothesis and is consistent with the predictions made in [Drakos et al. \(2019\)](#). This result can be used to study galaxies of interest. A halo reshaped by a merger will not resemble a halo that has been accreted from dark matter flow over time. Defining how halos are assembled has implications for determining the orbit history of satellites and how baryons are accreted onto the galaxy.

This triaxiality measurement neglects how ellipticity changes with radius. To address this, multiple radii could be selected to compare axial ratios. A "triaxiality profile" could be made to compare how the shape changes with radius. This measurement also ignores how the baryonic component of the combined galaxy would affect the inner halo shape; [Abadi et al. \(2010\)](#) found that the dark matter halo became more spherical as a result of the inclusion of the central galaxy. A second simulation, including all particles in the simulation, would likely make the inner halo appear more spherical.

ACKNOWLEDGEMENTS

I thank Dr. Gurtina Besla and Himansh Rathore for their help with troubleshooting code.

This work made use of the following software packages: `astropy` ([Astropy Collaboration et al. 2013, 2018, 2022](#)), `matplotlib` ([Hunter 2007](#)), `numpy` ([Harris et al. 2020](#)), `python` ([Van Rossum & Drake 2009](#)), `scipy` ([Virtanen et al. 2020; Gommers et al. 2025](#)), and `scikit-image` ([van der Walt et al. 2014](#)).

This research made use of Photutils, an Astropy package for detection and photometry of astronomical sources ([Bradley et al. 2025](#)).

Software citation information aggregated using [The Software Citation Station](#) ([Wagg & Broekgaarden 2024; Wagg et al. 2024](#)).

We respectfully acknowledge the University of Arizona is on the land and territories of Indigenous peoples. Today, Arizona is home to 22 federally recognized tribes, with Tucson being home to the O'odham and the Yaqui. The University strives to build sustainable relationships with sovereign Native Nations and Indigenous communities through education offerings, partnerships, and community service.

REFERENCES

- Abadi M. G., Navarro J. F., Fardal M., Babul A., Steinmetz M., 2010, *Monthly Notices of the Royal Astronomical Society*, 407, 435
- Astropy Collaboration et al., 2013, *A&A*, 558, A33
- Astropy Collaboration et al., 2018, *AJ*, 156, 123
- Astropy Collaboration et al., 2022, *ApJ*, 935, 167
- Bradley L., et al., 2025, `astropy/photutils`: 2.2.0, doi:10.5281/zenodo.14889440, <https://doi.org/10.5281/zenodo.14889440>
- Chua K. T. E., Pillepich A., Vogelsberger M., Hernquist L., 2019, *Monthly Notices of the Royal Astronomical Society*, 484, 476
- Drakos N. E., Taylor J. E., Berrouet A., Robotham A. S. G., Power C., 2019, *Monthly Notices of the Royal Astronomical Society*, 487, 993
- Gommers R., et al., 2025, `scipy/scipy`: SciPy 1.15.2, doi:10.5281/zenodo.14880408, <https://doi.org/10.5281/zenodo.14880408>
- Harris C. R., et al., 2020, *Nature*, 585, 357
- Hernquist L., 1990, *ApJ*, 356, 359
- Hunter J. D., 2007, *Computing in Science & Engineering*, 9, 90
- Van Rossum G., Drake F. L., 2009, *Python 3 Reference Manual*. CreateSpace, Scotts Valley, CA

- Virtanen P., et al., 2020, *Nature Methods*, 17, 261
- Wagg T., Broekgaarden F. S., 2024, arXiv e-prints, p. [arXiv:2406.04405](https://arxiv.org/abs/2406.04405)
- Wagg T., Broekgaarden F., Gültekin K., 2024, TomWagg/software-citation-station: v1.2, [doi:10.5281/zenodo.13225824](https://doi.org/10.5281/zenodo.13225824), <https://doi.org/10.5281/zenodo.13225824>
- Willman B., Strader J., 2012, *The Astronomical Journal*, 144, 76
- van der Marel R. P., Besla G., Cox T. J., Sohn S. T., Anderson J., 2012, *The Astrophysical Journal*, 753, 9
- van der Walt S., et al., 2014, *PeerJ*, 2, e453

This paper has been typeset from a \LaTeX file prepared by the author.