

Analytical solutions for acoustic vortex beam radiation from planar and spherically focused circular pistons

Chirag A. Gokani Michael R. Haberman Mark F. Hamilton

Applied Research Laboratories & Walker Department of Mechanical Engineering
University of Texas at Austin

188th ASA & 25th ICA Meeting
New Orleans, Louisiana



The University of Texas at Austin
Walker Department
of Mechanical Engineering
Cockrell School of Engineering

Outline

Introduction

Analytical solution of paraxial equation

Vortex ring radius

A simplified diffraction integral

Summary

Outline

Introduction

Analytical solution of paraxial equation

Vortex ring radius

A simplified diffraction integral

Summary

Our publications on this work

JASA EXPRESS LETTERS

ARTICLE

asa.scitation.org/journal/jel

CrossMark

Analytical solutions for acoustic vortex beam radiation from planar and spherically focused circular pistons

Chirag A. Gokani,^{1,2,a)} Michael R. Haberman,^{1,2}  and Mark F. Hamilton^{1,2} 

¹Applied Research Laboratories, The University of Texas at Austin, Austin, Texas 78766-9767, USA

²Walker Department of Mechanical Engineering, The University of Texas at Austin, Austin, Texas 78712-1063, USA

chiragokani@utexas.edu, haberman@utexas.edu, hamilton@mail.utexas.edu

[Editor: Régis Marchiano] <https://doi.org/10.1121/10.0034739>

Received: 19 October 2024 Accepted: 5 December 2024 Published Online: 24 December 2024

JASA Express Lett. **4** (12), 124001 (2024) © Author(s) 2024  4, 124001-1

JASA ARTICLE

CrossMark  click for updates

Paraxial and ray approximations of acoustic vortex beams

Chirag A. Gokani,^{1,2} Michael R. Haberman,^{1,2}  and Mark F. Hamilton^{1,2,a)} 

¹Applied Research Laboratories, The University of Texas at Austin, Austin, Texas 78766-9767, USA

²Walker Department of Mechanical Engineering, The University of Texas at Austin, Austin, Texas 78712-1063, USA

(Received 19 January 2024; revised 27 March 2024; accepted 27 March 2024; published online 22 April 2024)

[Editor: Philip L. Marston]



Pages: 2707–2723

J. Acoust. Soc. Am. **155** (4), April 2024 © 2024 Acoustical Society of America


What is an acoustic vortex beam?

Characterized by...

- ▶ helical wavefronts
- ▶ orbital number $\ell =$ number of equiphase wavefronts in \perp plane
- ▶ zero acoustic pressure on axis



C. Shi et al. *P. Natl. Acad. Sci. U.S.A.* 114 (2017), pp. 7250–7253




B. T. Hefner and P. L. Marston. *J. Acoust. Soc. Am.* 106 (1999), pp. 3313–3316


What is an acoustic vortex beam?

Used for...


- ▶ particle manipulation
- ▶ underwater communications
- ▶ therapeutic biomedical ultrasound
- ▶ sound diffusion




L. Zhang and
P. L. Marston.
Phys. Rev. E 84
(2011), pp. 1–5



M. E. Kelly and C. Shi.
JASA Express Lett. 3
(2023), pp. 1–5



S. Guo et al.
Ultrasound Med. Biol. 48
(2022),
pp. 1907–1917



N. Jiménez,
J. -P. Groby, and
V. Romero-García.
Sci. Rep. 11 (2021),
pp. 1–13

What is an acoustic vortex beam?

Generated by...

- ▶ phase plates
- ▶ transducer arrays
- ▶ metasurfaces



M. E. Terzi et al. *Moscow University Physics Bulletin* 72 (2017), pp. 61–67



A. Marzo, M. Caleap, and B. W. Drinkwater.
Phys. Rev. Lett. 120 (2018),
pp. 1–6



X. Jiang et al. *Phys. Rev. Lett.* 117 (2016), pp. 1–5

Previous analytical descriptions of vortex beams

- Bessel vortex beams are modes of the cylindrical wave equation:¹

$$p(r, \theta, z, t) = p_0 J_\ell(k_r r) e^{i(\ell\theta + k_z z - \omega t)}, \quad k = \frac{\omega}{c_0} = \sqrt{k_r^2 + k_z^2}. \quad (1)$$

- Gaussian vortex beams are generalizations of Gaussian beams:²

$$\begin{aligned} p(r, \theta, z, t) &= \sqrt{8\pi} \left(\frac{p_0 z}{k r^2} \right) \chi^{3/2} e^{-\chi} [I_{(\ell-1)/2}(\chi) - I_{(\ell+1)/2}(\chi)] \\ &\times e^{i[\ell\theta - (\ell+1)\pi/2 + kr^2/2z + k_z z - \omega t]}, \end{aligned} \quad (2)$$

$$\chi(r, z) = \frac{\frac{1}{8}(kar/z)^2}{1 - i(ka^2/2z)(1 - z/d)}.$$

- Fields described by Eqs. (1) and (2) require infinite source conditions.
- Equation (1) implies infinite energy,³ because $\int_0^\infty |J_\ell(k_r r)|^2 r dr \rightarrow \infty$.
- **Objective: derive solutions for vortex fields radiated by circular pistons**

¹N. Jiménez et al. *Phys. Rev. E* 94 (2016), pp. 1–9.

²C. A. Gokani, M. R. Haberman, and M. F. Hamilton. *J. Acoust. Soc. Am.* 155 (2024), pp. 2707–2723.

³M. R. Lapointe. *Opt. Laser Technol.* 24 (1992), pp. 315–321.

Outline

Introduction

Analytical solution of paraxial equation

Vortex ring radius

A simplified diffraction integral

Summary

Paraxial equation and its integral solution


- For $p = qe^{i(kz - \omega t)}$ and $|\partial^2 q / \partial z^2| \ll 2k |\partial q / \partial z|$, $\nabla^2 p - c_0^{-2} \ddot{p} = 0$ reduces to

$$i2k \frac{\partial q}{\partial z} + \nabla_{\perp}^2 q = 0. \quad (3)$$

- ∇_{\perp}^2 is the Laplacian in the plane perpendicular to the z axis.
- Equation (3) is solved by the Fresnel diffraction integral:

$$q(r, \theta, z) = -\frac{ik}{2\pi z} \int_0^{2\pi} \int_0^{\infty} q(r_0, \theta_0, 0) e^{i(k/2z)[r^2 + r_0^2 - 2rr_0 \cos(\theta_0 - \theta)]} r_0 dr_0 d\theta_0,$$

where $q(r_0, \theta_0, 0)$ is the prescribed pressure field in the plane $z = 0$.



Vortex radiation from unfocused circular piston

- A **circular vortex** source condition is first considered:

$$q(r, \theta, 0) = p_0 \operatorname{circ}(r/a) e^{i\ell\theta}, \quad (4)$$

where $\operatorname{circ}(x) = 1$ for $0 \leq x \leq 1$ and 0 for $x > 1$.

- Insertion of Eq. (4) in the Fresnel diffraction integral leads to

$$q = -ikp_0 \frac{e^{i(ka^2/2z)r^2/a^2}}{z} e^{i\ell(\theta-\pi/2)} \int_0^a e^{i(ka^2/2z)r_0^2/a^2} J_\ell(krr_0/z) r_0 dr_0. \quad (5)$$

- Using Watson's relation,⁴ Eq. (5) reduces to

$$q = -\frac{ikp_0}{z} e^{i\ell(\theta-\pi/2)} \int_0^a J_\ell(krr_0/z) r_0 dr_0 \quad (6)$$

for $z \gg z_R$, where $z_R = ka^2/2$ is the Rayleigh distance.

⁴G. N. Watson, *A Treatise on the Theory of Bessel Functions*, 2nd ed. Sec. 2.2, Eq. (5).

Vortex radiation from unfocused circular piston

- Taking the integral in Eq. (6) leads to an analytical solution:

$$q_\ell(r, \theta, z) = -ip_0 \frac{z}{kr^2} e^{i\ell(\theta-\pi/2)} F_\ell(kar/z), \quad z \gg z_R, \quad (7)$$

where⁵

$$F_\ell(\xi) = \int_0^\xi J_\ell(t) t dt = \xi \frac{\Gamma(\ell/2 + 1)}{\Gamma(\ell/2)} \sum_{k=0}^{\infty} \frac{(\ell + 2k + 1)\Gamma(\ell/2 + k)}{\Gamma(\ell/2 + 2 + k)} J_{\ell+2k+1}(\xi). \quad (8)$$

- Equation (8) equals the following closed-form expressions for $1 \leq \ell \leq 4$:

$$F_0(\xi) = \xi J_1(\xi), \quad \ell = 0 \quad (9a)$$

$$F_1(\xi) = \frac{\pi}{2}\xi [\mathbf{H}_0(\xi)J_1(\xi) - \mathbf{H}_1(\xi)J_0(\xi)], \quad \ell = 1 \quad (9b)$$

$$F_2(\xi) = 2 - 2J_0(\xi) - \xi J_1(\xi), \quad \ell = 2 \quad (9c)$$

$$F_3(\xi) = \left[\frac{3\pi}{2}\xi \mathbf{H}_0(\xi) - 8 \right] J_1(\xi) + \left[4\xi - \frac{3\pi}{2}\xi \mathbf{H}_1(\xi) \right] J_0(\xi), \quad \ell = 3 \quad (9d)$$


$$F_4(\xi) = 4 - 8J_1(\xi)/\xi - 4J_2(\xi) - \xi J_3(\xi), \quad \ell = 4 \quad (9e)$$

⁵M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions*, 4th ed. Item 11.1.1.

Verification of Eq. (7)

- The validity of Eq. (7) is assessed by comparison to

$$q(x, y, z) = \mathcal{F}_{xy}^{-1} \{ e^{ik_z z} \mathcal{F}_{xy}[q(x, y, 0)] \}, \quad k_z = k - (k_x^2 + k_y^2)/2k. \quad (10)$$



Vortex radiation from focused circular piston


- To describe spherical focusing at a geometric focal length d , source condition (4) is multiplied by $\exp(-ikr^2/2d)$:

$$q(r, \theta, 0) = p_0 \text{circ}(r/a) e^{i\ell\theta} e^{-ikr^2/2d}, \quad (11)$$


- An analytical solution of Eq. (3) is available at $z = d$:

$$q_\ell(r, \theta, d) = -ip_0 \frac{d}{kr^2} e^{ikr^2/2d} e^{i\ell(\theta-\pi/2)} F_\ell(kar/d), \quad (12)$$

where $F_\ell(\xi)$ is given by Eq. (8).



Verification of Eq. (12): analytical, Fourier



Outline

Introduction

Analytical solution of paraxial equation


Vortex ring radius

A simplified diffraction integral

Summary

Vortex ring radius


- ▶ The magnitudes of vortex beam fields are axisymmetric.
- ▶ In the far field of Eq. (7), the field is conical.
- ▶ In the geometric focal plane $z = d$ of Eq. (12), the field forms toroidal ring.
- ▶ Equations (7) and (12) can be used to find the radius of these features.



C. Zhou et al. *J. Appl. Phys.* 128 (2020), pp. 1–12



D. Baresch, J. -L. Thomas, and R. Marchiano. *Phys. Rev. Lett.* 116 (2016), pp. 1–6



M. E. Terzi et al. *Moscow University Physics Bulletin* 72 (2017), pp. 61–67

Vortex ring radius

- Maximizing Eqs. (7) and (12) in r amounts to solving

$$\frac{d|\xi^{-1} F_\ell(\xi)|}{d\xi} = 0,$$

where $\xi = kar/z$ (unfocused) and $\xi = kar/d$ (focused).

- Using Eq. (8) for F_ℓ and taking the derivative yields⁶

$$\sum_{k=0}^{\infty} \frac{(\ell + 2k + 1)\Gamma(\ell/2 + k)}{\Gamma(\ell/2 + 2 + k)} \left[\frac{J_{\ell+2k}(\xi) - J_{\ell+2k+2}(\xi)}{2\xi} - \frac{J_{\ell+2k+1}(\xi)}{\xi^2} \right] = 0. \quad (13)$$


- The roots ξ_ℓ of Eq. (13) are fit to a line: $\xi_\ell = 1.23\ell + 1.49$.
- Solving $\xi_\ell = kar_\ell/z$ and $\xi_\ell = kar_\ell/d$ for r_ℓ yields

$$r_\ell = \frac{\xi_\ell z}{ka}, \quad z \gg z_R, \quad (14a)$$

$$= \frac{\xi_\ell d}{ka}, \quad z = d. \quad (14b)$$

⁶I. S. Gradshteyn and I. M. Ryzhik. New York: Academic Press, 1980, Item 8.471-2.

Vortex ring radius



Comparison of roots ξ_ℓ (circles) with least-squares fit $\xi_\ell = 1.23\ell + 1.49$ (line)

Outline

Introduction

Analytical solution of paraxial equation

Vortex ring radius

A simplified diffraction integral

Summary

A simplified diffraction integral

- The angular spectrum method is equivalent to the first Rayleigh integral:


$$p(r, \theta, z) = \rho_0 c_0 k \mathcal{F}_{2D}^{-1}\{\mathcal{F}_{2D}\{u_z(r, \theta)\}\} e^{ik_z z} / k_z , \quad (15)$$

where

$$\mathcal{F}_{2D}\{f(r, \theta)\} = g(k_r, \psi) = \int_0^{2\pi} \int_0^{\infty} f(r, \theta) e^{-ik_r r \cos(\theta - \psi)} r dr d\theta , \quad (16a)$$

$$\mathcal{F}_{2D}^{-1}\{g(k_r, \psi)\} = f(r, \theta) = \int_0^{2\pi} \int_0^{\infty} g(k_r, \psi) e^{ik_r r \cos(\theta - \psi)} k_r dk_r d\psi . \quad (16b)$$

- Directly evaluating Eq. (15) can suffer from aliasing and discretization error.⁷



⁷J. Blackmore, R. O. Cleveland, and J. Mobley. *J. Acoust. Soc. Am.* 144 (2018), pp. 2947–2951.

A simplified diffraction integral


- In light of F_ℓ given by Eq. (8) and Watson's relation⁸

$$J_n(\beta) = \frac{1}{2\pi} \int_{\alpha}^{2\pi+\alpha} e^{i(n\phi - \beta \sin \phi)} d\phi, \quad (17)$$

Eq. (15) reduces to

$$p/\rho_0 c_0 u_0 = e^{i\ell\theta} \int_0^{\infty} (k/k_z) F_\ell(k_r a) J_\ell(k_r r) e^{ik_z z} dk_r / k_r. \quad (18)$$

- Equation (18) can be derived by considering the limiting case of a piston in a tube, amounting to an alternative theory of diffraction.



⁸G. N. Watson, *A Treatise on the Theory of Bessel Functions*, 2nd ed. Sec. 2.2, Eq. (5).

A simplified diffraction integral

- In terms of the dimensionless parameters

$$P \equiv p/\rho_0 c_0 u_0, \quad R \equiv r/a, \quad Z \equiv z/z_R, \quad K \equiv ka, \quad (19)$$

where z_R is the Rayleigh distance $ka^2/2$, and where

$$k_z/k = \sqrt{1 - (\zeta/K)^2}, \quad \zeta \equiv k_r a, \quad (20)$$

Eq. (18) becomes

$$P = e^{i\ell\theta} \int_0^\infty \frac{F_\ell(\zeta) J_\ell(\zeta R)}{\zeta \sqrt{1 - (\zeta/K)^2}} e^{iK^2 Z \sqrt{1 - (\zeta/K)^2}/2} d\zeta. \quad (21)$$

- Equation (21) is equivalent to and easier to evaluate than Eq. (15).



Mr. Jackson S. Hallveld



Dr. Randall P. Williams

A simplified diffraction integral

- Equation (21) can be evaluated analytically for $R = \ell = 0$:

$$P(Z) = \int_0^\infty \frac{J_1(\zeta)}{\sqrt{1 - (\zeta/K)^2}} e^{iK^2 Z \sqrt{1 - (\zeta/K)^2}/2} d\zeta. \quad (22)$$

- Equation (22) evaluates to⁹

$$P(Z) = -iK I_{1/2}[-i\chi_-(Z)] K_{1/2}[-i\chi_+(Z)], \quad (23)$$

where I_ν and K_ν are the modified Bessel functions of order ν , and where

$$\chi_\pm(Z) = \frac{K}{2} [\sqrt{1 + (KZ/2)^2} \pm KZ/2]. \quad (24)$$


- Bessel function identities reduce Eq. (23) to


$$P(Z) = -2i \sin[\chi_-(Z)] e^{i\chi_+(Z)}, \quad (25)$$

recovering the on-axis pressure radiated by a planar circular piston.¹⁰

⁹I. S. Gradshteyn and I. M. Ryzhik. New York: Academic Press, 1980, Item 6.637-1.

¹⁰A. D. Pierce. Secs. 8.2 and 8.5. Cham, Switzerland: Springer, 2019, Eq. (5.7.3).

Equation (21) for $\ell = 0$: semi-analytical, Fourier

Equation (21) for $\ell = 1$: semi-analytical, Fourier

Outline

Introduction

Analytical solution of paraxial equation

Vortex ring radius

A simplified diffraction integral

Summary

Thank you for listening!

Summary


- ▶ Solved paraxial equation for planar and focused circular vortex sources
- ▶ Calculated the ring radius for both solutions
- ▶ Derived simplified diffraction integral equivalent to Rayleigh integral

Further reading

- ▶ "Paraxial and ray approximations of acoustic vortex beams,"
J. Acoust. Soc. Am. **155**, 2707–2723 (2024).
- ▶ "Analytical solutions for acoustic vortex beam radiation from planar and spherically focused circular pistons," JASA Express Lett. **4**, 124001 (2024).

Acknowledgments

- ▶ Chester M. McKinney Graduate Fellowship in Acoustics at the Applied Research Laboratories



Mr. Chirag A. Gokani



Prof. Michael R. Haberman



Prof. Mark F. Hamilton

References I

-  C. Shi et al. "High-speed acoustic communication by multiplexing orbital angular momentum". *P. Natl. Acad. Sci. U.S.A.* 114 (2017), pp. 7250–7253.
-  B. T. Hefner and P. L. Marston. "An acoustical helicoidal wave transducer with applications for the alignment of ultrasonic and underwater systems". *J. Acoust. Soc. Am.* 106 (1999), pp. 3313–3316.
-  Zhang, L. and P. L. Marston. "Angular momentum flux of nonparaxial acoustic vortex beams and torques on axisymmetric objects". *Phys. Rev. E* 84 (2011), pp. 1–5.
-  M. E. Kelly and C. Shi. "Design and simulation of acoustic vortex wave arrays for long-range underwater communication". *JASA Express Lett.* 3 (2023), pp. 1–5.
-  S. Guo et al. "Enhanced Sonothrombolysis Induced by High-Intensity Focused Acoustic Vortex". *Ultrasound Med. Biol.* 48 (2022), pp. 1907–1917.
-  N. Jiménez, J. -P. Groby, and V. Romero-García. "Spiral sound-diffusing metasurfaces based on holographic vortices". *Sci. Rep.* 11 (2021), pp. 1–13.

References II

-  M. E. Terzi et al. "Generation of a vortex ultrasonic beam with a phase plate with an angular dependence of the thickness". *Moscow University Physics Bulletin* 72 (2017), pp. 61–67.
-  A. Marzo, M. Caleap, and B. W. Drinkwater. "Acoustic virtual vortices with tunable orbital angular momentum for trapping of mie particles". *Phys. Rev. Lett.* 120 (2018), pp. 1–6.
-  X. Jiang et al. "Convert acoustic resonances to orbital angular momentum". *Phys. Rev. Lett.* 117 (2016), pp. 1–5.
-  N. Jiménez et al. "Formation of high-order acoustic Bessel beams by spiral diffraction gratings". *Phys. Rev. E* 94 (2016), pp. 1–9.
-  C. A. Gokani, M. R. Haberman, and M. F. Hamilton. "Paraxial and ray approximations of acoustic vortex beams". *J. Acoust. Soc. Am.* 155 (2024), pp. 2707–2723.
-  M. R. Lapointe. "Review of non-diffracting Bessel beam experiments". *Opt. Laser Technol.* 24 (1992), pp. 315–321.

References III

-  C. Zhou et al. "Focused acoustic vortex generated by a circular array of planar sector transducers using an acoustic lens, and its application in object manipulation". *J. Appl. Phys.* 128 (2020), pp. 1–12.
-  D. Baresch, J. -L. Thomas, and R. Marchiano. "Observation of a single-beam gradient force acoustical trap for elastic particles: Acoustical tweezers". *Phys. Rev. Lett.* 116 (2016), pp. 1–6.
-  I. S. Gradshteyn and I. M. Ryzhik. *Table of Integrals, Series, and Products, 4th edition*. New York: Academic Press, 1980.
-  J. Blackmore, R. O. Cleveland, and J. Mobley. "Spatial filters suppress ripple artifacts in the computation of acoustic fields with the angular spectrum method". *J. Acoust. Soc. Am.* 144 (2018), pp. 2947–2951.
-  A. D. Pierce. *Acoustics, 3rd edition*. Secs. 8.2 and 8.5. Cham, Switzerland: Springer, 2019.

Notation |

| Symbol | Description | Dimensions |
|--------------|-----------------------------------------------------------|----------------------------------|
| a | source radius | m |
| c_0 | speed of sound | m s^{-1} |
| d | focal length | m |
| G | focusing gain $ka^2/2d$ | 1 |
| i | complex unit | 1 |
| k | wavenumber | m^{-1} |
| ℓ | orbital number | 1 |
| ρ_0 | ambient mass density | kg m^{-3} |
| p | acoustic pressure | $\text{kg m}^{-1} \text{s}^{-2}$ |
| q | paraxial pressure | $\text{kg m}^{-1} \text{s}^{-2}$ |
| \mathbf{R} | separation vector $\mathbf{R} = \mathbf{r} - \mathbf{r}'$ | m |
| \mathbf{r} | position vector | m |
| \mathbf{v} | particle velocity | m s^{-1} |
| z_R | Rayleigh distance, $ka^2/2$ | m |
| ω | angular frequency, $\omega = 2\pi f$ | s^{-1} |