

Effects of increasing orbital number on the field transformation in focused vortex beams

Chirag A. Gokani Michael R. Haberman Mark F. Hamilton

Applied Research Laboratories & Walker Department of Mechanical Engineering
University of Texas at Austin

5pPA, General Topics in Physical Acoustics
186th ASA Meeting, Ottawa



TEXAS
The University of Texas at Austin

Our paper on this topic



Paraxial and ray approximations of acoustic vortex beams

Chirag A. Gokani,^{1,2} Michael R. Haberman,^{1,2} and Mark F. Hamilton^{1,2,a)}

¹*Applied Research Laboratories, The University of Texas at Austin, Austin, Texas 78766-9767, USA*

²*Walker Department of Mechanical Engineering, The University of Texas at Austin, Austin, Texas 78712-1063, USA*

(Received 19 January 2024; revised 27 March 2024; accepted 27 March 2024; published online 22 April 2024)

[Editor: Philip L. Marston]

Pages: 2707–2723

Outline

1 Introduction

2 Ray theory

3 Gaussian vortex source

4 Uniform vortex source

5 Summary

Outline

1 Introduction

2 Ray theory

3 Gaussian vortex source



4 Uniform vortex source

5 Summary


What is an acoustic vortex beam?

Characterized by...

- helical wavefronts
- orbital number $\ell =$ number of equiphase wavefronts in \perp plane
- zero pressure on axis



Shi et al. 2017, *P. Natl. Acad. Sci. U.S.A.*



Hefner and Marston 1999,
J. Acoust. Soc. Am.

What is an acoustic vortex beam?

Generated by...

- phase plates
- transducer arrays
- metasurfaces



M. E. Terzi et al. 2017, *Moscow University Physics Bulletin*



Marzo, Caleap, and Drinkwater
2018, *Phys. Rev. Lett.*



Jiang et al. 2016,
Phys. Rev. Lett.


Focused vortex beams

- Focused beams used for particle manipulation
- Magnitudes of vortex beam fields are axisymmetric
- In geometric focal plane $z = d$, field forms toroidal ring
- Weak spherical focusing studied in paraxial approximation,


$$i2k \frac{\partial q}{\partial z} + \nabla_{\perp}^2 q = 0, \quad p = qe^{ikz} \quad (1)$$



Baresch, Thomas, and
Marchiano 2016,
Phys. Rev. Lett.



Zhou et al. 2020, *J. Appl. Phys.*



M. E. Terzi et al. 2017, *Moscow University Physics Bulletin*


Prefocal shadow zone

- Magnitude of solution of Eq. (1) for focused Gaussian vortex source is

$$|q(r, z)| = \sqrt{8\pi} \frac{p_0 z}{kr^2} \left| \chi^{3/2} e^{-\chi} [I_{(\ell-1)/2}(\chi) - I_{(\ell+1)/2}(\chi)] \right|, \quad (2)$$

$$\chi(r, z) = \frac{\frac{1}{8}(kar/z)^2}{1 - i(ka^2/2z)(1 - z/d)}$$

- For moderate values of focusing gain G , Eq. (2) reveals movement of vortex ring out of focal plane $z = d$ as ℓ increases



Outline

1 Introduction

2 Ray theory

3 Gaussian vortex source

4 Uniform vortex source

5 Summary

Focused vortex source condition

- To explain field transformation with increasing ℓ , appeal to *ray theory*:
 $ka \rightarrow \infty$
- In homogeneous media, rays travel in straight lines
- In the vicinity of source, pressure field is

$$p(r, \theta, z) \simeq p_0 f(r) e^{i\phi}, \quad z \simeq 0$$


where $f(r)$ = axisymmetric amplitude distribution in source plane

- Phase accounts for **focusing**, **helical wavefronts**, and **traveling wave motion**:

$$\phi(r, \theta, z) = -kr^2/2d + \ell\theta + kz$$



Sunbeams





Wave normal \mathbf{n} and annular channel radius Δ

- Wave normal in source plane at distance r_0 from origin is

$$\mathbf{n} = \frac{\nabla\phi}{|\nabla\phi|} = \frac{-(r_0/d)\mathbf{e}_r + (\ell/kr_0)\mathbf{e}_\theta + \mathbf{e}_z}{\sqrt{(r_0/d)^2 + (\ell/kr_0)^2 + 1}}$$

- Radius of circle formed by family of rays emanating from $r = r_0$ in source plane is

$$\Delta(r_0, z) = r_0 \left[(1 - z/d)^2 + (\ell d / kr_0^2)^2 (z/d)^2 \right]^{1/2}$$



Adapted from Richard et al. 2020, *New J. Phys.*

Pressure field according to ray theory

- Pressure field predicted by ray theory:


$$P(\Delta, z) = p_0 f(r_0) \sqrt{A(r_0, 0)/A(r_0, z)} \quad (3)$$

- Area of annular ray channel is

$$A(r_0, z) = A_z(r_0, z) \cos \psi(r_0, z) = 2\pi w \frac{\Delta(r_0, z)|\partial\Delta/\partial r_0|}{\sqrt{1 + (\partial\Delta/\partial z)^2}} \quad (4)$$

- Inserting Eq. (4) into Eq. (3) and calculating $\partial\Delta/\partial r_0$ and $\partial\Delta/\partial z$ yields

$$P(\Delta, z) = p_0 f(r_0) \left[\frac{\cos \psi(r_0, 0) / \cos \psi(r_0, z)}{|(1 - z/d)^2 - (\ell d / kr_0^2)^2 (z/d)^2|} \right]^{1/2}$$



Caustics

- Caustics occur when cross-sectional area vanishes, i.e.,

$$A_z(r_0, z) = 2\pi r_0 w \left| (1 - z/d)^2 - (\ell d / k r_0^2)^2 (z/d)^2 \right| = 0 \quad (5)$$

- Substitution of roots of Eq. (5) into $\Delta(r_0, z)$ gives caustic coordinates:


$$\Delta_c(z) = \sqrt{(2\ell d/k)(z/d)|1 - z/d|} \quad (6)$$

- Squaring Eq. (6), notating $G = ka^2/2d$, and identifying

$$\Delta_c^2 = x_c^2 + y_c^2, \quad a_c^2 = \ell a^2 / 4G, \quad c_c^2 = d^2 / 4$$


reveals that prefocal caustic is a spheroid of volume $V_c = \frac{1}{6}\ell\lambda d^2$:

$$\frac{x_c^2 + y_c^2}{a_c^2} + \frac{(z_c - d/2)^2}{c_c^2} = 1, \quad 0 \leq z_c \leq d$$




Prolate for $(d/a)^2 G > \ell$

Source
plane



Oblate for $(d/a)^2 G < \ell$



Outline

1 Introduction

2 Ray theory

3 Gaussian vortex source

4 Uniform vortex source

5 Summary

Paraxial field $q(r, z)$, ray paths Δ , and caustics Δ_c




Figure 1: Overlays of caustics Δ_c (thick lines) and ray channels Δ (thin lines) on color plots of the amplitude of the paraxial field $|q(r, z)|$ due to a focused Gaussian vortex source. Rays emanating from $r > a$ have been suppressed for visual clarity.

Ray pressure field $P(\Delta, z)$




Figure 2: Color plots for the ray field $P(\Delta, z)$ due to a focused Gaussian vortex source.

Unfocused limit, $d \rightarrow \infty$

- For unfocused vortex beams, $d \rightarrow \infty$, for which Δ , Δ_c , and P reduce to



$$\Delta(r_0, z) = r_0[1 + (\ell z / kr_0^2)^2]^{1/2}, \quad \text{ray channel radius}$$

$$\Delta_c(z) = \sqrt{2\ell z / k}, \quad \text{caustic radius}$$

$$P(\Delta, z) = p_0 f(r_0) \left[\frac{\cos \psi(r_0, 0) / \cos \psi(r_0, z)}{|1 - (\ell z / kr_0^2)^2|} \right]^{1/2}$$

- Caustic surface is a paraboloid, where $z_R = ka^2/2$:

$$\frac{z}{z_R} = \frac{x_c^2 + y_c^2}{\ell a^2}, \quad z \geq 0$$



Unfocused limit, $d \rightarrow \infty$

- For unfocused vortex beams, $d \rightarrow \infty$, for which Δ , Δ_c , and P reduce to



$$\Delta(r_0, z) = r_0[1 + (\ell z / kr_0^2)^2]^{1/2}, \quad \text{ray channel radius}$$

$$\Delta_c(z) = \sqrt{2\ell z / k}, \quad \text{caustic radius}$$

$$P(\Delta, z) = p_0 f(r_0) \left[\frac{\cos \psi(r_0, 0) / \cos \psi(r_0, z)}{|1 - (\ell z / kr_0^2)^2|} \right]^{1/2}$$

- Caustic surface is a paraboloid, where $z_R = ka^2/2$:

$$\frac{z}{z_R} = \frac{x_c^2 + y_c^2}{\ell a^2}, \quad z \geq 0$$



Outline

1 Introduction

2 Ray theory

3 Gaussian vortex source

4 Uniform vortex source

5 Summary

Paraxial field $q(r, z)$, ray paths Δ , and caustics Δ_c




Figure 3: Overlays of caustics Δ_c (thick lines) and ray channels Δ (thin lines) on color plots of the amplitude of the paraxial field $|q(r, z)|$ due to a uniform focused vortex source.

Unfocused limit, $d \rightarrow \infty$




Figure 4: Overlays of caustics Δ_c (thick lines) and ray channels Δ (thin lines) on color plots of the amplitude of the paraxial field $|q(r, z)|$ due to a uniform unfocused vortex source.

Outline

1 Introduction

2 Ray theory

3 Gaussian vortex source

4 Uniform vortex source

5 Summary

$$[T]\hbar an \cdot k \quad Y_0 \mathbf{u} \quad f(0) \mathbf{r} \quad \ell iSte^{ni} \mathbf{n} G!$$

Summary

- Developed ray theory to describe field transformation associated with increase in orbital number
- Showed that shadow zone in prefocal region is bounded by a spheroidal surface
- Calculated pressure field from ray theory

Acknowledgments

- Chester M. McKinney Graduate Fellowship in Acoustics at the Applied Research Laboratories



Chirag Gokani



Prof. Mark Hamilton



Prof. Michael Haberman

References I

-  Shi, Chengzhi et al. (2017). "High-speed acoustic communication by multiplexing orbital angular momentum". In: *P. Natl. Acad. Sci. U.S.A.* 114.28, pp. 7250–7253.
-  Hefner, Brian T and Philip L Marston (1999). "An acoustical helicoidal wave transducer with applications for the alignment of ultrasonic and underwater systems". In: *J. Acoust. Soc. Am.* 106.6, pp. 3313–3316.
-  M. E. Terzi et al. (2017). "Generation of a vortex ultrasonic beam with a phase plate with an angular dependence of the thickness". In: *Moscow University Physics Bulletin* 72, pp. 61–67.
-  Marzo, Asier, Mihai Caleap, and Bruce W Drinkwater (2018). "Acoustic virtual vortices with tunable orbital angular momentum for trapping of mie particles". In: *Phys. Rev. Lett.* 120.4, p. 044301.
-  Jiang, Xue et al. (2016). "Convert acoustic resonances to orbital angular momentum". In: *Phys. Rev. Lett.* 117.3, p. 034301.
-  Baresch, Diego, Jean-Louis Thomas, and Régis Marchiano (2016). "Observation of a single-beam gradient force acoustical trap for elastic particles: Acoustical tweezers". In: *Phys. Rev. Lett.* 116.2, p. 024301.

References II

-  Zhou, Chenchen et al. (2020). "Focused acoustic vortex generated by a circular array of planar sector transducers using an acoustic lens, and its application in object manipulation". In: *J. Appl. Phys.* 128, p. 084901.
-  Richard, Grace et al. (2020). "Twisting waves increase the visibility of nonlinear behaviour". In: *New J. Phys.* 22.6, p. 063021.
-  A. D. Pierce (2019). *Acoustics, 3rd edition*. Secs. 8.2 and 8.5. Cham, Switzerland: Springer.

Pressure field from ray theory (extra)


- Inserting $p(\mathbf{x}) = P(\mathbf{x}, \omega) e^{i\omega\tau(\mathbf{x})}$ into $\nabla^2 p + k^2 p = 0$ yields¹

$$\nabla^2 P + i\omega[2\nabla P \cdot \nabla \tau + P \nabla^2 \tau] - \omega^2 P[(\nabla \tau)^2 - c^{-2}] = 0. \quad (8.5.1)$$

- In limit that $\omega \rightarrow \infty$, it is necessary for $(\nabla \tau)^2 = c^{-2}$ and

$$\nabla \cdot (P^2 \nabla \tau) = 0 \quad (8.5.3b)$$

- Areas $A(\mathbf{x}_0)$ and $A(\mathbf{x})$ define the end caps of a *ray tube*



- Integrating Eq. (8.5.3b) over \mathcal{V} and applying Gauss's theorem yields

$$\oint_S (P^2 \nabla \tau) \cdot \mathbf{n} dS = 0 \implies P(\mathbf{x}) = P(\mathbf{x}_0) \sqrt{A(\mathbf{x}_0)/A(\mathbf{x})}.$$

¹Equation numbers refer to those in *Acoustics*, A. D. Pierce 2019.