Antibiotic Resistance in Osun State Abattoirs

Town: All

Site_Code: All

MDR Isolates

110


MDR %

40.15%


Town	AUGMENTIN	CEFTAZIDIME	CIPROFLOXACIN	GENTAMICIN	IMIPENEM
EDE					
Butcher table	33.33%	66.67%	47.62%	4.76%	4.76%
Concrete slab	71.43%	61.90%	52.38%	23.81%	0.00%
Soil	66.67%	80.95%	38.10%	4.76%	9.52%
IFE					
Butcher table	38.71%	48.39%	32.26%	19.35%	9.68%
Concrete slab	61.11%	94.44%	38.89%	27.78%	38.89%
Soil	25.00%	75.00%	45.83%	8.33%	8.33%
IWO					
Butcher table	70.37%	70.37%	59.26%	29.63%	33.33%
Concrete slab	50.00%	90.00%	50.00%	25.00%	20.00%
Soil	77.27%	95.45%	59.09%	22.73%	4.55%
OSU					
Butcher table	100.00%	78.26%	69.57%	21.74%	13.04%
Concrete slab	86.96%	95.65%	65.22%	13.04%	4.35%
Soil	30.43%	82.61%	56.52%	0.00%	4.35%

Cell colour: green = low resistance, red = high

MDR % by Site_Code



Resistance % by Antibiotics



key insights

- Overall MDR is 40.15% (110 of 274 isolates), highest on concrete slabs (50%).
- Ceftazidime resistance is highest at 77%, and Imipenem lowest at 12.4%.
- Hotspots include OSU butcher table with Augmentin at 100%, and Ceftazidime above 94% at IFE concrete slab, IWO soil and OSU concrete slab.

Source: Mendeley Data (DOI 10.17632/ccmrx8n7mk.1). Organism not recorded; CLSI Enterobacterales breakpoints assumed.