

A STATISTICAL MODEL OF THE GLOBAL CARBON BUDGET

Mikkel Bennedsen
Aarhus Universitet
and CREATES

Eric Hillebrand
Aarhus Universitet
and CREATES

Siem Jan Koopman
Vrije Universiteit Amsterdam
and CREATES

20. MAY 2021



CREATES
INSTITUT FOR ØKONOMI
AARHUS UNIVERSITET

Econometric Models of Climate Change


DATA




www.globalcarbonproject.org

Friedlingsstein et al. (2020),
The global carbon budget 2020,
Earth System Science Data 12,
3269-3340


Δ




II




II




I




I



+




ANTHROPOGENIC EMISSIONS





$$\Delta E_t = \frac{0.14}{(0.02)} + X_t^E,$$

$$X_{t+1}^E = \frac{0.06}{(0.26)} X_t^E + \eta_{5,t}, \eta_{5,t} \sim N(0, \frac{0.024}{(0.032)})$$




$$\Delta E_t = \frac{0.11}{(0.01)} \Delta GDP_t - \frac{0.29}{(0.06)} I_{1991} + X_t^E,$$

$$X_{t+1}^E = \frac{0.39}{(0.12)} X_t^E + \eta_{4,t}, \eta_{4,t} \sim N(0, \frac{0.004}{(0.001)})$$




Friedlingstein et al. (2020)
Bennedsen, Hillebrand,
Koopman (2021)

LAND SINK




$$S_{LND_t} = \frac{-7.22}{(0.05)} + \frac{7.48}{(0.97)} \frac{C_t}{C_0}$$




$$S_{LND_t} = \frac{-6.70}{(0.04)} + \frac{7.12}{(0.49)} \frac{C_t}{C_0} + \frac{0.57}{(0.09)} SOI_t$$

Raupach et al. (2014), Raupach (2013), Gloor et al. (2010), Canadell et al. (2007)




"Moisture sensitivities of both productivity and decomposition are important for capturing the response of the net flux to such [La Niña] events." Haverd et al. (2018, p. 3013)

OCEAN SINK




$$S_{OCN_t} = \frac{-4.93}{(0.04)} + \frac{5.22}{(0.41)} \frac{C_t}{C_0}$$





$$S_{OCN_t} = \frac{-4.63}{(0.03)} + \frac{4.99}{(0.29)} \frac{C_t}{C_0} - \frac{0.06}{(0.02)} SOI_t$$

Raupach et al. (2014), Raupach (2013), Gloor et al. (2010), Canadell et al. (2007)



"When the winds are strongest during the cold cycle of ENSO deep upwelling occurs and [ocean CO₂ partial pressure] values are at a maximum." Feely et al. (1999, p. 599)

$\Delta C = \text{GROWTH IN ATM. CONCENTRATIONS}$



$$\Delta C_t = E_t - S_{LND}t - S_{OCN}t$$

THE DYNAMICS OF C

$$\Delta C_t = E_t - S_LND_t - S_OCN_t$$

$$= E_t - c_1 - c_2 - \beta_1^* C_t - \beta_2^* C_t + \varepsilon_t, \quad \varepsilon_t \sim I(0)$$

$$(1 + \beta_1^* + \beta_2^*)C_t - C_{t-1} = c + dt + x_t + \varepsilon_t$$

$$(1 - qL)C_t = qc + qdt + qx_t + q\varepsilon_t$$

$$\beta_i^* = \frac{\beta_i}{C_0} \approx 0.01$$

$$x_t = \sum_{i=1}^t X_i^E$$

$$q := \frac{1}{1 + \beta_1^* + \beta_2^*} \approx \frac{1}{1.02}$$

$$c = E_0 - c_1 - c_2$$

Three insights:

$$\begin{aligned} C_t &= q^t \left[C_0 - \frac{qc}{1-q} + \frac{dq^2}{(1-q)^2} \right] + \left[\frac{qc}{1-q} - \frac{dq^2}{(1-q)^2} \right] + \frac{dq}{1-q} t + \sum_{j=0}^{t-1} q^{j+1} x_{t-j} + \sum_{j=0}^{t-1} q^{j+1} \varepsilon_{t-j} \\ &= o(1) + O(1) + O(t) + I(1) + I(0) = O(t) + I(1) \end{aligned}$$

Thus,

$$\Delta C_t = I(0)$$

But,

$$(1 - qL)(1 - L)C_t = qd + q\Delta x_t + q\Delta \varepsilon_t = I(0)$$

THE SYSTEM MODEL

State equation Model 1

$$S_LND_{t+1}^* = c_1 + \frac{\beta_1}{C_0} C_{t+1}^*$$

$$S_OCN_{t+1}^* = c_2 + \frac{\beta_2}{C_0} C_{t+1}^*$$

$$E_{t+1}^* = E_t^* + d + X_t^E$$

$$C_{t+1}^* = C_t^* + G_ATM_{t+1}^*$$

$$G_ATM_{t+1}^* = E_{t+1}^* - S_{LND_{t+1}}^* - S_{OCN_{t+1}}^* + \beta_7 I1991$$

$$X_{1,t} = \phi_1 X_{1,t-1} + \eta_{1,t}$$

$$X_{2,t} = \eta_{2,t}$$

$$X_{3,t} = \phi_3 X_{3,t-1} + \eta_{3,t}$$

$$X_t^E = \phi_E X_{t-1}^E + \eta_{4,t}$$

State equation Model 2

$$S_LND_{t+1}^* = c_1 + \frac{\beta_1}{C_0} C_{t+1}^* + \beta_3 SOI_{t+1}$$

$$S_OCN_{t+1}^* = c_2 + \frac{\beta_2}{C_0} C_{t+1}^* + \beta_4 SOI_{t+1}$$

$$E_{t+1}^* = E_t^* + \beta_5 \Delta GDP_{t+1}^{World} + \beta_8 I1991 + X_t^E$$

Measurement equation

$$C_t = C_t^* + X_{1,t}$$

$$S_LND_t = S_LND_t^* + X_{2,t}$$

$$S_OCN_t = S_OCN_t^* + X_{3,t}$$

$$E_t = E_t^* + \beta_6 I1997$$

$$\begin{bmatrix} \eta_{1,t} \\ \eta_{2,t} \\ \eta_{3,t} \\ \eta_{4,t} \end{bmatrix} \sim N \left(0, \begin{bmatrix} \sigma_1^2 & r_{12}\sigma_1\sigma_2 & r_{13}\sigma_1\sigma_3 & 0 \\ r_{12}\sigma_1\sigma_2 & \sigma_2^2 & 0 & 0 \\ r_{13}\sigma_1\sigma_3 & 0 & \sigma_3^2 & 0 \\ 0 & 0 & 0 & \sigma_4^2 \end{bmatrix} \right)$$

$$\eta_{4,t} \sim N(0, \sigma_4^2 s_E^2 I_{t \geq 1996})$$

RESIDUAL DIAGNOSTICS

Model 1

Residual	mean	std dev	skew	kurt	JB	LB(1)	DW
C	-0.183	0.964	0.205	2.765	0.568	1.607	1.653
E	-0.033	1.009	-1.448	8.506	98.38***	0.073	2.060
S_LND	0.061	1.022	0.047	2.615	0.399	0.054	2.037
S_OCN	0.093	0.978	0.377	2.988	1.447	0.116	1.908

Model 2

Residual	mean	std dev	skew	kurt	JB	LB(1)	DW
C	-0.140	0.979	0.167	3.272	0.471	0.350	1.840
E	0.004	0.983	-0.269	3.095	0.759	0.062	1.928
S_LND	0.164	0.995	0.285	2.530	1.387	0.223	2.111
S_OCN	0.120	0.951	0.440	3.394	2.360	0.648	2.191

PARAMETER ESTIMATES

Model 2

	Coefficients		Variances		
	estimate	std err	estimate	std err	
c_1 (filt.)	-6.70	0.04	$\sigma_{\eta_1}^2$	0.66	0.13
c_2 (filt.)	-4.63	0.03	$\sigma_{\eta_2}^2$	0.32	0.06
β_1	7.12	0.49	$\sigma_{\eta_3}^2$	0.01	0.002
β_2	4.99	0.29	$\sigma_{\eta_4}^2$	0.004	0.001
β_3 (filt.)	0.57	0.09	r_{12}	-0.60	0.09
β_4 (filt.)	-0.06	0.02	r_{13}	-0.01	0.12
β_5 (filt.)	0.11	0.01	s_E	2.28	0.51
β_6 (filt.)	0.39	0.06			
β_7 (filt.)	-2.04	0.66			
β_8 (filt.)	-0.29	0.06			
ϕ_1	0.77	0.08			
ϕ_3	0.62	0.12			
ϕ_E	0.39	0.12			

$$C_{t+1}^* = C_t^* + G_ATM_{t+1}^*$$

$$S_LND_{t+1}^* = c_1 + \frac{\beta_1}{C_0} C_{t+1}^* + \beta_3 SOI_{t+1}$$

$$S_OCN_{t+1}^* = c_2 + \frac{\beta_2}{C_0} C_{t+1}^* + \beta_4 SOI_{t+1}$$

$$S_OCN_t = S_OCN_t^* + X_{3,t}$$

$$E_{t+1}^* = E_t^* + \beta_5 \Delta GDP_{t+1}^{World} + \beta_8 I1991 + X_t^E$$

$$E_t = E_t^* + \beta_6 I1997$$

$$G_ATM_{t+1}^* = E_{t+1}^* - S_LND_{t+1}^* - S_OCN_{t+1}^* + \beta_7 I1991$$

$$X_{1,t} = \phi_1 X_{1,t-1} + \eta_{1,t}$$

$$X_{2,t} = \eta_{2,t}$$


$$X_{3,t} = \phi_3 X_{3,t-1} + \eta_{3,t}$$

$$X_t^E = \phi_E X_{t-1}^E + \eta_{4,t}$$


$$\begin{bmatrix} \eta_{1,t} \\ \eta_{2,t} \\ \eta_{3,t} \\ \eta_{4,t} \end{bmatrix} \sim N \left(0, \begin{bmatrix} \sigma_1^2 & r_{12}\sigma_1\sigma_2 & r_{13}\sigma_1\sigma_3 & 0 \\ r_{12}\sigma_1\sigma_2 & \sigma_2^2 & 0 & 0 \\ r_{13}\sigma_1\sigma_3 & 0 & \sigma_3^2 & 0 \\ 0 & 0 & 0 & \sigma_4^2 \end{bmatrix} \right)$$

$$\eta_{4,t} \sim N(0, \sigma_4^2 s_E^2 I_{t \geq 1996})$$


BUDGET IMBALANCE



(A) Smoothed
 $-(\Delta X_1 + X_2 + X_3) + \beta_6 I(1997) - \beta_7 I(1991)$




(B) One-year ahead
 predictions




(C) Components
 $-\Delta X_1, -X_2, -X_3$

AIRBORNE FRACTION AND SINK RATE

Airborne Fraction



Sink Rate



$$AF = \frac{\Delta C}{E}$$


$$SR = \frac{S_{LND} + S_{OCN}}{C}$$

Raupach et al. (2014), Bennedsen, Hillebrand, Koopman (2019)


NOW-/FORECASTS

Forecasts of World GDP growth from IMF and World Bank


	2020	2021	2022
IMF	-3.5%	5.5%	4.2%
World Bank	-4.3%	4.0%	3.8%



(A) C - IMF




(B) G_ATM* - IMF




(C) E - IMF

Forecasts of SOI from forecast model of monthly SOI data 1866-2020, with trigonometric seasonal and second-order trigonometric cycle w/ period about 4 years



(D) S_LND - IMF



(E) S_OCN - IMF

PROJECTIONS TO 2050

Scenarios:

$\beta_{5,t}$ decreases linearly to 0 until 2050

No other abatement


GDP

- 1% annual growth
- 3.4% annual growth
- 5% annual growth


IPCC SR15 (2018)
UNFCCC NDC Synthesis Report (2021)

$$E_{t+1}^* = E_t^* + \beta_{5,t} \Delta GDP_{2010,t+1} + \beta_8 I1991 + X_t^E,$$


$$\beta_{5,t+1} = \beta_{5,t} + \eta_{5,t},$$




(A) E^*




(B) S_{LND}^*



(C) S_{OCN}^*



(D) G_{ATM}^*



(E) C^*

PROJECTIONS TO 2050


Scenarios:

$\beta_{5,t}$ decreases linearly to 0 until 2050


E_t decreases linearly to 0 until 2050

GDP


- 1% annual growth
- 3.4% annual growth
- 5% annual growth




(A) E^*




(B) S_{LND}^*




(c) S_{OCN}^*



(D) G_{ATM}^*



(E) C^*



(F) Emission reductions

$$E_{t+1}^* = E_t^* + \beta_{9,t} + \beta_{5,t} \Delta GDP_{2010,t+1} + \beta_8 I1991 + X_t^E,$$

$$\beta_{9,t+1} = \beta_{9,t} + \eta_{6,t},$$

CONCLUSIONS

- Specification of state-space model for Global Carbon Budget
- World GDP as driver in emissions
- Sinks: linear in CO₂ concentrations and in SOI
- CO₂ concentrations are I(1) ranging on I(2)
- Model allows for study of key variables such as airborne fraction and sink rate, forecasting, projecting necessary emission reductions

Future directions

- Include ensemble members for S_LND and S_OCN
- Factor model for drift in emissions using large macroeconomic dataset
- Higher resolution on Global Carbon Cycle module (MAGICC)
- Connection to temperatures (Energy Balance Models)
- Cointegration analysis



CREATEs
INSTITUT FOR ØKONOMI
AARHUS UNIVERSITET

REFERENCES

- Bacastow, R. and C.D. Keeling (1973), Atmospheric carbon dioxide and radiocarbon in the natural cycle: II. Changes from A.D. 1700 to 2070 as deduced from a geochemical model, in: Carbon and the biosphere conference proceedings, Brookhaven symposia in biology, Upton, N.Y., 86-135.
- Bennedsen, M., Hillebrand, E. and S.J. Koopman (2019), Trend analysis of the airborne fraction and sink rate of anthropogenically realease CO₂, *Biogeosciences* 12(8), 2565-2584
- Bennedsen, M. Hillebrand, E., and S.J. Koopman (2021), Modeling, forecasting, and nowcasting U.S. CO₂ emissions using many macroeconomic predictors, *Energy Economics*, 96, 105118
- Canadell, J.G., et al. (2007), Contributions to accelerating atmospheric CO₂ growth from economic activity, carbon intensity, and efficiency of natural sinks, *PNAS* 104(47), 18866-18870
- Friedlingstein, P., et al. (2020), Global Carbon Budget 2020, *Earth System Science Data* 12(4), 3269-3340
- Gifford, R.M. (1993), Implications of CO₂ effects on vegetation for the global carbon budget, in M. Heimann (ed.): The global carbon cycle, Springer, 159-199
- IPCC SR15 (2018), Special report: Global warming of 1.5 degree C
- Joos, F., et al. (1996), An efficient and accurate representation of complex oceanic and biospheric models of anthropogenic carbon uptake, *Tellus B* 48(3), 397-417
- Joos, F. et al. (2001), Global warming feedbacks on terrestrial carbon uptake under the IPCC emissions scenarios, *Global Biogeochemical Cycles* 15(4), 891-907
- Raupach, M. (2013), The exponential eigenmodes of the carbon-climate system, and their implications for ratios of responses to forcings, *Earth System Dynamics* 4, 31-49
- Raupach, M., et al. (2014), The declining uptake rate of atmospheric CO₂ by land and ocean sinks, *Biogeosciences* 11(13), 3452-3475
- Gloor, M., et al. (2010), What can be learned about carbon cycle climate feedbacks from the CO₂ airborne fraction?, *Atmospheric Chemistry and Physics* 10, 7739-7751
- Stock, J.H. and M. Watson (1996), Evidence on structural instability in macroeconomic time series relations, *Journal of Business and Economic Statistics* 14(1), 11-30
- UNFCCC (2021), United Nations Framework Convention on Climate Change: National determined contributions synthesis report