

Review

Calpains in muscle wasting

Marc Bartoli, Isabelle Richard*

Généthon, Centre National de la Recherche Scientifique UMR 8115, 1 bis rue de l'Internationale, 91000 Evry, France

Received 8 October 2004; received in revised form 1 December 2004; accepted 31 December 2004

Abstract

Calpains are intracellular nonlysosomal Ca^{2+} -regulated cysteine proteases. They mediate regulatory cleavages of specific substrates in a large number of processes during the differentiation, life and death of the cell. The purpose of this review is to synthesize our current understanding of the participation of calpains in muscle atrophy. Muscle tissue expresses mainly three different calpains: the ubiquitous calpains and calpain 3. The participation of the ubiquitous calpains in the initial degradation of myofibrillar proteins occurring in muscle atrophy as well as in the necrosis process accompanying muscular dystrophies has been well characterized. Inactivating mutations in the calpain 3 gene are responsible for limb-girdle muscular dystrophy type 2A and calpain 3 has been found to be downregulated in different atrophic situations, suggesting that it has to be absent for the atrophy to occur. The fact that similar regulations of calpain activities occur during exercise as well as in atrophy led us to propose that the calpains control cytoskeletal modifications needed for muscle plasticity.

© 2005 Elsevier Ltd. All rights reserved.

Keywords: Ubiquitous calpains; Calpain 3; Muscle atrophy

Contents

1. The family of calpains	2116
2. Overview of muscle atrophy	2117
3. Confirmed and putative implications of the ubiquitous calpains in muscle atrophy	2118
3.1. Protein degradation	2118
3.2. Protein synthesis	2119
3.3. Cellular phenomena	2121
3.3.1. Necrosis	2121
3.3.2. Apoptosis	2122
3.3.3. Impairment in the regenerative capacity of the muscle	2122
3.4. Conclusion on ubiquitous calpains	2123

* Corresponding author. Tel.: +33 1 69 47 29 38; fax: +33 1 60 77 86 98.

E-mail address: richard@genethon.fr (I. Richard).

4.	The unique case of calpain 3	2124
4.1.	Inherited calpain 3 deficiency: limb-girdle muscular dystrophy type 2A	2124
4.2.	Downregulation of calpain 3 in atrophic conditions	2125
4.3.	Conclusion on calpain 3	2125
5.	General conclusion	2125
	Acknowledgment	2126
	References	2126

1. The family of calpains

The calpain family is comprised of 14 members of nonlysosomal Ca^{2+} -activated cysteine proteases showing various patterns of expression (for review, see Goll, Thompson, Li, Wei, & Cong, 2003). Muscle tissue expresses mainly three distinct calpains: the ubiquitous calpains 1 and 2 (also called mu- and m-) which are the best-characterized calpains and calpain 3 (also called p94) which is highly expressed in this tissue. The canonical structure of a calpain is composed of four distinct domains (Fig. 1). The N-terminal region presents no homology to domains identified in other proteins. Domain II is structurally similar to the catalytic domain of other cysteine proteases, particularly around the amino acids forming the catalytic triad and can be divided into two globular subdomains forming a catalytic cleft (Moldoveanu, Campbell, Cuerrier, & Davies, 2004; Reverter, Strobl, Fernandez-Catalan,

Sorimachi, Suzuki, & Bode, 2001). Domain III possesses a three-dimensional topology very similar to that of C2 domains, which are involved in protein–protein interaction and membrane targeting, and are regulated by Ca^{2+} through binding of this ion to an acidic loop (Tompa, Emori, Sorimachi, Suzuki, & Friedrich, 2001). Domain IV is a Ca^{2+} -binding domain structurally similar to calmodulin apart from the fact that it contains 5 EF-hands instead of four as in calmodulin (Babu, Bugg, & Cook, 1988; Cygler, Grochulski, & Blanchard, 2002; Minami, Emori, Kawasaki, & Suzuki, 1987). Calpains 1 and 2 associate non-covalently with a small subunit containing one domain rich in hydrophobic residues (domain V) and a domain similar to domain IV (domain VI) (Lin et al., 1997). Calpains 1 and 2 have different in vitro calcium sensibility, in the range of 5–50 μM for calpain 1 and 250–1000 μM for calpain 2 (Elce, Hegadorn, & Arthur, 1997). Calpain 3 possesses also the classical structure of a calpain except that it carries three unique sequences not found in any other calpains: NS at the N-terminus, IS1 within the catalytic domain and IS2 upstream of the Ca^{2+} -binding domain (Sorimachi et al., 1989) (Fig. 1). The latter two sequences confer specific properties on calpain 3: IS1 possesses autolytic sites and IS2 comprises a nuclear localization signal and a binding site to titin (also called connectin), a giant elastic protein of the sarcomere (Sorimachi et al., 1989, 1995, 1993).

Since proteolytic activities are potentially deleterious to the cell, calpains are highly regulated and are supposed to be in an inactivate state most of the time (Goll et al., 2003; Taveau, Bourg, Sillon, Roudaut, Bartoli, & Richard, 2003). Ca^{2+} spikes, removal of the N-terminus region by autolysis, phosphorylation and association with membrane, the small subunit or calpastatin (the only known endogenous protein inhibitor specific for ubiquitous calpains), all seem

Ubiquitous calpains

Large subunit of calpains 1 and 2

Small subunit of calpains 1 and 2 (calpain 4)

Calpain 3




Fig. 1. Organization of the ubiquitous calpains, the small subunit and calpain 3. The coordinates were determined by structural analysis for calpain 2 and predicted by sequence for the small subunit (also called calpain 4, see web site: <http://ag.arizona.edu/calpains>) and calpain 3. The 3 domains unique to calpain 3 are shown in red. NLS: nuclear localization sequence.

to be involved in the regulation of ubiquitous calpain activity. Controversial issues about their respective participation are reviewed in Goll et al. (2003). However, crystallographic studies have unraveled the structural basis of calpain activation by Ca^{2+} (Hosfield, Elce, Davies, & Jia, 1999; Strobl et al., 2000). In its absence, the residues forming the catalytic site are too far apart to form an active site due to the presence of salt bridges with other parts of the molecule. Upon Ca^{2+} binding to the EF-hands in the last domain and to the acidic loop in the C2-like domain, these salt bridges are disrupted allowing the two globular domains to come closer together. Further Ca^{2+} binding at both sides of the catalytic cleft completes the process, resulting in the formation of a fully active site (Moldoveanu, Hosfield, Lim, Elce, Jia, & Davies, 2002). Even though calpain 3 possesses Ca^{2+} -binding sites, it has a lower Ca^{2+} requirement for its activation (Branca, Gugliucci, Bano, Brini, & Carafoli, 1999). A supplementary regulatory mechanism seems to exist in the form of the internal propeptide, IS1, that has to be removed by autolysis for full activation (Diaz, Moldoveanu, Kuiper, Campbell, & Davies, 2004; Taveau et al., 2003).

Within the cell, the calpain system covers a broad range of physiological functions, most of them not fully understood. These functions include proteolysis of proteins involved in the cell cycle, apoptosis, cytoskeleton organization and signal transduction (Goll et al., 2003). The purpose of this review is to synthesize our current understanding of the participation of calpains in skeletal muscle atrophy since many of these biological functions could be involved in this condition. Because of their demonstrated implication in muscle protein breakdown and the huge amount of literature, this review focuses on ubiquitous calpains and on calpain 3. The role of other calpains found in muscle such as calpain 10 will not be discussed because there is no data to date that implicate these calpains in muscle wasting.

2. Overview of muscle atrophy

Skeletal muscle atrophy corresponds to a loss of muscle tissue that can include a hypotrophic component (a decrease in fiber size) and/or a hypoplastic component (a reduction in the numbers of fibers). Muscle

atrophy is associated with fasting, disuse, ageing and with several pathological conditions such as injury, sepsis, diabetes mellitus, AIDS and cancer (Childs, 2003). In particular, muscle atrophy is part of a multisystemic perturbation, known as cachexia, in critical illnesses. Muscle atrophy is also seen in genetically inherited myopathies and neuropathies (Anderson, Potter, Baban, & Davies, 2003; Stubgen, 1994; Walling, 1999).

The trigger of muscle wasting can be a circulating factor such as cytokines, of which the most important are tumor necrosis factor- α (TNF- α) and interleukin 6 (IL-6), catabolic hormones such as myostatin or glucocorticoids or a low level of anabolic hormones such as insulin or insulin-like growth factor-1 (IGF-1) (reviewed by Jackman & Kandarian, 2004). In these cases, the responses extend to muscles throughout the body. Atrophy can also arise by suppression of mechanical loading due to disuse or denervation and, in this case, the phenomenon occurs locally (Baldwin & Haddad, 2002). Increased Ca^{2+} permeability observed in muscular dystrophies or increase in free radicals as seen in ageing are also signals relevant for initiation of muscle wasting (Spencer, Croall, & Tidball, 1995; Weindruch, 1995).

These triggering signals activate different signaling pathways, some of which are directly linked to an atrophic program and others involved in various aspects of cell homeostasis. It is beyond the scope of this review to cover these points in details and the reader is invited to refer to recent reviews (Jackman & Kandarian, 2004; Kandarian & Stevenson, 2002; Sartorelli & Fulco, 2004). Nevertheless, it is worth mentioning that two E3 ubiquitin ligases, Muscle Ring Finger-1 protein (MuRF1) and Muscle Atrophy F-box (MAFbx), are invariably found upregulated in atrophic conditions, suggesting that they are part of a common response to atrophy (Bodine et al., 2001; Glass, 2003; Jagoe, Lecker, Gomes, & Goldberg, 2002; Lecker et al., 2004).

These signals will eventually lead to muscle atrophy through various mechanisms working in sequence or in parallel in the fibers: increase in protein degradation, decrease in gene expression leading to an imbalance of muscle protein amount, loss of fibers by apoptosis or necrosis and inability of satellite cells to counteract the decrease in fiber size and number (Glass, 2003; Jackman & Kandarian, 2004; Jejurikar & Kuzon, 2003).

3. Confirmed and putative implications of the ubiquitous calpains in muscle atrophy

Evidence that ubiquitous calpains are key players in atrophy is based on sets of data obtained by expression analysis, detection of activity of calpain and the use of calpain inhibitors. First of all, several studies have established that calpains are elevated in atrophic conditions like disuse, denervation, glucocorticoid treatment and sepsis (Haddad, Roy, Zhong, Edgerton, & Baldwin, 2003; Hong & Forsberg, 1995; Tang, Cheung, Ip, & Ip, 2000; Voisin et al., 1996; Williams, Decourten-Myers, Fischer, Luo, Sun, & Hasselgren, 1999). These data were not sufficient to definitively establish a role for calpains in muscle wasting because increases in RNA or/and protein expression are not necessarily correlated with increased activity since the net activity of calpains depends on the calpain/calpastatin ratio (Costelli, Tullio, Baccino, & Melloni, 2001; Wei, Zhou, Wang, & Schwartz, 2000).

Another argument in favor of the involvement of ubiquitous calpains in muscle wasting is based on the analysis of active calpains (Goll et al., 2003). Monitoring of calpain activity has been performed by various techniques such as degradation of fluorescent peptides, both in cells or in vitro, Western blot analysis revealing the shift in size of active calpains compared to inactive calpains or by analysis of substrate degradation. These studies demonstrated an increased activity in disuse, sepsis, fasting and muscular dystrophy (Alderton & Steinhardt, 2000b; Baker & Margolis, 1987; Huang & Forsberg, 1998; Purintrapiban, Wang, & Forsberg, 2003; Spencer et al., 1995; Voisin et al., 1996; Williams et al., 1999). One should be aware that disruption of muscle may non-specifically activate calpains, that, even if calpains are able to hydrolyze peptide in vitro, the natural substrate may not be accessible and that these tests do not necessarily take into account the regulation by association with inhibitors like calpastatin. Recently, several teams have tried to bypass these drawbacks by developing various assays based on fluorescence resonance energy transfer (Ono et al., 2004; Vanderklish, Krushel, Holst, Gally, Crossin, & Edelman, 2000). Using this technology combined with multiphoton microscopy, we were able to follow calpain activation in living animals, opening the way to future studies on the roles of calpains in pathophysiological conditions such as atrophy (Stockholm,

Bartoli, Sillon, Bourg, Davoust, & Richard, 2005).

Other data came from studies inhibiting calpains under atrophic conditions (Badalamente & Stracher, 2000; Whitehouse, Smith, Drake, & Tisdale, 2001). Unfortunately, the inhibitors used in these studies (like leupeptin and E-64) could also inhibit cathepsins and other cysteine proteases. Their lack of specificity for calpain prevents a clear confirmation that calpains are involved in muscle wasting (Donkor, 2000). Opportunely, the availability of transgenic mice overexpressing calpastatin confirmed that calpains are involved in muscle wasting (Tidball & Spencer, 2002). These transgenic animals exhibited a lower susceptibility to wasting by immobilization and these transgenic mice crossed with *mdx* mice had fewer fibers in regeneration, suggesting a reduction in necrosis (Spencer & Mellgren, 2002; Tews, 2002; Tidball & Spencer, 2002).

While the role of ubiquitous calpains in atrophy now seems unquestionable, it is difficult to distinguish the relative participation of each of the calpain isoforms. They may well have different roles in the development of atrophy since they have different activation signals and may be localized in distinct parts of the cell (see Goll et al., 2003 for a review). Hopefully, as illustrated in recent studies, the availability of siRNA specific for each calpain isoform is going to be a powerful tool to dissect their relative role in various atrophic conditions (Franco, Perrin, & Huttenlocher, 2004; Honda et al., 2004).

In the following chapters, we will briefly summarize the current knowledge about the different mechanisms intervening in atrophy before discussing the participation of the ubiquitous calpains in these phenomena.

3.1. Protein degradation

Protein degradation in muscle wasting has been shown by measuring either overall protein degradation using radioactive tracers or myofibrillar protein degradation by the excretion of 3-methylhistidine, a molecule obtained through a post-translational modification of histidine and found mostly in myosin and actin (Elia, Carter, Bacon, Winearls, & Smith, 1981; Huang & Forsberg, 1998; Johnson & Perry, 1970). Other studies have observed the concomitant disappearance of myofibrillar proteins and sarcomere disorganization at histological, immunohistochemical and electron

microscopical levels (Belozeroval, Shenkman, Mazin, & Leblanc, 2001; Massa, Carpenter, Holland, & Karpati, 1992). One type of evidence came from studies carried out by Forsberg's team. They used several strategies to modulate intracellular calpain activities: downregulation of calpain expression by use of antisense oligonucleotide, overexpression of a dominant-negative form of calpain 2 and overexpression of the calpastatin inhibitory domain. Under conditions of atrophy, inhibition of calpains decreased total protein degradation. Western blot analysis of the treated cells showed that inhibition of calpain 2 stabilized desmin, nebulin and fodrin. They conclude that calpains are involved in the disassembly of sarcomeric protein in atrophic models (Huang & Forsberg, 1998; Purintrapiban et al., 2003; Xiao, Wang, Purintrapiban, & Forsberg, 2003).

Numerous studies have evaluated the relative contribution of the three main cell proteolytic systems in protein degradation: the lysosomal enzymes, the Ca^{2+} -dependent calpains and the ATP-dependent ubiquitin-proteasome system (Costelli & Baccino, 2003; Furuno, Goodman, & Goldberg, 1990; Huang & Forsberg, 1998; Jackman & Kandarian, 2004). Increased lysosomal enzyme activity was observed in various atrophic conditions but does not seem to contribute significantly to cytosolic protein degradation (Fujita et al., 1996; Voisin et al., 1996). The two other systems are thought to work sequentially in muscle proteolysis during atrophy (Hasselgren & Fischer, 2001; Jackman & Kandarian, 2004).

Several studies have shown that the proteasome is not able to degrade intact myofibrils and that muscle proteins such as actin and myosin are released from the sarcomere by a Ca^{2+} /calpain-dependent mechanism before they undergo ubiquitination and degradation by the proteasome (Koochmaraie, 1992a; Solomon & Goldberg, 1996; Williams et al., 1999). This sequence of events is supported by the observation that fragments are still released from the sarcomere in ATP-depleted post-mortem muscle in which proteasome-dependent proteolysis cannot occur (Koochmaraie, 1992b). Moreover, in calpastatin overexpressing animals, this post-mortem degradation is reduced, confirming the involvement of calpains (Kent, Spencer, & Koochmaraie, 2004). Calpains can therefore be described as the initiators of myofibrillar degradation and the proteasome as the cleaner

removing all myofibrillar fragments to recycle amino acids.

In muscle wasting, the preferential site of ubiquitous calpain proteolysis seems to be the Z-disc, a dense multi-proteinaceous network located on both sides of the sarcomeres. First, the Z-disc is one of the possible subcellular localizations of ubiquitous calpains, at least of calpain 1, as it has been detected at this location by immunogold electron microscopy (Yoshimura, Murachi, Heath, Kay, Jasani, & Newman, 1986) and it binds to α -actinin, a major constituent of Z-discs (Ohtsuka, Yajima, Maruyama, & Kimura, 1997; Raynaud et al., 2003). Second, many calpain substrates associate with the Z-disc including titin, nebulin, filamin, troponin-T and the intermediate filament desmin that attaches the sarcolemma to the Z-disc. These data suggest that calpains could process proteins within the Z-disc directly and could participate in the disorganization of Z-discs observed by electron microscopy in experimental wasting conditions (Huff-Lonergan, Mitsuhashi, Beekman, Parrish, Olson, & Robson, 1996; Taylor, Geesink, Thompson, Koochmaraie, & Goll, 1995; Williams et al., 1999). When Z-discs are almost completely disrupted, actin and myosin are passively released in the cytoplasm together with other proteins from the sarcomere (Bullard, Sainsbury, & Miller, 1990; Tidball & Spencer, 2002).

In addition, ubiquitous calpains may indirectly influence the rate of protein degradation. First, it is possible that calpain cleavages facilitate their degradation by producing N-terminal extremities compatible with the N-end rule of ubiquitin-proteasome-dependent degradation of otherwise stable proteins (Hasselgren, 1999; Solomon, Baracos, Sarraf, & Goldberg, 1998a; Solomon, Lecker, & Goldberg, 1998b). According to this rule, the life span of a protein could be determined by its very N-terminal amino acid (a basic or a bulky hydrophobic amino acid boosts the degradation). Second, accumulation of myofibrillar protein fragments may generate a positive feedback resulting in an increase in proteasome activity by the stabilizing association of E3-ubiquitin ligases with their substrates (Li, Gazdaru, Pan, & Fuchs, 2004).

3.2. Protein synthesis

Another hallmark of muscle wasting is an alteration of protein synthesis. A selective downregulation

of transcripts from specific proteins corresponding either to myofibrillar proteins such as myosin and actin or to regulatory proteins such as the myogenic regulatory factors, myocyte enhancer factor 2C (MEF2C) and MyoD or cyclin D1, a substrate of calpains, has been observed in atrophy (Acharyya et al., 2004; Haddad et al., 2003; Yamakuchi et al., 2000). Since actin and myosin account for 40% of the total muscle protein content, even a slight decrease in their expression will have a profound impact in the total amount of protein. A general reduction in translational efficiency has also been reported (Haddad et al., 2003). The origin of this phenomenon could be due either to a decrease in ribosomal RNA, to an inactivation of factors implicated in translational initiation such as eukaryotic initiation factor 4E-binding protein 1 (4EBP-1) and ribosomal

protein S6 kinase (p70^{S6K}) or to an energy deficit due to a loss of mitochondria or to a decrease in nitrogen and glucose supply (Haddad et al., 2003; Heck & Davis, 1988; Hornberger, Hunter, Kandarian, & Esser, 2001; Jagoe et al., 2002).

To date, intervention of ubiquitous calpains has not been associated with any of these mechanisms in muscle atrophy. However, considering that calpains are well known to modulate several signaling pathways, some of which function as key mediators in atrophy (Fig. 2), there is a possibility that they may participate, one way or another, in the regulation of gene expression in atrophy. In particular, ubiquitous calpains are renowned for inducing limited cleavage of transcription factors, modulating their transcriptional activity (Hirai, Kawasaki, Yaniv, & Suzuki, 1991; Watt &




Fig. 2. Diagram showing some of the signaling pathways involved in atrophy. This diagram represents grossly the signaling pathways that transduce signals from cytokines (TNF- α /IL-6) and mechanical signals from both the sarcomere and the membrane (integrin). These pathways control, by the intermediary action of transcription factors, the expression of genes coding for myofibrillar proteins, the proteasome and apoptosis components. Proteins cleaved by calpains, which participate in this regulation, are boxed. Arrows indicate links between pathways. They do not necessarily indicate activation.

Molloy, 1993). Among the transcription factors known to be linked to the control of muscle mass and which are substrates of ubiquitous calpains, one can cite the signal transducer and activator of transcription family (STAT) which acts downstream of IL-6 receptor, a pathway known to be involved in cachexia and the nuclear factor- κ B (NF- κ B) family, transcription factors whose activity seems to be required for disuse and cytokine-induced muscle atrophy (Oda, Wakao, & Fujita, 2002; Walowitz, Bradley, Chen, & Lee, 1998). In addition, actors in signaling pathways, which are altered in some atrophies, are also known to be modified by calpain cleavage (the protein kinase C (PKC), calcineurin and Cdc42/RhoA, for example) (Chockalingam, Cholera, Oak, Zheng, Jarrett, & Thomason, 2002; Sneddon, Delday, & Maltin, 2000; Wang, Roufogalis, & Villalobo, 1989). Thus, signals inducing atrophy may be well relayed or interrupted in the cell by calpain activity. Therefore, it is possible that new concepts concerning calpain regulation of signaling pathways in muscle wasting will emerge in the near future.

3.3. Cellular phenomena

In muscle atrophy, several phenomena occur or can occur at the cellular level: (i) loss of myofibers by necrosis or apoptosis leading to hypoplasia, (ii) loss of myofiber segments by an apoptotic mechanism in order to adjust the nuclear domain to the reduced size of the fiber, (iii) decrease in the number of satellite cells by apoptosis and (iv) impairment of the regenerative capacity of the muscle (Allen, Linderman, Roy, Grindeland, Mukku, & Edgerton, 1997; Dupont-Versteegden, Murphy, Houle, Gurley, & Peterson, 1999; Hikida, Van Nostran, Murray, Staron, Gordon, & Kraemer, 1997; Mitchell & Pavlath, 2004).

3.3.1. Necrosis

Fiber loss by necrosis could engender muscle hypoplasia and is essentially linked to cell membrane damage, osmotic imbalance or ion fluxes that overwhelm the vital capacity of the fibers. Necrosis accounts for destructive changes associated with muscular dystrophies such as dystrophin or sarcoglycan deficiencies (Karpati, Hilton-Jones, & Griggs, 2001). Those proteins are associated in a complex linking the cytoskeleton to the extracellular matrix and protect the membrane against mechanical damage during muscle

contraction (for reviews, see Cohn & Campbell, 2000; Lapidus, Kakkar, & McNally, 2004). One of the functional repercussions of the absence of these proteins is an increase of Ca^{2+} influx resulting in an elevation of intracellular Ca^{2+} levels, at least at the submembranous level (Bertorini, Bhattacharya, Palmieri, Chesney, Pifer, & Baker, 1982; Gillis, 1999; Hopf, Turner, Denetclaw, Reddy, & Steinhardt, 1996; Jackson, Jones, & Edwards, 1985; Mongini et al., 1988; Robert et al., 2001; Turner, Fong, Denetclaw, & Steinhardt, 1991; Tutdibi, Brinkmeier, Rudel, & Fohr, 1999). Recent studies have associated this Ca^{2+} elevation with an increased permeability of the plasma membrane through leaky cationic channels, in particular of the transient receptor potential channel (TRP) family (Alderton & Steinhardt, 2000b; Iwata, Katanosaka, Arai, Komamura, Miyatake, & Shigekawa, 2003; Turner, Schultz, Ganguly, & Steinhardt, 1993; Vandebrouck, Martin, Colson-Van Schoor, Debaix, & Gailly, 2002). Several lines of evidence support the fact that this increase in Ca^{2+} induces a deregulated and sustained activation of ubiquitous calpains and that this participates in the pathology. Abnormal increase in calpain expression, increase in calpain activity and increase in protein degradation, all have been observed in dystrophic muscles (Alderton & Steinhardt, 2000b; Combaret, Taillandier, Voisin, Samuels, Boespflug-Tanguy, & Attaix, 1996; MacLennan, McArdle, & Edwards, 1991; Neerunjun & Dubowitz, 1979; Reddy, Anandavalli, & Anandaraj, 1986; Spencer et al., 1995; Tidball, Albrecht, Lokensgard, & Spencer, 1995; Tidball & Spencer, 2000; Turner, Westwood, Regen, & Steinhardt, 1988). These observations indicate a correlation with the course of the pathology. Confirmation came from the use of calcium channel blockers that were shown to inhibit the activation of calpains and to reduce the pathological signs and from the fact that mice transgenic for calpastatin have a reduced level of necrosis (Spencer & Mellgren, 2002). The relationship between ubiquitous calpains and fiber death may be directly due to increased and unregulated proteolysis but could also come from a perturbation of signaling pathways regulated by calpains.

It has been postulated that an initial influx of Ca^{2+} occurs at membrane wounds caused by mechanical stresses in the dystrophic fibers, that this flux activates calpains which, in turn, induces an increased opening of Ca^{2+} channels, possibly directly by proteolysis. Thus,

a positive feedback loop increasing Ca^{2+} influx could take place (Alderton & Steinhardt, 2000a; McCarter & Steinhardt, 2000). An increase in Ca^{2+} occurs also during muscle contraction, reaching the micromolar range (Berchtold, Brinkmeier, & Muntener, 2000). In this physiological condition, ubiquitous calpains seem to be tightly regulated to prevent unwanted damages whereas in dystrophic muscles, they seem to escape these regulations. Interestingly, calpastatin has been reported to be degraded in dystrophic muscles and was reported to be a substrate of calpains (De Tullio, Aversa, Salamino, Pontremoli, & Melloni, 2000; Nakamura, Imajoh-Ohmi, Suzuki, & Kawashima, 1991; Pontremoli, Melloni, Viotti, Michetti, Salamino, & Horecker, 1991).

3.3.2. Apoptosis

Apoptosis is a phenomenon that can participate in atrophy by leading to loss of myofibers (hypoplasia) or loss of myofiber segments (hypotrophy). Apoptosis is a genetically programmed form of cell death that can be triggered through death receptors such as the TNF-receptor or via mitochondrial pathways (Rossi & Gaidano, 2003). Initial stimuli are followed by a cascade of proteolytic events mediated by caspases and leading eventually to disassembly of architectural components of the cell and DNA fragmentation. It should be noted that a cell could die by necrosis even if the primary signal was an apoptotic one. In conditions where ATP is severely depleted, the energy-dependent apoptotic program cannot be pursued and necrosis occurs (Tidball et al., 1995). Apoptosis has been observed in dexamethasone-mediated atrophy, denervation, hind limb suspension and with ageing (Leeuwenburgh, 2003; Tews, 2002).

Increase in apoptotic cells external to the atrophied fibers could account for the reduced number of satellite cells seen in atrophy (Mitchell & Pavlath, 2004). In addition, in the atrophying fibers, nuclei are removed by an apoptotic-like pathway to maintain the same ratio between the number of nuclei and the volume of the fibers, with each nucleus controlling a defined volume known as the nucleus domain (Allen et al., 1997; Hall & Ralston, 1989). With the exception of the upregulation of Bax and Bcl-2 seen in denervated rat muscles which is indicative of a mitochondria-directed apoptosis, the mechanisms of apoptosis in muscle atrophy are not currently well-defined, nor is the possible participation of calpains in this phenomenon (Olive &

Ferrer, 2000). The only relationship between ubiquitous calpains and apoptosis in atrophy known to date is that induction of apoptosis by dexamethasone is dependent on calpain activation (Squier, Miller, Malkinson, & Cohen, 1994) and that some apoptosis-related proteins such as p53, cain/cabin1 and caspase 3 are substrates of ubiquitous calpains (Bizat et al., 2003; Kim et al., 2002; Piechaczyk, 2000). However, it is not known whether these cleavages could occur during atrophy or not.

3.3.3. Impairment in the regenerative capacity of the muscle

Muscle tissue possesses an intrinsic regenerative capacity thanks to the presence of satellite cells situated under the basal lamina (Bischoff, 1986). Those cells are normally quiescent but can be activated to regenerate the muscle. In muscle atrophy, the decrease in size (due to protein degradation and reduced protein synthesis) and/or number of myofibers (due to necrosis or apoptosis) is not compensated by this process. This impairment in the regenerative capacity of the muscle could be the result of a decrease in number of satellite cells or/and an insufficient potential of these cells to proliferate and fuse with existing myofibers. The former event has been observed in denervation, advanced age and muscular dystrophy (Hikida et al., 1997; Irintchev, Zwyer, & Wernig, 1997; Luz, Marques, & Santo Neto, 2002; Ontell, Hughes, Hauschka, & Ontell, 1992; Renault, Thornell, Eriksson, Butler-Browne, Mouly, & Thorne, 2002; Rodrigues Ade, Geuna, Rodrigues, Silva, & Aragon, 2002). For example, studies performed on *mdx* model mice have shown that after repeated cycles of degeneration–regeneration induced by toxins, muscle failed to regenerate because of the depletion of satellite cells (Luz et al., 2002). It has also been demonstrated that satellite cells from chronically denervated muscles have an increased susceptibility to apoptosis (Jejurikar, Marcelo, & Kuzon, 2002; Tews, 2002). The regenerative capacity of satellite cells can be negatively influenced by a reduction of anabolic factors and an increase of catabolic factors associated with atrophic conditions (McCroskery, Thomas, Maxwell, Sharma, & Kambadur, 2003; Melone, Peluso, Petillo, Galderisi, & Cotrufo, 1999; Stewart, Newcomb, & Holly, 2004). Finally, it has been shown that satellite cells isolated from immobilized muscles have impaired proliferative potential and are less effective in forming

myotubes in vitro (Jejurikar et al., 2002; Mitchell & Pavlath, 2004).

Modulation of ubiquitous calpain activity may have a place in the cellular events (impairment of proliferation, migration and fusion of satellite cells) associated with atrophy. Several lines of evidence point to a role of ubiquitous calpains in regulating cell proliferation through action at different points of the cell cycle. Relocation of ubiquitous calpains into the nuclear compartment has been observed during the cell cycle in various cell types (Choi et al., 1997; Zhang, Lu, Xie, & Mellgren, 1997). For instance, calpain 2 translocates into the nucleus during satellite cell proliferation both in cultured cells and in muscle, whereas it is located in the cytoplasm in quiescent cells (Raynaud, Carnac, Marcilhac, & Benyamin, 2004). Moreover, several studies using specific calpain inhibitors and/or in vitro assays with recombinant calpains established that ubiquitous calpains have an effect on cell cycle proteins such as cyclin D1, p53 or on the cell cycle itself, mainly in G1 (Choi et al., 1997; Zhang et al., 1997). Finally, downregulation of calpain 2 using RNA interference demonstrated that it plays a role in chromosome alignment on the metaphase plate during mitosis (Honda et al., 2004).

There is also compelling evidence that ubiquitous calpains are involved in migration and fusion of myogenic cells. Migration is an important step for myoblast alignment before fusion. Transfecting myoblast cells with calpastatin or with an antisense oligomer against ubiquitous calpains greatly reduces their migration (Dedieu, Mazeret, Poussard, Brustis, & Cottin, 2003). In addition, numerous studies have implicated ubiquitous calpains in cell mobility in other cell types and have led to a model in which calpain 2 regulates detachment at the rear of migrating cells and calpain 1 promotes adhesions at the leading edge (Glading, Lauffenburger, & Wells, 2002). Furthermore, many signaling and cytoskeletal proteins of the cell-matrix attachment sites such as talin, ezrin, focal adhesion kinase, integrin, filamin, vinculin and paxillin, have been identified as calpain substrates (Carragher, Levkau, Ross, & Raines, 1999; Dedieu et al., 2004; Goll et al., 2003). It has also been shown that calpain 2 activation at the rear edge of the cell is dependent on growth factors, suggesting that in atrophy, a trophic factor depleted condition, migration could be impaired.




Fig. 3. Possible action of ubiquitous calpains in atrophy. Upon receiving an activation signal, calpains could play their role in muscle wasting in numerous ways. Activated calpains can degrade sarcomeric proteins, control the expression of several genes by proteolysis of signaling proteins and transcription factors and also may participate in fiber necrosis. In muscle progenitor cells, calpains could control the cell cycle, migration and fusion.

Myoblast fusion, a Ca^{2+} -dependent process, requires an extensive reorganization of the cytoskeleton for redistribution of membrane proteins and leaving free patches of the lipid bilayer for the cells to detach from the extracellular matrix. Use of calpain inhibitors such as calpastatin (Barnoy, Zipser, Glaser, Grimberg, & Kosower, 1999; Kwak et al., 1993; Temm-Grove, Wert, Thompson, Allen, & Goll, 1999) or of antisense oligomers against calpain 2 (Balcerzak et al., 1995) completely prevents fusion while use of antisense oligomers against calpastatin leads to earlier and faster myoblast fusion (Balcerzak, Cottin, Poussard, Cucuron, Brustis, & Ducastaing, 1998). In addition, it has been shown that, in myoblasts, calpain 2 translocates from the cytosol to the membrane and that the calpastatin level decreases at the time of cell fusion. Altogether, these data point to an essential role of calpains, at least calpain 2, in myoblast fusion (Barnoy, Glaser, & Kosower, 1997).

3.4. Conclusion on ubiquitous calpains

The proven and putative actions of the ubiquitous calpains on processes occurring during muscle atrophy are summarized in Fig. 3. The ubiquitous calpains are undoubtedly involved in the initial breakdown of

myofibrils observed in atrophy and in the necrosis of fibers observed in muscular dystrophies. Besides these actions, they could have a multi-faceted function in the regulation of gene expression during muscle wasting, although much work needs to be done to challenge these hypotheses. Participation of the ubiquitous calpains in atrophy-associated apoptosis and in the impairment of the regenerative capacity of the muscle is also possible, albeit less clear. Conversely, all evidence points to a positive role of calpains in myoblast fusion, setting aside a role of calpains at this stage in muscle atrophy.

4. The unique case of calpain 3

Calpain 3 studies result in a completely different picture compared to the ubiquitous calpains. Both in genetically modified conditions and in physiological or pathological models, it has never been possible to observe an increase in calpain 3 activity in relation to muscle wasting. In fact, it seems quite the opposite.

4.1. *Inherited calpain 3 deficiency: limb-girdle muscular dystrophy type 2A*

Inactivating mutations in the calpain 3 gene are responsible for limb-girdle muscular dystrophy (LGMD2A), a recessive muscle disorder characterized by a progressive weakness and atrophy of the proximal limb muscles, especially those of the posterior compartment (Fardeau, Eymard, Mignard, Tome, Richard, & Beckmann, 1996; Richard et al., 1995). The disease is also characterized by an elevation of serum creatine kinase level. At the histological level, the muscle shows inflammatory infiltrates and areas of degeneration/regeneration. These observations are indicative of some muscle fiber death. In calpain 3 knockout mice, a decrease in muscle weight has been observed in extensor digitorum longus (EDL) and soleus muscles associated with a decrease in fiber diameter, indicating hypotrophy (personal communication).

Several studies of ours have provided some clues about possible molecular mechanisms involved in the muscle wasting occurring in LGMD2A. We have shown that the level of apoptotic myonuclei was 100 times higher in LGMD2A patients than in other myopathies (Baghdiguian et al., 1999). Even if the number

of apoptotic nuclei is not very high from an absolute point of view, the accumulation of such events over time may be sufficient to participate in the atrophying phenotype in this slowly progressive disease. This increase in apoptosis is accompanied by a perturbation of the NF- κ B pathway. Immunohistochemistry showed that I κ B α accumulates in apoptotic and preapoptotic myonuclei whereas NF- κ B accumulates in the subsarcolemmal areas (Baghdiguian et al., 1999; Richard et al., 2000).

It is interesting to note that many studies reported the requirement for NF- κ B in the process of muscle wasting. Despite our observations seems to be in contradiction with these data, it has to be reminded that there is not a unique NF- κ B pathway. In fact, these factors exist as homo- or heterodimers of five different proteins (p65/RelA, relB, cREL, p50 and p52) (Perkins, 2000) and I κ B is also a family of proteins. Degradation of I κ B α required for NF- κ B activation can be induced by a large variety of stimuli and is not only dependent on the proteasome but also can be performed by ubiquitous calpains (Shumway, Maki, & Miyamoto, 1999). Several regulatory mechanisms such as selective activation of different NF- κ B dimers, modification of promoter accessibility or interaction with various co-activators, will then ensure the specificity and selectivity of the response (Perkins, 1997). This versatility of the NF- κ B pathway is perfectly illustrated in skeletal muscles. Indeed, cytokine induction in cachexia activates a “classical NF- κ B” pathway involving p65 and leading to protein loss, while in disuse or denervation atrophies, the activation of an alternative NF- κ B pathway involving p50 leads to expression of anti-apoptotic genes (Ghosh, May, & Kopp, 1998; Hunter, Stevenson, Koncarevic, Mitchell-Felton, Essig, & Kandarian, 2002). Whereas most studies indicate a positive influence of NF- κ B activation towards muscle atrophy (for recent examples, see Cai et al., 2004; Hunter & Kandarian, 2004), other ones indicate that NF- κ B may have a beneficial effect on cell survival since its inhibition in the presence of TNF- α increases myogenic cell death (Stewart et al., 2004) and that it may inhibit muscle proteolysis by negatively influences the expression of the proteasome (Luo, Sun, Hungness, & Hasselgren, 2001). Considering this pleiotropy where NF- κ B can be considered as a switchboard, the observation made in LGMD2A suggests that the impaired NF- κ B pathway is of an anti-apoptotic nature. Thus, LGMD2A would correspond to

a condition where NF- κ B is unable to exert an antiapoptotic response, leading to fiber loss and hypotrophy.

Global proteosynthesis and proteolysis were measured in muscles of calpain 3 deficient mice without showing any significant difference (Combaret, Bechet, Claustre, Taillandier, Richard, & Attaix, 2003). However, a reduction was observed in the expression of several components of proteolytic systems such as cathepsin L, the 14-kDa E2 conjugating enzyme, a protein involved in polyubiquitination, and the alpha-type C2 subunit of the 20S catalytic core of the proteasome. Of interest, the 14-kDa E2 was shown to increase in a model of cachexia with upregulation of IL-6 and TNF- α (Catalano et al., 2003; Taillandier et al., 1996; Temparis et al., 1994; Wing & Banville, 1994). Conjugation mediated by this E2 is supposed to be a rate-limiting step in the N-end rule mediated degradation of proteins (Wing & Bedard, 1996), suggesting that a perturbation of proteasome activity could occur in LGMD2A.

4.2. Downregulation of calpain 3 in atrophic conditions

Besides this genetic condition, several studies have demonstrated a decrease of calpain 3 at the messenger level. First, it was observed in two conditions related to an increase in cytokine level. In fact, calpain 3 has been shown to be downregulated in experimental cancer cachexia in Yoshida hepatoma-bearing rats and in the atrophying skeletal muscle of IL-6 transgenic mice (Busquets, Garcia-Martinez, Alvarez, Carbo, Lopez-Soriano, & Argiles, 2000; Tsujinaka et al., 1996). The decrease of the mRNA level of calpain 3 in IL-6 transgenic mice was restored after injection of anti IL-6 receptor antibody. Second, this reduction was also observed in two conditions related to loss of muscle activity: disuse and denervation (Jones, Hill, Krasney, O'Conner, Peirce, & Greenhaff, 2004; Stockholm et al., 2001; Tang et al., 2000). Considering the consequences of the deficiency in calpain 3 in LGMD2A, it can be suggested that the decreased expression of calpain 3 observed in these conditions may participate in muscle wasting by deregulation of an antiapoptotic response of the NF- κ B pathway or through modification of proteasome activity. The former is supported by the fact that disuse atrophy is associated with an increase in IKK- β transcripts (Jones et al., 2004).

4.3. Conclusion on calpain 3

The fact that deficiency in calpain 3 leads to a muscular dystrophy phenotype demonstrates that its activity is necessary for the homeostasis of skeletal muscle. Calpain 3 seems to exert a protective effect against atrophy by directing the NF- κ B pathway towards an antiapoptotic response and may also have some links with the proteasome activity. In addition, it looks like in atrophy the activity of calpain 3 has to be dismissed.

Besides these observations, it is worth mentioning the relationship between calpain 3 and the cytoskeleton since muscle atrophy is intimately linked to the regulation of this structure. Three elements are important to consider. First, the relationship of calpain 3 with titin, a major component of muscle (for review, see Granzier & Labeit, 2004). Second, the fact that calpain 3 has the ability to cleave a number of cytoskeletal proteins such as ezrin, talin, filamin, vinexin, fodrin, and titin (Guyon et al., 2003; Taveau et al., 2003). Third, the possible perturbation of cytoskeletal architecture in calpain 3-deficient muscle. Kramerova, Kudryashova, Tidball, & Spencer (2004) observed a misalignment of A-band in the sarcomere of calpain 3-deficient mice. Furthermore, a frequent feature seen in LGMD2A biopsies is lobulated fibers, a non-specific morphological alteration of the cytoarchitecture with abnormal subsarcolemmal distribution of mitochondria and focal Z streaming (Chae et al., 2001). To date, it is difficult to place all these observations in a precise figure or model to explain the function of calpain 3 and further studies are still necessary.

5. General conclusion

To date, the ubiquitous calpains have been clearly implicated in the initial degradative events of the cytoskeleton observed in atrophic conditions and in the necrotic process secondary to a deregulation of Ca²⁺ homeostasis in muscular dystrophies. Calpain 3 shares with the ubiquitous calpains the fact that they have cytoskeletal proteins as substrates. However, they present a major difference with respect to the regulation of their expression. Ubiquitous calpain activities are turned on in all the models of atrophy studied to date, whereas calpain 3 activity is turned down. One can address several questions: whether their effects on their common

substrates are different or identical, do they respond to antagonist signals and how could they influence each other? It has been shown that calpain 3 is able to cleave calpains and calpastatin suggesting a possible cross-talk between ubiquitous calpains and calpain 3 (Ono et al., 2004).

Another notable condition showing a different behavior of ubiquitous calpains and calpain 3 is mechanical stress. Using eccentric exercise as a model, our studies showed that calpain 3 is downregulated immediately after exercise whereas calpain 2 mRNA levels increase dramatically as early as one day after the exercise (Feasson et al., 2002). An increase in calpain 2 expression was also observed in muscle overloading (Spencer, Lu, & Tidball, 1997). In addition, an increasing ubiquitous calpain activity associated with a redistribution of the protein from the cytosol to the membrane was observed after level running (Arthur, Booker, & Belcastro, 1999; Belcastro, Gilchrist, & Scrubb, 1993).

Interestingly, some parallels can be drawn between the observations made at cellular and molecular levels during atrophy and those during mechanical stress such as exercise. Intense exercise induces cytoskeletal disruption with, in particular, Z-disc streaming (Sorichter, Puschendorf, & Mair, 1999). Apoptosis was observed in fibers and satellite cells after exercise (Arslan, Erdem, Sivri, Hascelik, & Tan, 2002; Podhorska-Okolow, Sandri, Zampieri, Brun, Rossini, & Carraro, 1998). Finally, the NF- κ B signaling pathway can be modulated during stretching (Kumar & Boriek, 2003; Lammerding, Kamm, & Lee, 2004; McClung, Lee, Thompson, Lowe, & Carson, 2003).

However, the final outcome is completely different as exercise directs skeletal muscle towards hypertrophy. This suggests that, depending on the cellular context, different processes will take place following an initial phase of sarcomere disruption. Therefore, we would like to propose that calpain-induced myofibril cleavage is a preliminary step allowing either the further degradation of myofibrillar proteins for amino acid supply or the qualitative remodeling of the muscle (Fig. 4). This preliminary step may be crucial in muscle, in which the protein content is mainly immobilized into the sarcomere, a quasi-crystalline structure. This sarcomere disruption will allow neosarcomerogenesis or substitution of one protein isoform with a different one more appropriate for a new physiologic condition.




Fig. 4. General model. After cleavage of sarcomeric proteins by calpains, the various fragments produced could be directed to one of two possible pathways according to the signal received: (1) In the case of an atrophying signal, they are sent to the proteasome for further degradation, leading to muscle atrophy. (2) In the absence of such a signal, they could be reincorporated/replaced in the sarcomere in order to adapt to new conditions.

As a final conclusion, despite the numerous data that have been generated on calpains and atrophy, much work is still needed to complete the picture of calpain functions, especially in the case of the puzzling calpain 3.

Acknowledgment

This work was funded by the Association Française contre les Myopathies. We would like to thank Dr. Nathalie Daniele and Dr. Susan Cure for critical reading of the manuscript, Carinne Roudaut for different aspects of the *capn3*^{−/−} animal model studies and Dr. Béatrice Benayoun for studies on the NF- κ B pathway. We also wish to acknowledge Prof. Stephen Baghdigian and Prof. Gerard Lefranc for their collaborations and Dr. Olivier Danos for his constant support.

References

- Acharyya, S., Ladner, K. J., Nelsen, L. L., Damrauer, J., Reiser, P. J., Swoap, S., et al. (2004). Cancer cachexia is regulated by selective targeting of skeletal muscle gene products. *Journal of Clinical Investigation*, 114, 370–378.

- Alderton, J. M., & Steinhardt, R. A. (2000a). How calcium influx through calcium leak channels is responsible for the elevated levels of calcium-dependent proteolysis in dystrophic myotubes. *Trends Cardiovascular Medicine*, 10, 268–272.
- Alderton, J. M., & Steinhardt, R. A. (2000b). Calcium influx through calcium leak channels is responsible for the elevated levels of calcium-dependent proteolysis in dystrophic myotubes. *Journal of Biological Chemistry*, 275, 9452–9460.
- Allen, D. L., Linderman, J. K., Roy, R. R., Grindeland, R. E., Mukku, V., & Edgerton, V. R. (1997). Growth hormone/IGF-I and/or resistive exercise maintains myonuclear number in hindlimb unweighted muscles. *Journal of Applied Physiology*, 83, 1857–1861.
- Anderson, K., Potter, A., Baban, D., & Davies, K. E. (2003). Protein expression changes in spinal muscular atrophy revealed with a novel antibody array technology. *Brain*, 126, 2052–2064.
- Arslan, S., Erdem, S., Sivri, A., Hascelik, Z., & Tan, E. (2002). Exercise-induced apoptosis of rat skeletal muscle and the effect of meloxicam. *Rheumatology International*, 21, 133–136.
- Arthur, G. D., Booker, T. S., & Belcastro, A. N. (1999). Exercise promotes a subcellular redistribution of calcium-stimulated protease activity in striated muscle. *Canadian Journal of Physiology and Pharmacology*, 77, 42–47.
- Babu, Y. S., Bugg, C. E., & Cook, W. J. (1988). Structure of calmodulin refined at 2.2 Å resolution. *Journal of Molecular Biology*, 204, 191–204.
- Badalamente, M. A., & Stracher, A. (2000). Delay of muscle degeneration and necrosis in mdx mice by calpain inhibition. *Muscle & Nerve*, 23, 106–111.
- Baghdiguian, S., Martin, M., Richard, I., Pons, F., Astier, C., Bourg, N., et al. (1999). Calpain 3 deficiency is associated with myonuclear apoptosis and profound perturbation of the IkappaB alpha/NF-kappaB pathway in limb-girdle muscular dystrophy type 2A. *Nature Medicine*, 5, 503–511.
- Baker, J. H., & Margolis, R. N. (1987). Calcium-activated protease activity in tenotomized muscle. *Muscle & Nerve*, 10, 34–40.
- Balcerzak, D., Cottin, P., Poussard, S., Cucuron, A., Brustis, J. J., & Ducastaing, A. (1998). Calpastatin-modulation of m-calpain activity is required for myoblast fusion. *European Journal of Cell Biology*, 75, 247–253.
- Balcerzak, D., Poussard, S., Brustis, J. J., Elamrani, N., Soriano, M., Cottin, P., et al. (1995). An antisense oligodeoxynucleotide to m-calpain mRNA inhibits myoblast fusion. *Journal of Cell Science*, 108(Pt 5), 2077–2082.
- Baldwin, K. M., & Haddad, F. (2002). Skeletal muscle plasticity: cellular and molecular responses to altered physical activity paradigms. *American Journal of Physical Medicine Rehabilitation*, 81, 40–51.
- Barnoy, S., Glaser, T., & Kosower, N. S. (1997). Calpain and calpastatin in myoblast differentiation and fusion: effects of inhibitors. *Biochimica Biophysica Acta*, 1358, 181–188.
- Barnoy, S., Zipser, Y., Glaser, T., Grimberg, Y., & Kosower, N. S. (1999). Association of calpain (Ca(2+)-dependent thiol protease) with its endogenous inhibitor calpastatin in myoblasts. *Journal of Cellular Biochemistry*, 74, 522–531.
- Belcastro, A. N., Gilchrist, J. S., & Scrubb, J. (1993). Function of skeletal muscle sarcoplasmic reticulum vesicles with exercise. *Journal of Applied Physiology*, 75, 2412–2418.
- Belozero, I., Shenkman, B., Mazin, M., & Leblanc, A. (2001). Effects of long-duration bed rest on structural compartments of m. soleus in man. *Journal of Gravitational Physiology*, 8, P71–P72.
- Berchtold, M. W., Brinkmeier, H., & Muntener, M. (2000). Calcium ion in skeletal muscle: its crucial role for muscle function, plasticity, and disease. *Physiological Reviews*, 80, 1215–1265.
- Bertorini, T. E., Bhattacharya, S. K., Palmieri, G. M., Chesney, C. M., Pifer, D., & Baker, B. (1982). Muscle calcium and magnesium content in Duchenne muscular dystrophy. *Neurology*, 32, 1088–1092.
- Bischoff, R. (1986). Proliferation of muscle satellite cells on intact myofibers in culture. *Developmental Biology*, 115, 129–139.
- Bizat, N., Hermel, J. M., Humbert, S., Jacquard, C., Creminon, C., Escartin, C., et al. (2003). In vivo calpain/caspase cross-talk during 3-nitropropionic acid-induced striatal degeneration: implication of a calpain-mediated cleavage of active caspase-3. *Journal of Biological Chemistry*, 278, 43245–43253.
- Bodine, S. C., Latres, E., Baumhueter, S., Lai, V. K., Nunez, L., Clarke, B. A., et al. (2001). Identification of ubiquitin ligases required for skeletal muscle atrophy. *Science*, 294, 1704–1708.
- Branca, D., Gugliucci, A., Bano, D., Brini, M., & Carafoli, E. (1999). Expression, partial purification and functional properties of the muscle-specific calpain isoform p94. *European Journal of Biochemistry*, 265, 839–846.
- Bullard, B., Sainsbury, G., & Miller, N. (1990). Digestion of proteins associated with the Z-disc by calpain. *Journal of Muscle Research and Cell Motility*, 11, 271–279.
- Busquets, S., Garcia-Martinez, C., Alvarez, B., Carbo, N., Lopez-Soriano, F. J., & Argiles, J. M. (2000). Calpain-3 gene expression is decreased during experimental cancer cachexia. *Biochimica Biophysica Acta*, 1475, 5–9.
- Cai, D., Frantz, J. D., Tawa, N. E., Jr., Melendez, P. A., Oh, B. C., Lidov, H. G., et al. (2004). IKKbeta/NF-kappaB activation causes severe muscle wasting in mice. *Cell*, 119, 285–298.
- Carragher, N. O., Levkau, B., Ross, R., & Raines, E. W. (1999). Degraded collagen fragments promote rapid disassembly of smooth muscle focal adhesions that correlates with cleavage of pp125(FAK), paxillin, and talin. *Journal of Cellular Biology*, 147, 619–630.
- Catalano, M. G., Fortunati, N., Arena, K., Costelli, P., Aragno, M., Danni, O., et al. (2003). Selective up-regulation of tumor necrosis factor receptor I in tumor-bearing rats with cancer-related cachexia. *International Journal of Oncology*, 23, 429–436.
- Chae, J., Minami, N., Jin, Y., Nakagawa, M., Murayama, K., Igarashi, F., et al. (2001). Calpain 3 gene mutations: genetic and clinicopathologic findings in limb-girdle muscular dystrophy. *Neuromuscular Disorders*, 11, 547–555.
- Childs, S. G. (2003). Muscle wasting. *Orthopedics and Nursing*, 22, 251–257.
- Chockalingam, P. S., Cholera, R., Oak, S. A., Zheng, Y., Jarrett, H. W., & Thomason, D. B. (2002). Dystrophin-glycoprotein complex and Ras and Rho GTPase signaling are altered in muscle atrophy. *American Journal of Physiology and Cell Physiology*, 283, C500–C511.

- Choi, Y. H., Lee, S. J., Nguyen, P., Jang, J. S., Lee, J., Wu, M. L., et al. (1997). Regulation of cyclin D1 by calpain protease. *Journal of Biological Chemistry*, 272, 28479–28484.
- Cohn, R. D., & Campbell, K. P. (2000). Molecular basis of muscular dystrophies. *Muscle & Nerve*, 23, 1456–1471.
- Combaret, L., Bechet, D., Claustre, A., Taillandier, D., Richard, I., & Attaix, D. (2003). Down-regulation of genes in the lysosomal and ubiquitin-proteasome proteolytic pathways in calpain-3-deficient muscle. *International Journal of Biochemistry and Cell Biology*, 35, 676–684.
- Combaret, L., Taillandier, D., Voisin, L., Samuels, S. E., Boespflug-Tanguy, O., & Attaix, D. (1996). No alteration in gene expression of components of the ubiquitin-proteasome proteolytic pathway in dystrophin-deficient muscles. *FEBS Letters*, 393, 292–296.
- Costelli, P., & Baccino, F. M. (2003). Mechanisms of skeletal muscle depletion in wasting syndromes: role of ATP-ubiquitin-dependent proteolysis. *Current Opinion in Clinical Nutrition Metabolism Care*, 6, 407–412.
- Costelli, P., Tullio, R. D., Baccino, F. M., & Melloni, E. (2001). Activation of Ca(2+)-dependent proteolysis in skeletal muscle and heart in cancer cachexia. *British Journal of Cancer*, 84, 946–950.
- Cyglér, M., Grochulski, P., & Blanchard, H. (2002). Crystallization and structural details of Ca(2+)-induced conformational changes in the EF-hand domain VI of calpain. *Methods in Molecular Biology*, 172, 243–260.
- De Tullio, R., Avena, M., Salamino, F., Pontremoli, S., & Melloni, E. (2000). Differential degradation of calpastatin by mu- and m-calpain in Ca(2+)-enriched human neuroblastoma LAN-5 cells. *FEBS Letters*, 475, 17–21.
- Dedieu, S., Mazerès, G., Poussard, S., Brustis, J. J., & Cottin, P. (2003). Myoblast migration is prevented by a calpain-dependent accumulation of MARCKS. *Biology of the Cell*, 95, 615–623.
- Dedieu, S., Poussard, S., Mazerès, G., Grise, F., Dargelos, E., Cottin, P., et al. (2004). Myoblast migration is regulated by calpain through its involvement in cell attachment and cytoskeletal organization. *Experimental Cell Research*, 292, 187–200.
- Diaz, B. G., Moldoveanu, T., Kuiper, M. J., Campbell, R. L., & Davies, P. L. (2004). Insertion sequence 1 of muscle-specific calpain, p94, acts as an internal propeptide. *Journal of Biological Chemistry*, 27656–27666.
- Donkor, I. O. (2000). A survey of calpain inhibitors. *Current Medicinal Chemistry*, 7, 1171–1188.
- Dupont-Versteegden, E. E., Murphy, R. J., Houle, J. D., Gurley, C. M., & Peterson, C. A. (1999). Activated satellite cells fail to restore myonuclear number in spinal cord transected and exercised rats. *American Journal of Physiology*, 277, C589–C597.
- Elce, J. S., Hegadorn, C., & Arthur, J. S. (1997). Autolysis, Ca2+ requirement, and heterodimer stability in m-calpain. *Journal of Biological Chemistry*, 272, 11268–11275.
- Elia, M., Carter, A., Bacon, S., Winearls, C. G., & Smith, R. (1981). Clinical usefulness of urinary 3-methylhistidine excretion in indicating muscle protein breakdown. *British Medical Journal (Clinical Research Education)*, 282, 351–354.
- Fardeau, M., Eymard, B., Mignard, C., Tome, F. M., Richard, I., & Beckmann, J. S. (1996). Chromosome 15-linked limb-girdle muscular dystrophy: clinical phenotypes in Reunion Island and French metropolitan communities. *Neuromuscular Disorders*, 6, 447–453.
- Feasson, L., Stockholm, D., Freyssen, D., Richard, I., Duguez, S., Beckmann, J. S., et al. (2002). Molecular adaptations of neuromuscular disease-associated proteins in response to eccentric exercise in human skeletal muscle. *Journal of Physiology*, 543, 297–306.
- Franco, S., Perrin, B., & Huttenlocher, A. (2004). Isoform specific function of calpain 2 in regulating membrane protrusion. *Experimental Cell Research*, 299, 179–187.
- Fujita, J., Tsujinaka, T., Yano, M., Ebisui, C., Saito, H., Katsume, A., et al. (1996). Anti-interleukin-6 receptor antibody prevents muscle atrophy in colon-26 adenocarcinoma-bearing mice with modulation of lysosomal and ATP-ubiquitin-dependent proteolytic pathways. *International Journal of Cancer*, 68, 637–643.
- Furuno, K., Goodman, M. N., & Goldberg, A. L. (1990). Role of different proteolytic systems in the degradation of muscle proteins during denervation atrophy. *Journal of Biological Chemistry*, 265, 8550–8557.
- Ghosh, S., May, M. J., & Kopp, E. B. (1998). NF-kappa B and Rel proteins: evolutionarily conserved mediators of immune responses. *Annual Review of Immunology*, 16, 225–260.
- Gillis, J. M. (1999). Understanding dystrophinopathies: an inventory of the structural and functional consequences of the absence of dystrophin in muscles of the mdx mouse. *Journal of Muscle Research and Cell Motility*, 20, 605–625.
- Glading, A., Lauffenburger, D. A., & Wells, A. (2002). Cutting to the chase: calpain proteases in cell motility. *Trends in Cell Biology*, 12, 46–54.
- Glass, D. J. (2003). Molecular mechanisms modulating muscle mass. *Trends in Molecular Medicine*, 9, 344–350.
- Goll, D. E., Thompson, V. F., Li, H., Wei, W., & Cong, J. (2003). The calpain system. *Physiological Reviews*, 83, 731–801.
- Granzier, H. L., & Labeit, S. (2004). The giant protein titin: a major player in myocardial mechanics, signaling, and disease. *Circulation Research*, 94, 284–295.
- Guyon, J. R., Kudryashova, E., Potts, A., Dalkilic, I., Brosius, M. A., Thompson, T. G., et al. (2003). Calpain 3 cleaves filamin C and regulates its ability to interact with gamma- and delta-sarcoglycans. *Muscle & Nerve*, 28, 472–483.
- Haddad, F., Roy, R. R., Zhong, H., Edgerton, V. R., & Baldwin, K. M. (2003). Atrophy responses to muscle inactivity. II. Molecular markers of protein deficits. *Journal of Applied Physiology*, 95, 791–802.
- Hall, Z. W., & Ralston, E. (1989). Nuclear domains in muscle cells. *Cell*, 59, 771–772.
- Hasselgren, P. O. (1999). Role of the ubiquitin-proteasome pathway in sepsis-induced muscle catabolism. *Molecular Biology Reports*, 26, 71–76.
- Hasselgren, P. O., & Fischer, J. E. (2001). Muscle cachexia: current concepts of intracellular mechanisms and molecular regulation. *Annals of Surgery*, 233, 9–17.
- Heck, C. S., & Davis, H. L. (1988). Effect of denervation and nerve extract on ultrastructure of muscle. *Experimental Neurology*, 100, 139–153.
- Hikida, R. S., Van Nostran, S., Murray, J. D., Staron, R. S., Gordon, S. E., & Kraemer, W. J. (1997). Myonuclear loss in

- atrophied soleus muscle fibers. *Anatomical Records*, 247, 350–354.
- Hirai, S., Kawasaki, H., Yaniv, M., & Suzuki, K. (1991). Degradation of transcription factors, c-Jun and c-Fos, by calpain. *FEBS Letters*, 287, 57–61.
- Honda, S., Marumoto, T., Hirota, T., Nitta, M., Arima, Y., Ogawa, M., et al. (2004). Activation of *m*-calpain is required for chromosome alignment on the metaphase plate during mitosis. *Journal of Biological Chemistry*, 279, 10615–10623.
- Hong, D. H., & Forsberg, N. E. (1995). Effects of dexamethasone on protein degradation and protease gene expression in rat L8 myotube cultures. *Molecular and Cell Endocrinology*, 108, 199–209.
- Hopf, F. W., Turner, P. R., Denetclaw, W. F., Jr., Reddy, P., & Steinhardt, R. A. (1996). A critical evaluation of resting intracellular free calcium regulation in dystrophic mdx muscle. *American Journal of Physiology*, 271, C1325–C1339.
- Hornberger, T. A., Hunter, R. B., Kandarian, S. C., & Esser, K. A. (2001). Regulation of translation factors during hindlimb unloading and denervation of skeletal muscle in rats. *American Journal of Physiology and Cell Physiology*, 281, C179–C187.
- Hosfield, C. M., Elce, J. S., Davies, P. L., & Jia, Z. (1999). Crystal structure of calpain reveals the structural basis for Ca(2+)-dependent protease activity and a novel mode of enzyme activation. *EMBO Journal*, 18, 6880–6889.
- Huang, J., & Forsberg, N. E. (1998). Role of calpain in skeletal-muscle protein degradation. *Proceedings of the National Academy of Sciences of United States of America*, 95, 12100–12105.
- Huff-Lonergan, E., Mitsuhashi, T., Beekman, D. D., Parrish, F. C., Jr., Olson, D. G., & Robson, R. M. (1996). Proteolysis of specific muscle structural proteins by mu-calpain at low pH and temperature is similar to degradation in postmortem bovine muscle. *Journal of Animal Sciences*, 74, 993–1008.
- Hunter, R. B., & Kandarian, S. C. (2004). Disruption of either the Nfkb1 or the Bcl3 gene inhibits skeletal muscle atrophy. *Journal of Clinical Investigation*, 114, 1504–1511.
- Hunter, R. B., Stevenson, E., Koncarevic, A., Mitchell-Felton, H., Essig, D. A., & Kandarian, S. C. (2002). Activation of an alternative NF-kappaB pathway in skeletal muscle during disuse atrophy. *FASEB Journal*, 16, 529–538.
- Irintchev, A., Zwyer, M., & Wernig, A. (1997). Impaired functional and structural recovery after muscle injury in dystrophic mdx mice. *Neuromuscular Disorders*, 7, 117–125.
- Iwata, Y., Katanosaka, Y., Arai, Y., Komamura, K., Miyatake, K., & Shigekawa, M. (2003). A novel mechanism of myocyte degeneration involving the Ca2+-permeable growth factor-regulated channel. *Journal of Cell Biology*, 161, 957–967.
- Jackman, R. W., & Kandarian, S. C. (2004). The molecular basis of skeletal muscle atrophy. *American Journal of Physiology and Cell Physiology*, 287, C834–C843.
- Jackson, M. J., Jones, D. A., & Edwards, R. H. (1985). Measurements of calcium and other elements in muscle biopsy samples from patients with Duchenne muscular dystrophy. *Clinical Chimica Acta*, 147, 215–221.
- Jagoe, R. T., Lecker, S. H., Gomes, M., & Goldberg, A. L. (2002). Patterns of gene expression in atrophying skeletal muscles: response to food deprivation. *FASEB Journal*, 16, 1697–1712.
- Jejurikar, S. S., Marcelo, C. L., & Kuzon, W. M., Jr. (2002). Skeletal muscle denervation increases satellite cell susceptibility to apoptosis. *Plastics and Reconstructing Surgery*, 110, 160–168.
- Jejurikar, S. S., & Kuzon, W. M., Jr. (2003). Satellite cell depletion in degenerative skeletal muscle. *Apoptosis*, 8, 573–578.
- Johnson, P., & Perry, S. V. (1970). Biological activity and the 3-methylhistidine content of actin and myosin. *Biochemistry Journal*, 119, 293–298.
- Jones, S. W., Hill, R. J., Krasney, P. A., O'Conner, B., Peirce, N., & Greenhaff, P. L. (2004). Disuse atrophy and exercise rehabilitation in humans profoundly affects the expression of genes associated with the regulation of skeletal muscle mass. *FASEB Journal*, 18, 1025–1027.
- Kandarian, S. C., & Stevenson, E. J. (2002). Molecular events in skeletal muscle during disuse atrophy. *Exercise and Sport Sciences Reviews*, 30, 111–116.
- Karpati, G., Hilton-Jones, D., & Griggs, R. C. (2001). *Disorders of voluntary muscle: vol. xiv*. Cambridge, New York: Cambridge University Press, p. 775.
- Kent, M. P., Spencer, M. J., & Koohmaraie, M. (2004). Postmortem proteolysis is reduced in transgenic mice overexpressing calpastatin. *Journal of Animal Sciences*, 82, 794–801.
- Kim, M. J., Jo, D. G., Hong, G. S., Kim, B. J., Lai, M., Cho, D. H., et al. (2002). Calpain-dependent cleavage of cain/cabin1 activates calcineurin to mediate calcium-triggered cell death. *Proceedings of the National Academy of Sciences of United States of America*, 99, 9870–9875.
- Koohmaraie, M. (1992a). Ovine skeletal muscle multicatalytic proteinase complex (proteasome): purification, characterization, and comparison of its effects on myofibrils with mu-calpains. *Journal of Animal Sciences*, 70, 3697–3708.
- Koohmaraie, M. (1992b). The role of Ca(2+)-dependent proteases (calpains) in post mortem proteolysis and meat tenderness. *Biochimie*, 74, 239–245.
- Kramerova, I., Kudryashova, E., Tidball, J. G., & Spencer, M. J. (2004). Null mutation of calpain 3 (p94) in mice causes abnormal sarcomere formation in vivo and in vitro. *Human Molecular Genetics*, 13, 1373–1388.
- Kumamoto, T., Ueyama, H., Watanabe, S., Yoshioka, K., Miike, T., Goll, D. E., et al. (1995). Immunohistochemical study of calpain and its endogenous inhibitor in the skeletal muscle of muscular dystrophy. *Acta Neuropathology (Berlin)*, 89, 399–403.
- Kumar, A., & Boriek, A. M. (2003). Mechanical stress activates the nuclear factor-kappaB pathway in skeletal muscle fibers: a possible role in Duchenne muscular dystrophy. *FASEB Journal*, 17, 386–396.
- Kwak, K. B., Chung, S. S., Kim, O. M., Kang, M. S., Ha, D. B., & Chung, C. H. (1993). Increase in the level of m-calpain correlates with the elevated cleavage of filamin during myogenic differentiation of embryonic muscle cells. *Biochimica Biophysica Acta*, 1175, 243–249.
- Lammerding, J., Kamm, R. D., & Lee, R. T. (2004). Mechanotransduction in cardiac myocytes. *Annals of New York Academy of Sciences*, 1015, 53–70.

- Lapidos, K. A., Kakkar, R., & McNally, E. M. (2004). The dystrophin glycoprotein complex: signaling strength and integrity for the sarcolemma. *Circular Research*, 94, 1023–1031.
- Lecker, S. H., Jagoe, R. T., Gilbert, A., Gomes, M., Baracos, V., Bailey, J., et al. (2004). Multiple types of skeletal muscle atrophy involve a common program of changes in gene expression. *FASEB Journal*, 18, 39–51.
- Leeuwenburgh, C. (2003). Role of apoptosis in sarcopenia. *Journal of Gerontology A: Biological Sciences and Medicine Sciences*, 58, 999–1001.
- Li, Y., Gazdoui, S., Pan, Z. Q., & Fuchs, S. Y. (2004). Stability of homologue of Slimb F-box protein is regulated by availability of its substrate. *Journal of Biological Chemistry*, 279, 11074–11080.
- Lin, G. D., Chattopadhyay, D., Maki, M., Wang, K. K., Carson, M., Jin, L., et al. (1997). Crystal structure of calcium bound domain VI of calpain at 1.9 Å resolution and its role in enzyme assembly, regulation, and inhibitor binding. *Natural and Structural Biology*, 4, 539–547.
- Luo, G., Sun, X., Hungness, E., & Hasselgren, P. O. (2001). Heat shock protects L6 myotubes from catabolic effects of dexamethasone and prevents downregulation of NF-kappaB. *American Journal of Physiology: Regulatory Integrative and Comparative Physiology*, 281, R1193–R1200.
- Luz, M. A., Marques, M. J., & Santo Neto, H. (2002). Impaired regeneration of dystrophin-deficient muscle fibers is caused by exhaustion of myogenic cells. *Brazilian Journal of Medical and Biological Research*, 35, 691–695.
- MacLennan, P. A., McArdle, A., & Edwards, R. H. (1991). Effects of calcium on protein turnover of incubated muscles from mdx mice. *American Journal of Physiology*, 260, E594–E598.
- Massa, R., Carpenter, S., Holland, P., & Karpati, G. (1992). Loss and renewal of thick myofilaments in glucocorticoid-treated rat soleus after denervation and reinnervation. *Muscle & Nerve*, 15, 1290–1298.
- McCarter, G. C., & Steinhardt, R. A. (2000). Increased activity of calcium leak channels caused by proteolysis near sarcolemmal ruptures. *Journal of Membrane Biology*, 176, 169–174.
- McClung, J. M., Lee, W. J., Thompson, R. W., Lowe, L. L., & Carson, J. A. (2003). RhoA induction by functional overload and nandrolone decanoate administration in rat skeletal muscle. *Pflugers Archives*, 447, 345–355.
- McCroskery, S., Thomas, M., Maxwell, L., Sharma, M., & Kambadur, R. (2003). Myostatin negatively regulates satellite cell activation and self-renewal. *Journal of Cell Biology*, 162, 1135–1147.
- Melone, M. A., Peluso, G., Petillo, O., Galderisi, U., & Cotrufo, R. (1999). Defective growth in vitro of Duchenne Muscular Dystrophy myoblasts: the molecular and biochemical basis. *Journal of Cellular Biochemistry*, 76, 118–132.
- Minami, Y., Emori, Y., Kawasaki, H., & Suzuki, K. (1987). E-F hand structure-domain of calcium-activated neutral protease (CANP) can bind Ca²⁺ ions. *Journal of Biochemistry (Tokyo)*, 101, 889–895.
- Mitchell, P. O., & Pavlath, G. K. (2004). Skeletal muscle atrophy leads to loss and dysfunction of muscle precursor cells. *American Journal of Physiology and Cell Physiology*, C1753–C1762.
- Moldoveanu, T., Campbell, R. L., Cuerrier, D., & Davies, P. L. (2004). Crystal structures of calpain-E64 and -leupeptin inhibitor complexes reveal mobile loops gating the active site. *Journal of Molecular Biology*, 343, 1313–1326.
- Moldoveanu, T., Hosfield, C. M., Lim, D., Elce, J. S., Jia, Z., & Davies, P. L. (2002). A Ca(2+) switch aligns the active site of calpain. *Cell*, 108, 649–660.
- Mongini, T., Ghigo, D., Doriguzzi, C., Bussolino, F., Pescarmona, G., Pollo, B., et al. (1988). Free cytoplasmic Ca⁺⁺ at rest and after cholinergic stimulus is increased in cultured muscle cells from Duchenne muscular dystrophy patients. *Neurology*, 38, 476–480.
- Nakamura, M., Imajoh-Ohmi, S., Suzuki, K., & Kawashima, S. (1991). An endogenous inhibitor of calcium-activated neutral protease in UMX 7.1 hamster dystrophy. *Muscle & Nerve*, 14, 701–708.
- Neerunjun, J. S., & Dubowitz, V. (1979). Increased calcium-activated neutral protease activity in muscles of dystrophic hamsters and mice. *Journal of the Neurological Sciences*, 40, 105–111.
- Oda, A., Wakao, H., & Fujita, H. (2002). Calpain is a signal transducer and activator of transcription (STAT) 3 and STAT5 protease. *Blood*, 99, 1850–1852.
- Ohtsuka, H., Yajima, H., Maruyama, K., & Kimura, S. (1997). The N-terminal Z repeat 5 of connectin/titin binds to the C-terminal region of alpha-actinin. *Biochemical and Biophysical Research Communications*, 235, 1–3.
- Olive, M., & Ferrer, I. (2000). Bcl-2 and bax immunohistochemistry in denervation-reinnervation and necrosis-regeneration of rat skeletal muscles. *Muscle & Nerve*, 23, 1862–1867.
- Ono, Y., Kakinuma, K., Torii, F., Irie, A., Nakagawa, K., Labeit, S., et al. (2004). Possible regulation of the conventional calpain system by skeletal muscle-specific calpain, p94/calpain 3. *Journal of Biological Chemistry*, 279, 2761–2771.
- Ontell, M. P., Hughes, D., Hauschka, S. D., & Ontell, M. (1992). Transient neonatal denervation alters the proliferative capacity of myosatellite cells in dystrophic (129ReJdy/dy) muscle. *Journal of Neurobiology*, 23, 407–419.
- Perkins, N. D. (1997). Achieving transcriptional specificity with NF-kappa B. *International Journal of Biochemistry and Cell Biology*, 29, 1433–1448.
- Perkins, N. D. (2000). The Rel/NF-kappa B family: friend and foe. *Trends in Biochemical Sciences*, 25, 434–440.
- Piechaczyk, M. (2000). Proteolysis of p53 protein by ubiquitous calpains. *Methods in Molecular Biology*, 144, 297–307.
- Podhorska-Okolow, M., Sandri, M., Zampieri, S., Brun, B., Rossini, K., & Carraro, U. (1998). Apoptosis of myofibres and satellite cells: exercise-induced damage in skeletal muscle of the mouse. *Neuropathology and Applied Neurobiology*, 24, 518–531.
- Pontremoli, S., Melloni, E., Viotti, P. L., Michetti, M., Salamino, F., & Horecker, B. L. (1991). Identification of two calpastatin forms in rat skeletal muscle and their susceptibility to digestion by homologous calpains. *Archives of Biochemistry and Biophysics*, 288, 646–652.
- Purintrapiban, J., Wang, M. C., & Forsberg, N. E. (2003). Degradation of sarcomeric and cytoskeletal proteins in cultured skeletal muscle cells. *Comparative Biochemistry and Physiology B: Biochemistry and Molecular Biology*, 136, 393–401.

- Raynaud, F., Bonnal, C., Fernandez, E., Bremaud, L., Cerutti, M., Lebart, M. C., et al. (2003). The calpain I-alpha-actinin interaction: Resting complex between the calcium-dependent protease and its target in cytoskeleton. *European Journal of Biochemistry*, 270, 4662–4670.
- Raynaud, F., Carnac, G., Marcilhac, A., & Benyamin, Y. (2004). m-Calpain implication in cell cycle during muscle precursor cell activation. *Experimental Cell Research*, 298, 48–57.
- Reddy, P. A., Anandavalli, T. E., & Anandaraj, M. P. (1986). Calcium activated neutral proteases (milli- and micro-CANP) and endogenous CANP inhibitor of muscle in Duchenne muscular dystrophy (DMD). *Clinical Chimica Acta*, 160, 281–288.
- Renault, V., Thornell, L. E., Eriksson, P. O., Butler-Browne, G., Mouly, V., & Thorne, L. E. (2002). Regenerative potential of human skeletal muscle during aging. *Aging Cell*, 1, 132–139.
- Reverter, D., Strobl, S., Fernandez-Catalan, C., Sorimachi, H., Suzuki, K., & Bode, W. (2001). Structural basis for possible calcium-induced activation mechanisms of calpains. *Biological Chemistry*, 382, 753–766.
- Richard, I., Broux, O., Allamand, V., Fougereuse, F., Chianilulchai, N., Bourg, N., et al. (1995). Mutations in the proteolytic enzyme, calpain 3, cause limb-girdle muscular dystrophy type 2A. *Cell*, 81, 27–40.
- Richard, I., Roudaut, C., Marchand, S., Baghdiguian, S., Herasse, M., Stockholm, D., et al. (2000). Loss of calpain 3 proteolytic activity leads to muscular dystrophy and to apoptosis-associated IkappaBalpha/nuclear factor kappaB pathway perturbation in mice. *Journal of Cell Biology*, 151, 1583–1590.
- Robert, V., Massimino, M. L., Tosello, V., Marsault, R., Cantini, M., Sorrentino, V., et al. (2001). Alteration in calcium handling at the subcellular level in mdx myotubes. *Journal of Biological Chemistry*, 276, 4647–4651.
- Rodrigues Ade, C., Geuna, S., Rodrigues, S. P., Silva, M. D., & Aragon, F. F. (2002). Satellite cells and myonuclei in neonatally denervated rat muscle. *Italian Journal of Anatomic Embryology*, 107, 51–56.
- Rossi, D., & Gaidano, G. (2003). Messengers of cell death: apoptotic signaling in health and disease. *Haematologica*, 88, 212–218.
- Sartorelli, V., & Fulco, M. (2004). Molecular and cellular determinants of skeletal muscle atrophy and hypertrophy. *Sciences of STKE*, re11.
- Shumway, S. D., Maki, M., & Miyamoto, S. (1999). The PEST domain of IkappaBalpha is necessary and sufficient for in vitro degradation by mu-calpain. *Journal of Biological Chemistry*, 274, 30874–30881.
- Sneddon, A. A., Delday, M. I., & Maltin, C. A. (2000). Amelioration of denervation-induced atrophy by clenbuterol is associated with increased PKC-alpha activity. *American Journal of Physiology: Endocrinology and Metabolism*, 279, E188–E195.
- Solomon, V., Baracos, V., Sarraf, P., & Goldberg, A. L. (1998). Rates of ubiquitin conjugation increase when muscles atrophy, largely through activation of the N-end rule pathway. *Proceedings of the National Academy of Sciences of United States of America*, 95, 12602–12607.
- Solomon, V., & Goldberg, A. L. (1996). Importance of the ATP-ubiquitin-proteasome pathway in the degradation of soluble and myofibrillar proteins in rabbit muscle extracts. *Journal of Biological Chemistry*, 271, 26690–26697.
- Solomon, V., Lecker, S. H., & Goldberg, A. L. (1998). The N-end rule pathway catalyzes a major fraction of the protein degradation in skeletal muscle. *Journal of Biological Chemistry*, 273, 25216–25222.
- Sorichter, S., Puschendorf, B., & Mair, J. (1999). Skeletal muscle injury induced by eccentric muscle action: muscle proteins as markers of muscle fiber injury. *Exercise Immunology Review*, 5, 5–21.
- Sorimachi, H., Imajoh-Ohmi, S., Emori, Y., Kawasaki, H., Ohno, S., Minami, Y., et al. (1989). Molecular cloning of a novel mammalian calcium-dependent protease distinct from both m- and mu- type. Specific expression of the mRNA in skeletal muscle. *Journal of Biological Chemistry*, 264, 20106–20111.
- Sorimachi, H., Kimbara, K., Kimura, S., Takahashi, M., Ishiura, S., Sasagawa, N., et al. (1995). Muscle-specific calpain, p94, responsible for limb-girdle muscular dystrophy type 2A, associates with connectin through IS2, a p94-specific sequence. *Journal of Biological Chemistry*, 270, 31158–31162.
- Sorimachi, H., Toyama-Sorimachi, N., Saido, T. C., Kawasaki, H., Sugita, H., Miyasaka, M., et al. (1993). Muscle-specific calpain, p94, is degraded by autolysis immediately after translation, resulting in disappearance from muscle. *Journal of Biological Chemistry*, 268, 10593–10605.
- Spencer, M. J., Croall, D. E., & Tidball, J. G. (1995). Calpains are activated in necrotic fibers from mdx dystrophic mice. *Journal of Biological Chemistry*, 270, 10909–10914.
- Spencer, M. J., Lu, B., & Tidball, J. G. (1997). Calpain II expression is increased by changes in mechanical loading of muscle in vivo. *Journal of Cellular Biochemistry*, 64, 55–66.
- Spencer, M. J., & Mellgren, R. L. (2002). Overexpression of a calpastatin transgene in mdx muscle reduces dystrophic pathology. *Human Molecular Genetics*, 11, 2645–2655.
- Squier, M. K., Miller, A. C., Malkinson, A. M., & Cohen, J. J. (1994). Calpain activation in apoptosis. *Journal of Cell Physiology*, 159, 229–237.
- Stewart, C. E., Newcomb, P. V., & Holly, J. M. (2004). Multifaceted roles of TNF-alpha in myoblast destruction: a multitude of signal transduction pathways. *Journal of Cell Physiology*, 198, 237–247.
- Stockholm, D., Bartoli, M., Sillon, G., Bourg, N., Davoust, J., & Richard, I. (2005). Imaging calpain protease activity by multi-photon FRET in living mice. *Journal of Molecular Biology*, 346, 15–22.
- Stockholm, D., Herasse, M., Marchand, S., Praud, C., Roudaut, C., Richard, I., et al. (2001). Calpain 3 mRNA expression in vivo after denervation and during muscle regeneration. *American Journal of Physiology and Cellular Physiology*, 280, C1561–C1569.
- Strobl, S., Fernandez-Catalan, C., Braun, M., Huber, R., Masumoto, H., Nakagawa, K., et al. (2000). The crystal structure of calcium-free human m-calpain suggests an electrostatic switch mechanism for activation by calcium. *Proceedings of the National Academy of Sciences of United States of America*, 97, 588–592.
- Stubgen, J. P. (1994). Limb girdle muscular dystrophy: description of a phenotype. *Muscle & Nerve*, 17, 1449–1455.

- Taillandier, D., Aourousseau, E., Meynial-Denis, D., Bechet, D., Ferrara, M., Cottin, P., et al. (1996). Coordinate activation of lysosomal, Ca²⁺-activated and ATP-ubiquitin-dependent proteinases in the unweighted rat soleus muscle. *Biochemical Journal*, 316(Pt 1), 65–72.
- Tang, H., Cheung, W. M., Ip, F. C., & Ip, N. Y. (2000). Identification and characterization of differentially expressed genes in denervated muscle. *Molecular and Cellular Neuroscience*, 16, 127–140.
- Taveau, M., Bourg, N., Sillon, G., Roudaut, C., Bartoli, M., & Richard, I. (2003). Calpain 3 is activated through autolysis within the active site and lyses sarcomeric and sarcolemmal components. *Molecular and Cellular Biology*, 23, 9127–9135.
- Taylor, R. G., Geesink, G. H., Thompson, V. F., Koohmaraie, M., & Goll, D. E. (1995). Is Z-disk degradation responsible for postmortem tenderization? *Journal of Animal Sciences*, 73, 1351–1367.
- Temme-Grove, C. J., Wert, D., Thompson, V. F., Allen, R. E., & Goll, D. E. (1999). Microinjection of calpastatin inhibits fusion in myoblasts. *Experimental Cell Research*, 247, 293–303.
- Temparis, S., Asensi, M., Taillandier, D., Aourousseau, E., Larbaud, D., Obled, A., et al. (1994). Increased ATP-ubiquitin-dependent proteolysis in skeletal muscles of tumor-bearing rats. *Cancer Research*, 54, 5568–5573.
- Tews, D. S. (2002). Apoptosis and muscle fibre loss in neuromuscular disorders. *Neuromuscular Disorders*, 12, 613–622.
- Tidball, J. G., Albrecht, D. E., Lokensgard, B. E., & Spencer, M. J. (1995). Apoptosis precedes necrosis of dystrophin-deficient muscle. *Journal of Cell Science*, 108(Pt 6), 2197–2204.
- Tidball, J. G., & Spencer, M. J. (2000). Calpains and muscular dystrophies. *International Journal of Biochemistry and Cell Biology*, 32, 1–5.
- Tidball, J. G., & Spencer, M. J. (2002). Expression of a calpastatin transgene slows muscle wasting and obviates changes in myosin isoform expression during murine muscle disuse. *Journal of Physiology*, 545, 819–828.
- Tompa, P., Emori, Y., Sorimachi, H., Suzuki, K., & Friedrich, P. (2001). Domain III of calpain is a Ca²⁺-regulated phospholipid-binding domain. *Biochemical and Biophysical Research Communications*, 280, 1333–1339.
- Tsujinaka, T., Fujita, J., Ebisui, C., Yano, M., Kominami, E., Suzuki, K., et al. (1996). Interleukin 6 receptor antibody inhibits muscle atrophy and modulates proteolytic systems in interleukin 6 transgenic mice. *Journal of Clinical Investigation*, 97, 244–249.
- Turner, P. R., Fong, P. Y., Denetclaw, W. F., & Steinhardt, R. A. (1991). Increased calcium influx in dystrophic muscle. *Journal of Cell Biology*, 115, 1701–1712.
- Turner, P. R., Schultz, R., Ganguly, B., & Steinhardt, R. A. (1993). Proteolysis results in altered leak channel kinetics and elevated free calcium in mdx muscle. *Journal of Membrane Biology*, 133, 243–251.
- Turner, P. R., Westwood, T., Regen, C. M., & Steinhardt, R. A. (1988). Increased protein degradation results from elevated free calcium levels found in muscle from mdx mice. *Nature*, 335, 735–738.
- Tutdibi, O., Brinkmeier, H., Rudel, R., & Fohr, K. J. (1999). Increased calcium entry into dystrophin-deficient muscle fibres of MDX and ADR-MDX mice is reduced by ion channel blockers. *Journal of Physiology*, 515(Pt 3), 859–868.
- Vandebrouck, C., Martin, D., Colson-Van Schoor, M., Debaix, H., & Gailly, P. (2002). Involvement of TRPC in the abnormal calcium influx observed in dystrophic (mdx) mouse skeletal muscle fibers. *Journal of Cell Biology*, 158, 1089–1096.
- Vanderklish, P. W., Krushel, L. A., Holst, B. H., Gally, J. A., Crossin, K. L., & Edelman, G. M. (2000). Marking synaptic activity in dendritic spines with a calpain substrate exhibiting fluorescence resonance energy transfer. *Proceedings of the National Academy of Sciences of United States of America*, 97, 2253–2258.
- Voisin, L., Breuille, D., Combaret, L., Pouyet, C., Taillandier, D., Aourousseau, E., et al. (1996). Muscle wasting in a rat model of long-lasting sepsis results from the activation of lysosomal, Ca²⁺-activated, and ubiquitin-proteasome proteolytic pathways. *Journal of Clinical Investigation*, 97, 1610–1617.
- Walling, A. D. (1999). Amyotrophic lateral sclerosis: Lou Gehrig's disease. *American of Family Physician*, 59, 1489–1496.
- Walowitz, J. L., Bradley, M. E., Chen, S., & Lee, T. (1998). Proteolytic regulation of the zinc finger transcription factor YY1, a repressor of muscle-restricted gene expression. *Journal of Biological Chemistry*, 273, 6656–6661.
- Wang, K. K., Roufogalis, B. D., & Villalobo, A. (1989). Characterization of the fragmented forms of calcineurin produced by calpain I. *Biochemistry and Cell Biology*, 67, 703–711.
- Watt, F., & Molloy, P. L. (1993). Specific cleavage of transcription factors by the thiol protease, m-calpain. *Nucleic Acids Research*, 21, 5092–5100.
- Wei, L., Zhou, W., Wang, L., & Schwartz, R. J. (2000). beta(1)-integrin and PI 3-kinase regulate RhoA-dependent activation of skeletal alpha-actin promoter in myoblasts. *American Journal of Physiology: Heart and Circulatory Physiology*, 278, H1736–H1743.
- Weindruch, R. (1995). Interventions based on the possibility that oxidative stress contributes to sarcopenia. *Journal of Gerontology A: Biological Sciences and Medicinal Sciences*, 50 Spec No, 157–161.
- Whitehouse, A. S., Smith, H. J., Drake, J. L., & Tisdale, M. J. (2001). Mechanism of attenuation of skeletal muscle protein catabolism in cancer cachexia by eicosapentaenoic acid. *Cancer Research*, 61, 3604–3609.
- Williams, A. B., Decourten-Myers, G. M., Fischer, J. E., Luo, G., Sun, X., & Hasselgren, P. O. (1999). Sepsis stimulates release of myofilaments in skeletal muscle by a calcium-dependent mechanism. *FASEB Journal*, 13, 1435–1443.
- Wing, S. S., & Banville, D. (1994). 14-kDa ubiquitin-conjugating enzyme: structure of the rat gene and regulation upon fasting and by insulin. *American Journal of Physiology*, 267, E39–E48.
- Wing, S. S., & Bedard, N. (1996). Insulin-like growth factor I stimulates degradation of an mRNA transcript encoding the 14 kDa ubiquitin-conjugating enzyme. *Biochemical Journal*, 319(Pt 2), 455–461.
- Xiao, Y. Y., Wang, M. C., Purintrapiban, J., & Forsberg, N. E. (2003). Roles of mu-calpain in cultured L8 muscle cells: application of a skeletal muscle-specific gene expression system. *Comparative Biochemistry and Physiology C: Toxicology and Pharmacology*, 134, 439–450.

- Yamakuchi, M., Higuchi, I., Masuda, S., Ohira, Y., Kubo, T., Kato, Y., et al. (2000). Type I muscle atrophy caused by microgravity-induced decrease of myocyte enhancer factor 2C (MEF2C) protein expression. *FEBS Letters*, 477, 135–140.
- Yoshimura, N., Murachi, T., Heath, R., Kay, J., Jasani, B., & Newman, G. R. (1986). Immunogold electron-microscopic localisation of calpain I in skeletal muscle of rats. *Cell Tissue Research*, 244, 265–270.
- Zhang, W., Lu, Q., Xie, Z. J., & Mellgren, R. L. (1997). Inhibition of the growth of WI-38 fibroblasts by benzyloxycarbonyl-Leu-Leu-Tyr diazomethyl ketone: evidence that cleavage of p53 by a calpain-like protease is necessary for G1 to S-phase transition. *Oncogene*, 14, 255–263.