

MATH 140 Practice Set #1

Calculus 1 (McGill University)



Scan to open on Studocu

MATH 140 Practice Set #1

MATH 140

These are a few problems to illustrate some types of limits and derivatives. The level of difficulty of these problems is similar to the exercises in your book. More problems will be posted throughout the semester.

1. Evaluate each of the following limits:

(a)
$$\lim_{x \to -2} \frac{\frac{x-2}{x+3} + 4}{x+2}$$

(b)
$$\lim_{x \to \infty} \left(\sqrt{x^2 - 3x} - x \right)$$

(c)
$$\lim_{x \to 0} \frac{\sin^2(2x)\tan(3x)}{x^3}$$

2. Derive each of the following functions:

(a)
$$f(x) = \sin(3\tan(2x))$$

(b)
$$f(x) = \sqrt{1 + \sqrt{1 + \sqrt{x}}}$$

(c)
$$g(x) = \sec^2(5x)$$

(d)
$$g(x) = \frac{1 - \tan(2x)}{1 + x^e + e^x}$$

(e)
$$h(x) = (1+3^x)\cos(x^3)e^{2x-x^3}$$

3. (Harder) Evaluate $\lim_{x \to \pi/4} \frac{\tan x - 1}{x - \pi/4}$

1

MATH 140 - Solutions

Evaluate each of the following Limits % Limit

Thank you for attending my PREP101 session!

antoine.prep101@gmail.com

(a)
$$\lim_{x \to -2} \frac{\frac{x-2+4(x+3)}{x+3}}{\frac{x+2}{1}} = \lim_{x \to -2} \frac{x-2+4x+12}{x+3} = \frac{1}{x+2}$$

$$= \lim_{X \to -2} \frac{5x + 10}{(x+3)(x+2)} = \lim_{X \to -2} \frac{5(x+2)}{(x+3)(x+2)} = \lim_{X \to -2} \frac{5}{x+3} = 5$$

Note: If
$$f(x) = \frac{x-2}{x+3}$$
; the above limit is $f'(-2) = \lim_{x \to -2} \frac{f(x) - f(-2)}{x - (-2)}$

(b)
$$\lim_{X\to\infty} \sqrt{X^2-3X} - X = \lim_{X\to\infty} \sqrt{X^2-3X} - X \cdot \frac{\sqrt{X^2-3X} + X}{\sqrt{X^2-3X} + X}$$

$$= \lim_{X \to \infty} \frac{x^2 - 3x - x^2}{\sqrt{x^2 - 3x} + x} = \lim_{X \to \infty} \frac{-3x}{\sqrt{x^2 - 3x} + x}$$

$$= \lim_{X \to \infty} \frac{x (-3)}{\sqrt{x^2(1-\frac{3}{4})} + X} = \lim_{X \to \infty} \frac{x (-3)}{|x|\sqrt{1-\frac{3}{4}} + x} \qquad \text{note: } \sqrt{x^2} = |x|$$

$$= \lim_{X \to \infty} \frac{x (-3)}{\sqrt{x^2(1-\frac{3}{4})} + X} = \lim_{X \to \infty} \frac{x (-3)}{|x|\sqrt{1-\frac{3}{4}} + x} \qquad \text{note: } \sqrt{x^2} = |x|$$

note:
$$\sqrt{x^2} = |x|$$
 $|x| = \begin{cases} x & \text{if } x \neq 0 \\ -x & \text{if } x \neq 0 \end{cases}$

$$= \lim_{X \to \infty} \frac{\chi(-3)}{\chi \sqrt{1-3/\chi} + \chi} = \lim_{X \to \infty} \frac{\chi(-3)}{\chi \left(\sqrt{1-3/\chi} + 1\right)} = \lim_{X \to \infty} \frac{-3}{\sqrt{1-3/\chi} + 1} = \frac{-3}{2}.$$

Question: What happens if x - 00?

ANS
$$\lim_{x\to -\infty} \sqrt{x^2-3x} - x = \infty - (-\infty) = \infty$$

(c)
$$\lim_{X\to 0} \frac{\sin(2X).\sin(2X).\sin(3X)}{x.x.x.x.\cos(3X)}$$

Thank you for attending my **PREP101** session!
Best of luck!

antoine.prep101@gmail.com

$$= \lim_{X \to 0} \frac{sin(2x)}{2x} \cdot \frac{sin(2x)}{2x} \cdot \frac{sin(3x)}{3x} \cdot \frac{(2x)(3x)}{(2x)}$$

$$\frac{1}{\cos 0}$$
 $\frac{1}{\cos 0}$ $\frac{1}{\cos 0}$ $\frac{1}{\cos 0}$

Derive each of the following functions

(a)
$$f'(x) = \cos(3\tan(2x)) \cdot (3\tan(2x))'$$

 $= \cos(3\tan(2x)) \cdot 3 \sec^2(2x) \cdot (2x)'$
 $= \cos(3\tan(2x)) \cdot 3 \sec^2(2x) \cdot (2)$
 $= 6 \cos(3\tan(2x)) \cdot \sec^2(2x)$

(b)
$$f(x) = \frac{1}{2\sqrt{1+\sqrt{1+\sqrt{x}}}} \cdot (1+\sqrt{1+\sqrt{x}})'$$
$$= \frac{1}{2\sqrt{1+\sqrt{1+\sqrt{x}}}} \cdot \frac{1}{2\sqrt{1+\sqrt{x}}} \cdot (1+\sqrt{x})'$$

$$=\frac{1}{2\sqrt{1+\sqrt{1+\sqrt{2}}}}\cdot\frac{1}{2\sqrt{1+\sqrt{2}}}\cdot\frac{1}{2\sqrt{2}}$$

(c)
$$g'(x) = 2 \operatorname{Sec}(5x) (\operatorname{sec}(5x))'$$

= $2 \operatorname{Sec}(5x) \cdot \operatorname{Sec}(5x) \cdot \tan(5x) \cdot (5x)'$
= $2 \operatorname{sec}^2(5x) \cdot \tan(5x) \cdot 5$
= $10 \operatorname{Sec}^2(5x) \cdot \tan(5x)$

Thank you for attending my **PREP101** session!
Best of luck!

antoine.prep101@gmail.com

$$(d)g(x) = \frac{(1 - \tan(2x))'(1 + x^{e} + e^{x}) - (1 - \tan(2x)) \cdot (1 + x^{e} + e^{x})'}{(1 + x^{e} + e^{x})^{2}}$$

$$g'(x) = \frac{-2 \sec^2(2x) (1 + x^e + e^x) - (1 - \tan(2x)) (1 + e^{x^{e-1}} + e^x)}{(1 + x^e + e^x)^2}$$

(e)
$$h'(x) = \frac{\left((1+3^{x})\cos(x^{3})\right)'e^{2x-x^{3}} + (1+3^{x})\cos(x^{3}).(e^{2x-x^{3}})'}{e^{2x-x^{3}}\cos(x^{3}) + (1+3^{x})\cos(x^{3}).(e^{2x-x^{3}})'}$$

$$= \frac{\left((1+3^{x})^{2}\cos(x^{3}) + (1+3^{x})(\cos(x^{3}))^{2}\right)e^{2x-x^{3}} + (1+3^{x})\cos(x^{3}).e^{2x-x^{3}}}{(2-3x^{3})}$$

$$= \left(3^{x} \ln 3 \cos(x^{3}) - (1+3^{x}) \sin(x^{3}) \cdot (3x^{2})\right) e^{2x-x^{3}} + (1+3^{x}) \cos(x^{3}) \cdot e^{2x-x^{3}}$$

$$= 3^{x} \ln 3 \cos(x^{3}) \cdot e^{2x-x^{3}} - (1+3^{x}) \sin(x^{3}) \cdot (3x^{2}) \cdot e^{2x-x^{3}} + (1+3^{x}) \cos(x^{3}) \cdot e^{2x-x^{3}} (2-3x^{2})$$

NOTE:
$$(FG)' = FG' + FG'$$

 $(FGH)' = FGH + FGH' + FGH'$

3. Remember the definition of the derivative

$$f'(a) = \lim_{x \to a} \frac{f(x) - f(a)}{x - a}$$

Here
$$f(x) = \tan x \quad \alpha = \pi/4$$

$$f'(\pi/4) = \lim_{X \to \pi/4} \frac{\tan x - \tan(\pi/4)}{x - \pi/4}$$

$$= \lim_{X \to \pi/4} \frac{\tan x - 1}{x - \pi/4} = \sec^2(\pi/4) = (\sqrt{2})^2 = 2.$$