


A.


Cells called
Median UMI / cell
Median genes / cell
Mean mito %
Doublet rate %
GEX01
GEX02
GEX09
GEX10


B.


Male (ZZ)
Female (ZW)
W-gene UMI fraction
(ZW female ↑, ZZ male ≈ 0)
Z-gene UMI fraction
(ZZ male ≈ 2× ZW female)
donor7 early/GFP+ (n=1,959)
donor0 early/GFP+ (n=1,618)
donor9 early/WT (n=3,744)
donor10 early/WT (n=4,323)
donor1 early/WT (n=2,026)
donor11a early/GFP+ (n=1,425)
donor6 early/WT (n=3,645)
donor4 early/GFP+ (n=3,192)
donor2b HH25/WT (n=2,661)
donor2a HH25/WT (n=2,496)
donor8 HH25/GFP+ (n=1,543)
donor11b HH25/GFP+ (n=284)


C.


GEX01 ×GEX02 same 0.94 diff 0.42
GEX0S-chick ×GEX01 same 0.84 diff 0.31
GEX0S-chick ×GEX02 same 0.84 diff 0.31
pseudobulk AF correlation r
embryo (same 12 across all lanes)

D.


Group A
Group B
HH13 (4)
HH15-16 (4)
HH25 (4)
Cells per half
Left (designed)
Right (designed)
half — side not per-embryo resolvable

E.


E1
Cells per library
E2
ATAC raw reads / cell
E3
ATAC duplicate rate / cell
E4
HQ fragments / cell
E5
TSS-fragment fraction / cell
E6
FRIP — arc peaks / cell
E7
AMULET doublet rate (%)
ARC01
ARC02
ARC0S-chick
ARC09
ARC10