



universität
wien

BACHELORARBEIT

Titel der Bachelorarbeit

Analytic sets

Verfasser

Charlie Ahrendts

angestrebter akademischer Grad

Bachelor of Science (BSc.)

Wien, im Monat Juni 2013

Studienkennzahl lt. Studienblatt: UA 033621

Studienrichtung lt. Studienblatt: Mathematik

Betreuer: Dipl.-Phys. Dipl.-Math. Dr. Peter Elbau, Privatdoz.

Abriss

Dies ist ein Template für Abschlussarbeiten an der Fakultät für Mathematik der Universität Wien

Abstract

This is a template for theses at the Faculty of Mathematics of the University of Vienna.

Contents

1	A short history lesson	1
2	Polish spaces and Analytic sets	1
3	Borel-gedöns	5
4	Measurability	6
5	Alternative description	6
5.1	Souslin operation	6
5.2	Luzins construction ??	6
5.3	Projections	7
6	K-analytic sets??? prettyprettyplease?	7

1 A short history lesson

As it is so often the case with new and interesting discoveries, mathematical areas of study often come about unplanned or on accident. One prime example of this is the field of Analytic sets. Studying it provides us, next to some important results in among others set theory, probability theory and functional analysis with a deeper understanding of the way mathematics is done in practice.

Our story starts in the Year 1905 with famous French mathematician Henri Lebesgue and his paper "Sur les fonctions représentables analytiquement" [Leb05]. In it, Lebesgue proved many statements of great importance. He did however include one small remark which stated, without proof, that the Borel sets are closed under Projections.

About 10 years later, Russian mathematician Nikolai Luzin ordered his student Souslin to study Lebesgues paper. When the still very young Souslin returned, saying he had found an error in Lebesgues work, Luzin and his colleague Sierpiński to their credit believed their student. [Sie50] Soon after, they would be the ones leading the charge on this young new field of mathematics that had just opened itself up to them. As Souslin had correctly identified, there was no justification of the claim that projections of Borel sets would again be Borel.

In a short publication in the French Journal "Comptes Rendu" in 1917, Souslin announced the falsity of Lebesgues statement and gave a rough idea of his proof. [cite sur une] These new sets that arise out of projections of Borel sets are called Analytic sets or Souslin sets.

Much more thorough treatments of the topic as well as the first construction of an analytic set that is not Borel were later given by Luzin and Sierpinski after Souslins early death in 19 [cite sur quelques proprietes. . .], [Lus23] [Lus27] It took until after the end of World War I for western mathematicians to start engaging with this new field. But once they did, it quickly started to become apparent that Analytic sets were going to become a very important tool for many different areas of mathematics. [Mos87] [RS80].

This thesis aims to give an overview over some of the basic concepts, theorems and proofs in the field of Analytic sets. Through chapters 2-4, we will mainly follow the conventions and proofs of [Coh13]

2 Polish spaces and Analytic sets

As previously stated, Analytic sets arose out of the question, whether Projections of Borel sets are again Borel and Souslins counterproof of that claim. So if we seek to study the greater class of sets that is obtained as projections of Borel sets, it would only make sense to define them exactly as such. As it turns out however,

there are many different ways to define these sets, all of them useful in their own ways. The one that will be used throughout the following chapters is one that allows for particularly nice versions of many of the basic results considered here. We will define Analytic sets as continuous images of Polish spaces, named in honor of the Polish mathematicians who were the first to extensively study them. [cite]

Definition (Polish space). A topological space X is called a Polish space, if it is completely metrizable and separable (contains a countable dense subset)

Interesting to note here is the difference between a complete metric space and a completely metrizable space. For the latter, we only require the existence of a complete metric on X , but we do not need to choose a concrete one. This means of course that all complete metric spaces are Polish

There are a few different ways to view analytic sets, as well as different ways to define them. Souslin, for example in his early works on the topic defined them as arising from a series of unions and intersections of certain families of sets. We shall here however stick to the most common, and often most useful definition, which is that of Analytic sets being continuous images of Polish spaces

Definition (Analytic set). Let X be a Polish space, $A \subset X$. We call A analytic, if there exists a Polish space Y and $f : Y \rightarrow X$ continuous, such that $f(Y) = A$

Throughout this section, we will use some standard results about topological and metric spaces without proof. These are:

Corollary 2.1. *In metric spaces, separability and second countability are equivalent*

Corollary 2.2. *closed subsets of complete metric spaces are again complete metric spaces*

Proofs for these theorems can be found in [cite/Anhang]

Lemma 2.3. *Finite and countable products of Polish spaces are Polish.*

Proof. Let X_1, X_2, \dots be a sequence of (nonempty) Polish spaces. We can choose a complete metric \overline{d}_i for each of the spaces X_i . Now for each i , let $d_i(x, y) := \min \{1, \overline{d}_i(x, y)\}$

This again defines a complete metric with the additional property that $d_i(x, y) \leq 1$ for all $x, y \in X_i$ [Genauer beweisen? Anhang?] This new metric retains only information about small distances in \overline{d}_i .

Now we can turn towards the cartesian product $X := \prod X_i$. Let $d(x, y) :=$

$\sum \frac{1}{2^n} d_i(x_i, y_i)$ where $x = (x_1, x_2, \dots), y = (y_1, y_2, \dots) \in X$. This sum converges for all $x, y \in X$ since we constructed the d_i to be bounded by 1. Moreover, d defines a metric on X . This is easily seen by the fact that positive definiteness, symmetry and the triangle inequality all hold in each term of our sum individually and thus for the whole sum. The topology generated by d is exactly the product topology on X [Genauer?].

[Ab hier nicht im Buch, hoffe der Beweis passt so]

To show that d is indeed complete, we take an arbitrary Cauchy sequence $(x_n)_{n \in \mathbb{N}}$ in X . The projection onto each space X_i is also a Cauchy sequence x_{i_n} . Since X_i is complete, x_{i_n} converges to some value x_i . Let $x := (x_1, x_2, \dots)$ be the componentwise limit of our Cauchy sequences. We need to show that x_n converges to x in the metric d :

Choose an arbitrary $\varepsilon > 0$ and $m \in \mathbb{N}$, such that $\sum_{i \geq m} \frac{1}{2^i} \leq \frac{\varepsilon}{2}$. For each $i \leq m$, we can find an N_i , such that $d_i(x_{i_n}, x_i) \leq \frac{\varepsilon}{2}$ for all $n \geq N_i$. If we choose $N := \max_{i \leq m} \{N_i\}$, we get the following estimate for $n \geq N$:

$$\begin{aligned} d(x_n, x) &= \sum \frac{1}{2^i} d_i(x_{i_n}, x_i) \\ &= \sum_{i \leq m} \frac{1}{2^i} d_i(x_{i_n}, x_i) + \sum_{i \geq m} \frac{1}{2^i} d_i(x_{i_n}, x_i) \\ &\leq \sum_{i \leq m} \frac{1}{2^i} \frac{\varepsilon}{2} + \sum_{i \geq m} \frac{1}{2^i} \\ &\leq \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon \end{aligned}$$

So we get convergence of x_n and thus completeness of the metric space (X, d)

What remains to be shown is separability. We want to make use of the equivalence of separability and second countability for metric spaces. [cite] For each i , we can find a countable Basis \mathcal{U}_i for the topology on X_i . The sets of the form $U_1 \times U_2 \times \dots \times U_N \times X_{N+1} \times X_{N+2} \times \dots$ form a countable Basis for the product topology on X , so X is a separable metrisable space and thus Polish.

□

Corollary 2.4. *Open and closed subsets of Polish spaces are Polish*

Proof. Let X be a Polish space, A be an open or closed subset of X . By Corollary 2.1, X has a countable basis of open sets for its Topology. Then the restrictions of those sets to A form a countable basis for the subspace topology on A . By the same equivalence, we get separability of A .

If A is closed, it is according to Corollary 2.2 completely metrizable by any

complete metric on X , restricted to A .

Now only the case of A being an open subset of X remains. Let

$$d_0(x, y) := d(x, y) + \left| \frac{1}{d(x, A^C)} - \frac{1}{d(y, A^C)} \right|.$$

where d is a complete metric on X and $d(x, A^C) := \inf \{d(x, z) : z \in A^C\}$.
 d_0 defines a metric on A . [proof?]

□

Theorem 2.5. *Open and closed subsets of Polish spaces are analytic*

Proof. Let X be a Polish space, $A \subset X$ open or closed. We know that A , as a space is polish. So there exists a polish space Y and a continuous function $f : Y \rightarrow A$, such that $f(Y) = A$. Let $\iota : A \hookrightarrow X$ be the canonical embedding of A into X . Then we can define $\tilde{f} := f \circ \iota$. Then $\tilde{f} : Y \rightarrow X$ and $\tilde{f}(Y) = A$ hold. So A is an analytic subset of X

□

Two particular spaces that are of great interest in the study of analytic sets are $\mathbb{N}^{\mathbb{N}}$ and $\{0, 1\}^{\mathbb{N}}$. We will see that the polish space Z in our definition of analytic set can always be replaced by the space $\mathbb{N}^{\mathbb{N}}$. But first we need to verify that they are in fact polish:

Theorem 2.6. $\mathbb{N}^{\mathbb{N}}$ is polish

Proof. by \mathbb{N} we always mean the natural numbers together with the discrete topology in which every subset is open. Since \mathbb{N} is countable, separability immediately follows. The discrete metric $d(m, n) = 1 - \delta_{mn}$, which equals is a complete metric on \mathbb{N} . In this metric the only Cauchy sequences are those that are eventually constant which obviously converge. It now follows from 2.3, that the cartesian product $\mathbb{N}^{\mathbb{N}}$ is also Polish.

□

Theorem 2.7. $\{0, 1\}^{\mathbb{N}}$ is polish

Proof. The proof for this is equivalent to that of $\mathbb{N}^{\mathbb{N}}$ being polish. We again choose the discrete topology and discrete metric on $\{0, 1\}$ and use 2.3 to get that $\{0, 1\}^{\mathbb{N}}$ is Polish.

□

Theorem 2.8. *Let X be a Polish space. Then there is a continuous function $f : \mathbb{N}^{\mathbb{N}} \rightarrow X$, such that $f(\mathbb{N}^{\mathbb{N}}) = X$*

Theorem 2.9. *Let A be a nonempty analytic subset of a polish space X . Then there exists continuous function $f : \mathbb{N}^{\mathbb{N}} \rightarrow X$, such that $f(\mathbb{N}^{\mathbb{N}}) = A$*

3 Borel-gedöns

The sets maybe closest in nature to the analytic sets are the Borel sets. This of course is a natural consequence of the way analytic sets were discovered. In some sense, Analytic sets are a just a generalization of the Borel sets which are closed under projections. We will try to formalise this notion as well as some other results about these two classes of sets in this section.

Definition (Borel set). We call a subset of a topological space Borel, if it is a member of \mathcal{B} , the σ -Algebra generated by the open sets

Theorem 3.1. *Let B be a Borel subset of a Polish space X . Then B is analytic*

Definition (Zero-Dimensional space?). .

Theorem 3.2. *Let B be a Borel subset of a Polish space X . Then there exists a zero-dimensional space Z , such that $f(Z) = B$*

Theorem 3.3. *Let B be an uncountable Borel subset of a polish space X . Then B contains a subset which is homeomorphic to $\{0,1\}^{\mathbb{N}}$*

Theorem 3.4 (Separation theorem). .

Theorem 3.5. *Let A be a subset of a polish space X . If A and A^C are analytic, then A is Borel.*

Theorem 3.6. *There exists.*

Definition (Borel isomorphic). We call two Borel subsets A, B of a Polish space X Borel isomorphic, if there exists a bijective, Borel measurable function $f : A \rightarrow B$

Theorem 3.7. *Two Borel subsets of a polish space X are Borel isomorphic iff they have the same cardinality*

We have seen that all Borel sets are analytic, but have not yet said anything about the converse. So we turn to the foundational theorem, introduced by souslin, that makes this field of analytic sets worth studying:

Definition (Universal set).

Theorem 3.8 (Souslin). *There exist Analytic sets which are not Borel sets.*

Kechris p.85.

□

Example. As almost all 'nice' sets are Borel, we can assume that most constructions of Analytic non-borelian sets are fairly complicated. One of the earliest such examples was provided in 1936 by polish mathematician Stefan Mazurkiewicz

[cite]:

Let \mathbb{R}^I denote the space of real-valued functions, which are continuous in the closed interval $I := [0, 1]$. Let Γ be the set of functions $f \in \mathbb{R}^I$, which are differentiable in I . This set is Co-analytic, meaning it is the complement of an analytic set, but is itself not analytic. Thus, its complement cannot be Borel (or Γ would be Borel as well, and thus Analytic, leading to a contradiction).

The proof of this is rather long and technical and beyond the scope of this thesis, but can be found in [cite]

4 Measurability

Definition (μ -Measurable).

Definition (Universally measurable).

Theorem 4.1. *Every finite Borel measure on Polish space is regular*

Theorem 4.2. *Let B be an analytic subset of a polish space X . Then B is universally measurable.*

Theorem 4.3. *Let (X, \mathcal{A}) , (Y, \mathcal{B}) be measurable spaces, that is, spaces endowed with a σ -Algebra. Let \mathcal{A}_* and \mathcal{B}_* be the σ -Algebras of universally measurable sets. If $f : X \rightarrow Y$ is $\mathcal{A} - \mathcal{B}$ -measurable, then it is also $\mathcal{A}_* - \mathcal{B}_*$ -measurable*

Definition (Analytic space).

Theorem 4.4. *Let X, A be analytic meas. space, Y polish, f measurable then $f(A)$ analytic.*

5 Alternative description

5.1 Souslin operation

[Rogers, p.319 ?]

5.2 Luzins construction ??

[Sur les ensembles analytiques]

5.3 Projections

6 K-analytic sets??? prettyprettyplease?

References

- [Coh13] D.L. Cohn. *Measure Theory: Second Edition*. Birkhäuser Advanced Texts Basler Lehrbücher. Springer New York, 2013. URL: <https://books.google.at/books?id=PEC3BAAQBAJ>.
- [Leb05] Henri Lebesgue. “Sur les fonctions représentables analytiquement”. fre. In: *Journal de Mathématiques Pures et Appliquées* 1 (1905), pp. 139–216. URL: <http://eudml.org/doc/234955>.
- [Lus23] Nicolas Lusin. “Sur un ensemble non mesurable B”. fre. In: *Journal de Mathématiques Pures et Appliquées* (1923), pp. 53–72. URL: <http://eudml.org/doc/234332>.
- [Lus27] Nicolas Lusin. “Sur les ensembles analytiques”. fre. In: *Fundamenta Mathematicae* 10.1 (1927), pp. 1–95. URL: <http://eudml.org/doc/211183>.
- [Mos87] Y N Moschovakis. *Descriptive Set Theory*. Studies in Logic and the Foundations of Mathematics. North-Holland, Jan. 1987.
- [RS80] C.A. Rogers and London Mathematical Society. *Analytic Sets*. London Mathematical Society Symposia Series. Academic Press, 1980. URL: <https://books.google.at/books?id=zQHvAAAAAAAJ>.
- [Sie50] W. Sierpiński. *Les ensembles projectifs et analytiques*. fre. Gauthier-Villars, 1950. URL: <http://eudml.org/doc/192631>.