

# Methods 2 - 4

**Chris Mathys**



BSc Programme in Cognitive Science

Spring 2022

# Math basics – Gill, chapter 1

- Arithmetic
- Notation
- Functions
- Polynomials
- Exponents
- Logarithms

## The Summation Operator

- If  $X_1, X_2, \dots, X_n$  are  $n$  numerical values,
- then their sum can be represented by  $\sum_{i=1}^n X_i$ ,
- where  $i$  is an indexing variable to indicate the starting and stopping points in the series  $X_1, X_2, \dots, X_n$ .

## The Product Operator

- If  $X_1, X_2, \dots, X_n$  are  $n$  numerical values,
- then their product can be represented by  $\prod_{i=1}^n X_i$ ,
- where  $i$  is an indexing variable to indicate the starting and stopping points in the series  $X_1, X_2, \dots, X_n$ .

# Numbers

Symbol	Explanation
$\mathfrak{R}$	the set of real numbers
$\mathfrak{R}^+$	the set of positive real numbers
$\mathfrak{R}^-$	the set of negative real numbers
$\mathcal{I}$	the set of integers
$\mathcal{I}^+$ or $\mathbb{Z}^+$	the set of positive integers
$\mathcal{I}^-$ or $\mathbb{Z}^-$	the set of negative integers
$\mathbb{Q}$	the set of rational numbers
$\mathbb{Q}^+$	the set of positive rational numbers
$\mathbb{Q}^-$	the set of negative rational numbers
$\mathcal{C}$	the set of complex numbers (those based on $\sqrt{-1}$ ).

# Symbols

Symbol	Explanation
$\neg$	logical negation statement
$\in$	is an element of, as in $3 \in \mathcal{I}^+$
$\ni$	such that
$\therefore$	therefore
$\because$	because
$\implies$	logical “then” statement
$\iff$	if and only if, also abbreviated “iff”
$\exists$	there exists
$\forall$	for all
$\oslash$	between
$\parallel$	parallel
$\angle$	angle

# Symbols

Symbol	Explanation
$\emptyset$	the empty set (sometimes used with the Greek phi: $\phi$ )
$\cup$	union of sets
$\cap$	intersection of sets
$\setminus$	subtract from set
$\subset$	subset
$\complement$	complement

# Symbols

Symbol	Explanation
$\emptyset$	the empty set (sometimes used with the Greek phi: $\phi$ )
$\cup$	union of sets
$\cap$	intersection of sets
$\setminus$	subtract from set
$\subset$	subset
$\complement$	complement



# Symbols

Symbol	Explanation
$\propto$	is proportional to
$\doteq$	equal to in the limit (approaches)
$\perp$	perpendicular
$\infty$	infinity
$\infty^+, +\infty$	positive infinity
$\infty^-, -\infty$	negative infinity
$\Sigma$	summation
$\Pi$	product
$\lfloor \rfloor$	floor: round down to nearest integer
$\lceil \rceil$	ceiling: round up to nearest integer
$ $	given that: $X Y = 3$

# Symbols

Symbol	Explanation
$\vee$	maximum of two values
$\max()$	maximum value from list
$\wedge$	minimum of two values
$\min()$	minimum value from list
$\operatorname{argmax}_x f(x)$	the value of $x$ that maximizes the function $f(x)$
$\operatorname{argmin}_x f(x)$	the value of $x$ that minimizes the function $f(x)$

# Symbols

Symbol	Meaning
$<$	less than
$\leq$	less than or equal to
$\ll$	much less than
$>$	greater than
$\geq$	greater than or equal to
$\gg$	much greater than
$\approx$	approximately the same
$\approx$	approximately equal to
$\approx$	approximately less than (also $\lesssim$ )
$\approx$	approximately greater than (also $\gtrsim$ )
$\equiv$	equivalent by assumption


# Functions

## Properties of Functions, Given for $g(x) = y$

- A function is **continuous** if it has no “gaps” in its mapping from  $x$  to  $y$ .
- A function is **invertible** if its reverse operation exists:  
 $g^{-1}(y) = x$ , where  $g^{-1}(g(x)) = x$ .

# Functions

Fig. 1.2. A RELATION THAT IS NOT A FUNCTION



## Key Properties of Powers and Exponents

- Zero Property  $x^0 = 1$
- One Property  $x^1 = x$
- Power Notation  $\text{power}(x, a) = x^a$
- Fraction Property  $\left(\frac{x}{y}\right)^a = \left(\frac{x^a}{y^a}\right) = x^a y^{-a}$
- Nested Exponents  $(x^a)^b = x^{ab}$
- Distributive Property  $(xy)^a = x^a y^a$
- Product Property  $x^a \times x^b = x^{a+b}$
- Ratio Property  $x^{\frac{a}{b}} = (x^a)^{\frac{1}{b}} = \left(x^{\frac{1}{b}}\right)^a = \sqrt[b]{x^a}$

## Basic Properties of Logarithms

→ Zero/One       $\log_b(1) = 0$

→ Multiplication       $\log(x \cdot y) = \log(x) + \log(y)$

→ Division       $\log(x/y) = \log(x) - \log(y)$

→ Exponentiation       $\log(x^y) = y \log(x)$

→ Basis       $\log_b(b^x) = x$ , and  $b^{\log_b(x)} = x$