

# Edexcel A Level Maths

Core 1, Core 2, Core 3, Core 4 and two units from (Decision 1, Decision 2, Mechanics 1, Mechanics 2, Statistics 1 and Statistics 2). Each unit is out of 100 UMS, giving a total of 600 UMS. 480 and above is grade A. To obtain an A\*, total score must be 480 or more, and total of C3 and C4 must be 180 or more.

## Unit 1 Core 1

### Chapter 1 Sequence and Series 1

#### Lesson 1 Sequence and Summation

**Question 1** Experience: 10 Order: Level:

A sequence is defined by  $a_n = 3n^2 - 4$ , find the value of  $a_2$ .

**Solution 1**

$$a_2 = 3(2)^2 - 4 = 8$$

$$a_2 = -1$$

$$a_2 = 23$$

$$a_2 = 5$$

$$a_2 = 10$$

$$a_2 = 8$$

**Question 2** Experience: 25 Order: Level:

A sequence is defined by  $x_n = 3n^2 - 5n + 2$ , find the value of  $n$  such that  $x_n = 14$ .

**Solution 2**

$$x_n = 3n^2 - 5n + 2 = 14$$

$$3n^2 - 5n - 12 = 0 \quad S = -5 \quad P = -36$$

$$\left(n + \frac{4}{3}\right)(n - 3) = 0 \quad (4, -9) \quad \left(\frac{4}{3}, -3\right)$$

$$n = 3$$

$$n = 4$$

$$n = 2$$

$$n = 5$$

$$n = 6$$

$$n = 3$$

**Question 3** Experience: 15 Order: Level:

A sequence is defined by  $x_n = 6n - 3$ , find the value of  $x_3$  and  $x_5$ .

**Solution 3**

$$x_3 = 6(3) - 3 = 15$$

$$x_5 = 6(5) - 3 = 27$$

$$x_3 = 3 \quad x_5 = 15$$

$$x_3 = 9 \quad x_5 = 21$$

$$x_3 = 15 \quad x_5 = 27$$

$$x_3 = 3 \quad x_5 = 21$$

$$x_3 = 15 \quad x_5 = 27$$

**Question 4** Experience: 10 Order: Level:

A sequence is defined by  $X_n = 2n - 1$ , find the value of  $n$  such that  $a_n = 15$ .

**Solution 4**

$$15 = 2n - 1$$

$$n = 8$$

$$n = 6$$

$$n = 7$$

$$n = 3$$

$$n = 9$$

$$n = 8$$

**Question 5** Experience: 15 Order: Level:

A sequence is defined by  $u_n = an - b$ , find the sum of the first four terms in terms of  $a$  and  $b$ .

**Solution 5**

$$u_1 + u_2 + u_3 + u_4 = (a - b) + (2a - b) + (3a - b) + (4a - b) = 10a - 4b$$

$$10a - 6b$$

$$6a - 4b$$

$$6a - 6b$$

$$10a - 8b$$

$$10a - 4b$$

**Question 6** Experience: 15 Order: Level:

A sequence is defined by  $x_n = an^2 - 4$ , find the sum of the first three terms in terms of  $a$ .

**Solution 6**

$$x_1 + x_2 + x_3 = (a - 4) + (4a - 4) + (9a - 4) = 14a - 12$$

$$14a - 8$$

$$5a - 12$$

$$5a - 8$$

$$5a - 14$$

$$14a - 12$$

**Question 7** Experience: 25 Order: Level:

A sequence is defined by  $y_n = an^2 + bn + c$ , find the sum of the first three terms in terms of  $a, b$  and  $c$ .

**Solution 7**

$$y_1 + y_2 + y_3 = (a + b + c) + (4a + 2b + c) + (9a + 3b + c) = 14a + 6b + 3c$$

wrong choice

$$14a - 6b + 3c$$

$$6a + 6b + 3c$$

$$6a + 4b + 2c$$

$$6a + 4b + 3c$$

$$14a + 6b + 3c$$

**Question 8** Experience: 10 Order: Level:

A sequence is defined by  $x_n = 4n - b$ , find the third term in terms of  $b$ .

**Solution 8**

$$x_3 = 4(3) - b$$

$$x_3 = 12 - b$$

$$x_3 = 12 - 3b$$

$$x_3 = 6 - b$$

$$x_3 = 6 - 3b$$

$$x_3 = 8 - 3b$$

$$x_3 = 12 - b$$

**Question 9** Experience: 10 Order: Level:

A sequence is defined by  $U_n = \frac{a}{n} + b$ , find the fourth term in terms of  $a$  and  $b$ .

**Solution 9**

$$U_4 = \frac{a}{4} + b$$

$$U_4 = \frac{a + b}{4}$$

$$U_4 = \frac{a}{4} + 4b$$

$$U_4 = \frac{a}{4} + 2b$$

$$U_4 = \frac{a}{4} + b$$

$$U_4 = \frac{a + 4b}{8}$$

**Question 10** Experience: 15 Order: Level:

A sequence is defined by  $y_n = \frac{a - 3b}{n^2}$ , find the fifth term in terms of  $a$  and  $b$ .

**Solution 10**

$$y_5 = \frac{a-3b}{(5)^2}$$

$$y_5 = \frac{a-3b}{25}$$

$$y_5 = \frac{5a-3b}{25}$$

$$y_5 = \frac{5a-3b}{16}$$

$$y_5 = \frac{a-3b}{16}$$

$$y_5 = \frac{5a-b}{25}$$

$$y_5 = \frac{a-3b}{25}$$

**Question 11** Experience: 50 Order: Level:

A sequence is defined by  $U_n = an + 2b$ , given the Sum of the first four terms is 26 and the fifth term is 9, find the values of  $a$  and  $b$ .

**Solution 11**

$$S_4 = (a + 2b) + (2a + 2b) + (3a + 2b) + (4a + 2b)$$

$$S_4 = 10a + 8b \quad S_4 = 26$$

$$10a + 8b = 26$$

$$5a + 4b = 13 \quad (1)$$

$$U_5 = 5a + 2b \quad U_5 = 9$$

$$5a + 2b = 9 \quad (2)$$

$$(1) - (2) \quad 5a + 4b - (5a + 2b) = 13 - 9$$

$$2b = 4$$

$$b = 2$$

$$\text{sub into } (2) \quad 5a + 2(2) = 9$$

$$5a = 5$$

$$a = 1$$

$$a = 1 \quad b = 3$$

$$a = 2 \quad b = 3$$

$$a = 2 \quad b = 2$$

$$a = 3 \quad b = 2$$

$$a = 1 \quad b = 2$$

**Question 12** Experience: 25 Order: Level:

A sequence is defined by  $U_{n+1} = U_n - 4$ ,  $U_1 = 20$ , find the values of  $U_2$ ,  $U_3$  and  $U_4$ .

**Solution 12**

$$U_2 = U_1 - 4 = 20 - 4 = 16$$

$$U_3 = U_2 - 4 = 16 - 4 = 12$$

$$U_4 = U_3 - 4 = 12 - 4 = 8$$

$$U_2 = 16 \quad U_3 = 12 \quad U_4 = 8$$

$$U_2 = 12 \quad U_3 = 8 \quad U_4 = 4$$

$$U_2 = 12 \quad U_3 = 4 \quad U_4 = 0$$

$$U_2 = 16 \quad U_3 = 4 \quad U_4 = 4$$

$$U_2 = 16 \quad U_3 = 12 \quad U_4 = 8$$

**Question 13**      Experience: 25      Order:      Level:

A sequence is defined by  $X_{n+1} = X_n + 5$ ,  $X_4 = 17$ , find the values of  $X_1$ ,  $X_2$  and  $X_3$ .

**Solution 13**

$$X_4 = X_3 + 5$$

$$17 = X_3 + 5$$

$$X_3 = 12$$

$$X_3 = X_2 + 5$$

$$12 = X_2 + 5$$

$$X_2 = 7$$

$$X_2 = X_1 + 5$$

$$7 = X_1 + 5$$

$$X_1 = 2$$

$$X_1 = 2 \quad X_2 = 7 \quad X_3 = 12$$

$$X_1 = 5 \quad X_2 = 8 \quad X_3 = 11$$

$$X_1 = 5 \quad X_2 = 7 \quad X_3 = 9$$

$$X_1 = 5 \quad X_2 = 6 \quad X_3 = 10$$

$$X_1 = 2 \quad X_2 = 7 \quad X_3 = 12$$

**Question 14**      Experience: 30      Order:      Level:

A sequence is defined by  $a_{n+1} = (a_n)^2 - 4$ ,  $a_1 = 2$ , find the values of  $a_2$ ,  $a_3$  and  $a_4$ .

**Solution 14**

$$a_2 = (a_1)^2 - 4 = 4 - 4 = 0$$

$$a_3 = (a_2)^2 - 4 = 0 - 4 = -4$$

$$a_4 = (a_3)^2 - 4 = (-4)^2 - 4 = 16 - 4 = 12$$

$$a_2 = 0 \quad a_3 = -4 \quad a_4 = 12$$

$$a_2 = 4 \quad a_3 = -4 \quad a_4 = 8$$

$$a_2 = 4 \quad a_3 = 8 \quad a_4 = -8$$

$$a_2 = 0 \quad a_3 = 8 \quad a_4 = 12$$

$$a_2 = 0 \quad a_3 = -4 \quad a_4 = 12$$

**Question 15**      Experience: 25      Order:      Level:

A sequence is defined by  $y_{n+2} = 3y_{n+1} - y_n$ ,  $y_1 = 3$ ,  $y_2 = 2$ , find the values of  $y_3$ ,  $y_4$  and  $y_5$ .

**Solution 15**

$$y_3 = 3(y_2) - y_1 = 3(2) - 3 = 3$$

$$y_4 = 3(y_3) - y_2 = 3(3) - 2 = 7$$

$$y_5 = 3(y_4) - y_3 = 3(7) - 3 = 18$$

$$y_3 = 3 \quad y_4 = 7 \quad y_5 = 18$$

$$y_3 = 7 \quad y_4 = 4 \quad y_5 = 5$$

$$y_3 = 7 \quad y_4 = 8 \quad y_5 = 5$$

$$y_3 = 3 \quad y_4 = 7 \quad y_5 = 5$$

$$y_3 = 3 \quad y_4 = 7 \quad y_5 = 18$$

**Question 16**      Experience: 15      Order:      Level:

Calculate the following sum:

$$\sum_{r=2}^5 (r-1)$$

**Solution 16**

$$\sum_{r=2}^5 (r-1) = (2-1) + (3-1) + (4-1) + (5-1)$$

$$= 1 + 2 + 3 + 4$$

$$= 10$$

9

8

11

12

10

**Question 17**      Experience: 15      Order:      Level:

Calculate the following sum:

$$\sum_{r=4}^8 (r^2 - 2r + 1)$$

**Solution 17**

$$\begin{aligned}
& \sum_{r=4}^8 (r^2 - 2r + 1) \\
&= \sum_{r=4}^8 (r - 1)^2 \\
&= (4 - 1)^2 + (5 - 1)^2 + (6 - 1)^2 + (7 - 1)^2 + (8 - 1)^2 \\
&= 9 + 16 + 25 + 36 + 49 \\
&= 135
\end{aligned}$$

137

128

130

136

135

**Question 18**      Experience: 30      Order:      Level:

Calculate the following sum:

$$\sum_{r=5}^9 U_r \quad U_r = 3r^2 + 4$$

**Solution 18**

$$\begin{aligned}
& \sum_{r=5}^9 U_r \\
&= \sum_{r=5}^9 3r^2 + 4 \\
&= (3(5)^2 + 4) + (3(6)^2 + 4) + (3(7)^2 + 4) + (3(8)^2 + 4) + (3(9)^2 + 4) \\
&= 785
\end{aligned}$$

795

790

780

800

785

**Question 19**      Experience: 30      Order:      Level:

Calculate the following sum:

$$\sum_{r=1}^3 a_r \quad a_r = 4r - 1$$

**Solution 19**

$$\begin{aligned}
 & \sum_{r=1}^3 a_r \\
 &= \sum_{r=1}^3 4r - 1 \\
 &= (4(1) - 1) + (4(2) - 1) + (4(3) - 1) \\
 &= 21
 \end{aligned}$$

22

19

20

18

21

**Question 20**      Experience: 45      Order:      Level:

A sequence is defined by  $U_{n+1} = 3(U_n - 1)$ ,  $U_1 = 2$ , find the following sum:  $\sum_2^4 (U_r + 2)^2$

**Solution 20**

$$U_2 = 3(2 - 1)$$

$$U_2 = 3$$

$$U_3 = 3(3 - 1)$$

$$U_3 = 6$$

$$U_4 = 3(6 - 1)$$

$$U_4 = 15$$

$$\begin{aligned}
 \sum_2^4 (U_r + 2)^2 &= (U_2 + 2)^2 + (U_3 + 2)^2 + (U_4 + 2)^2 \\
 &= (3 + 2)^2 + (6 + 2)^2 + (15 + 2)^2 \\
 &= 5^2 + 8^2 + 17^2 \\
 &= 378
 \end{aligned}$$

380

377

379

381

378

**Question 21**      Experience: 15      Order:      Level:

Calculate the following sum:

$$\sum_{r=1}^4 (2r + 4)$$

**Solution 21**



$$\begin{aligned}
& \sum_{r=1}^4 (2r + 4) \\
&= \sum_{r=1}^4 2(r + 2) \\
&= 2 \sum_{r=1}^4 (r + 2) \\
&= 2[(1 + 2) + (2 + 2) + (3 + 2) + (4 + 2)] \\
&= 2(3 + 4 + 5 + 6) \\
&= 36
\end{aligned}$$

37

35

34

33

36

**Question 22**    Experience: 25    Order:    Level:

Calculate the following sum:

$$\sum_{r=3}^6 (r^2 - 1)$$

**Solution 22**

$$\begin{aligned}
& \sum_{r=3}^6 (r^2 - 1) \\
&= (3^2 - 1) + (4^2 - 1) + (5^2 - 1) + (6^2 - 1) \\
&= 8 + 15 + 24 + 35 \\
&= 82
\end{aligned}$$

81

80

83

84

82

**Question 23**    Experience: 15    Order:    Level:

Calculate the following sum:

$$\sum_{r=1}^{45} 2$$

**Solution 23**

$$\sum_{r=1}^{45} 2$$

$$= 2 + 2 + 2 + 2 + 2 + \dots + 2$$

$$= 2 \times 45$$

$$= 90$$

94

92

88

86

90

**Question 24**      Experience: 15      Order:      Level:

Calculate the following sum:

$$\sum_{r=1}^{100} 5$$

**Solution 24**

$$\sum_{r=1}^{100} 5$$

$$= 5 + 5 + 5 + 5 + 5 + 5 + \dots + 5$$

$$= 5 \times 100$$

$$= 500$$

495

490

480

500

485

**Question 25**      Experience: 25      Order:      Level:

A sequence is defined by  $U_n = 2n + 3$ , find the value of  $U_2$ ,  $U_4$  and  $U_5$ .

**Solution 25**

$$U_2 = 2(2) + 3 = 7$$

$$U_4 = 2(4) + 3 = 11$$

$$U_5 = 2(5) + 3 = 13$$

$$U_2 = 5 \quad U_4 = 9 \quad U_5 = 11$$

$$U_2 = 7 \quad U_4 = 9 \quad U_5 = 13$$

$$U_2 = 7 \quad U_4 = 10 \quad U_5 = 15$$

$$U_2 = 7 \quad U_4 = 11 \quad U_5 = 13$$

$$U_2 = 5 \quad U_4 = 11 \quad U_5 = 13$$

**Question 26**    Experience: 25    Order:    Level:

A sequence is defined by  $u_n = 2n^2 - 5n - 3$ , find the value of  $n$  such that  $u_n = 9$ .

**Solution 26**

$$u_n = 2n^2 - 5n - 3 = 9$$

$$2n^2 - 5n - 12 = 0 \quad S = -5 \quad P = -24$$

$$\left(n + \frac{3}{2}\right)(n - 4) = 0 \quad (3, -8) \quad \left(\frac{3}{2}, -4\right)$$

$$n = 4$$

$$n = 5$$

$$n = 2$$

$$n = 3$$

$$n = 6$$

$$n = 4$$

**Question 27**    Experience: 50    Order:    Level:

A sequence is defined by  $a_n = an^2 + b$ , given the Sum of the first five terms is  $-5$  and the sixth term is 4, find the values of  $a$  and  $b$ .

**Solution 27**

$$S_5 = (a + b) + (4a + b) + (9a + b) + (16a + b)$$

$$S_5 = 30a + 5b \quad S_5 = -5$$

$$30a + 5b = -5$$

$$6a + b = -1 \quad (1)$$

$$a_6 = 25a + b \quad a_6 = 4$$

$$36a + b = 4 \quad (2)$$

$$(2) - (1) \quad 36a + b - (6a + b) = 4 - (-1)$$

$$30a = 5$$

$$a = \frac{1}{6}$$

$$\text{sub into (1)} \quad 6\left(\frac{1}{6}\right) + b = -1$$

$$b = -2$$

$$a = \frac{1}{6} \quad b = 2$$

$$a = 1 \quad b = -2$$

$$a = 1 \quad b = 2$$

$$a = 2 \quad b = \frac{1}{6}$$

$$a = \frac{1}{6} \quad b = -2$$

## Lesson 2 Arithmetic Sequence 1

**Question 1** Experience: 30 Order: Level:

Evaluate  $\sum_{r=1}^{15} (5r + 2)$

**Solution 1**

$$\sum_{r=1}^{15} (5r + 2) = 7 + 12 + 17 + 22 + \dots + 77$$

$$a = 7 \quad l = 77 \quad n = 15$$

$$\sum_{r=1}^{15} (5r + 2) = \frac{15}{2}(7 + 77)$$

$$= 630$$

$$\sum_{r=1}^{15} (5r + 2) = 625$$

$$\sum_{r=1}^{15} (5r + 2) = 620$$

$$\sum_{r=1}^{15} (5r + 2) = 615$$

$$\sum_{r=1}^{15} (5r + 2) = 635$$

$$\sum_{r=1}^{15} (5r + 2) = 630$$

**Question 2** Experience: 30 Order: Level:

How many terms are there in the arithmetic sequence 19,21,23,...,87

**Solution 2**

$$a = 19 \quad d = 2$$

$$U_n = a + (n - 1)d$$

$$87 = 19 + (n - 1)2$$

$$n - 1 = 34$$

$$n = 35$$

$$n = 38$$

$$n = 37$$

$$n = 36$$

$$n = 34$$

$$n = 35$$

**Question 3** Experience: 30 Order: Level:

How many terms are there in the arithmetic sequence 21,26,31,...,256

**Solution 3**

$$a = 21 \quad d = 5$$

$$U_n = a + (n - 1)d$$

$$256 = 21 + (n - 1)5$$

$$n - 1 = 47$$

$$n = 48$$

$$n = 51$$

$$n = 47$$

$$n = 50$$

$$n = 49$$

$$n = 48$$

**Question 4** Experience: 35 Order: Level:

How many terms are there in the arithmetic sequence 88,86,84,...,22

**Solution 4**

Reverse the order of the sequence 22,24,26,28...88

$$a = 88 \quad d = 2$$

$$n = 36 \quad U_n = a + (n - 1)d$$

$$n = 35 \quad 88 = 22 + (n - 1)2$$

$$n = 32 \quad n - 1 = 33$$

$$n = 33$$

$$n = 34 \quad n = 34$$

**Question 5** Experience: 30 Order: Level:

Evaluate  $S = 1 + 2 + 3 + 4 + \dots + 50$

**Solution 5**

$$S = 1 + 2 + 3 + 4 + \dots + 50$$

$$S = 50 + 49 + 48 + 47 + \dots + 1$$

$$2S = 51 \times 50$$

$$S = \frac{51 \times 50}{2}$$

$$S = 1275$$

$$S = 1270$$

$$S = 1280$$

$$S = 1285$$

$$S = 1290$$

$$S = 1275$$

**Question 6** Experience: 30 Order: Level:

Evaluate  $T = 2 + 4 + 6 + 8 + \dots + 100$

**Solution 6**

$$T = 2 + 4 + 6 + 8 + \dots + 100$$

$$T = 100 + 98 + 96 + 94 + \dots + 2$$

$$2T = 102 \times 50$$

$$T = \frac{102 \times 50}{2}$$

$$T = 2550$$

$$T = 2565$$

$$T = 2560$$

$$T = 2555$$

$$T = 2545$$

$$T = 2550$$

**Question 7**      Experience: 30      Order:      Level:

Evaluate  $R = 1 + 3 + 5 + 7 + \dots + 99$

**Solution 7**

$$R = 1 + 3 + 5 + 7 + \dots + 99$$

$$R = 99 + 97 + 95 + 93 + \dots + 1$$

$$2R = 100 \times 100$$

$$R = \frac{100 \times 100}{2}$$

$$R = 5000$$

$$R = 5015$$

$$R = 5010$$

$$R = 5005$$

$$R = 4995$$

$$R = 5000$$

**Question 8**      Experience: 30      Order:      Level:

Evaluate  $S = 1 + 2 + 3 + 4 + \dots + 200$

**Solution 8**

$$S = 1 + 2 + 3 + 4 + \dots + 200$$

$$S = 200 + 199 + 198 + 197 + \dots + 1$$

$$2S = 201 \times 200$$

$$S = \frac{201 \times 200}{2}$$

$$S = 20100$$

$$S = 20115$$

$$S = 20110$$

$$S = 20105$$

$$S = 20095$$

$$S = 20100$$

**Question 9** Experience: 30 Order: Level:

Evaluate  $T = 102 + 104 + 106 + 108 + \dots + 200$

**Solution 9**

$$T = 102 + 104 + 106 + 108 + \dots + 200$$

$$T = 200 + 198 + 196 + 194 + \dots + 102$$

$$2T = 302 \times 50$$

$$T = \frac{302 \times 50}{2}$$

$$T = 7550$$

$$T = 7565$$

$$T = 7560$$

$$T = 7555$$

$$T = 7545$$

$$T = 7550$$

**Question 10** Experience: 40 Order: Level:

Find the sum of all numbers divisible by 5 between 1 and 300

**Solution 10**

$$300 \div 5 = 60$$

$$\Rightarrow \text{last term} = 300$$

$$S = 5 + 10 + 15 + \dots + 300$$

$$a = 5 \quad d = 5 \quad U_n = 300$$

$$U_n = a + (n - 1)d$$

$$300 = 5 + (n - 1)5$$

$$n = 60$$

$$S = \frac{n}{2}(a + l)$$

$$S = \frac{60}{2}(5 + 300)$$

$$S = 9165$$

$$S = 9160$$

$$S = 9155$$

$$S = 9145$$

$$S = 9150$$

**Question 11** Experience: 40 Order: Level:

Find the sum of all numbers divisible by 7 between 1 and 200

**Solution 11**

$$200 \div 7 = 28 \text{ remainder } 4$$

$$\Rightarrow \text{last term} = 7 \times 28 = 196$$

$$S = 7 + 14 + 21 + \dots + 196$$

$$a = 7 \quad d = 7 \quad U_n = 196$$

$$U_n = a + (n - 1)d$$

$$S = 2806 = 7 + (n - 1)7$$

$$S = 2856$$

$$S = 2849$$

$$S = 2835$$

$$S = \frac{n}{2}(a + l)$$

$$S = 2842$$

**Question 12** Experience: 45 Order: Level:

$$\text{Evaluate } S = 27 + 31 + 35 + 39 + \dots + 107$$

$$S = 2842$$

**Solution 12**

$$S = 27 + 31 + 35 + 39 + \dots + 107$$

$$a = 27 \quad d = 4 \quad U_n = 107$$

$$U_n = a + (n - 1)d$$

$$S = 1407 = 27 + (n - 1)4$$

$$S = 1386$$

$$S = 1393$$

$$S = 1400$$

$$S = \frac{n}{2}(a + l)$$

$$S = 1407$$

**Question 13** Experience: 45 Order: Level:

$$\text{Evaluate } T = 31 + 33 + 35 + 37 + \dots + 81$$

$$S = 1407$$

**Solution 13**

$$T = 31 + 33 + 35 + 37 + \dots + 81$$

$$a = 31 \quad d = 2 \quad U_n = 81$$

$$U_n = a + (n - 1)d$$

$$S = 1462 = 31 + (n - 1)2$$

$$S = 1460$$

$$S = 1458$$

$$S = 1454$$

$$S = \frac{n}{2}(a + l)$$

$$S = 1456$$

**Question 14** Experience: 45 Order: Level:

$$\text{Evaluate } R = 97 + 92 + 87 + 82 + \dots + 22$$

$$S = 1456$$

**Solution 14**



Reverse the sequence:  $R = 22 + 27 + 32 + 27 + \dots + 97$

$$R = 22 + 27 + 32 + 27 + \dots + 97$$

$$a = 22 \quad d = 5 \quad U_n = 97$$

$$U_n = a + (n - 1)d$$

$$97 = 22 + (n - 1)5$$

$$n = 16$$

$$S = \frac{n}{2}(a + l)$$

$$S = \frac{16}{2}(22 + 97)$$

$$S = 952$$

$$S = 950$$

$$S = 954$$

$$S = 948$$

$$S = 950$$

$$S = 952$$

**Question 15**      Experience: 30      Order:      Level:

Evaluate  $\sum_{r=9}^{35} (3r - 1)$

**Solution 15**

$$\sum_{r=9}^{35} (3r - 1) = 26 + 29 + 32 + 35 + \dots + 104$$

$$a = 26 \quad l = 104 \quad n = 27$$

$$\sum_{r=9}^{35} (3r - 1) = \frac{27}{2}(26 + 104)$$

$$= 1755$$

$$\sum_{r=9}^{35} (3r - 1) = 1760$$

$$\sum_{r=9}^{35} (3r - 1) = 1740$$

$$\sum_{r=9}^{35} (3r - 1) = 1745$$

$$\sum_{r=9}^{35} (3r - 1) = 1750$$

$$\sum_{r=9}^{35} (3r - 1) = 1755$$

**Question 16**      Experience: 30      Order:      Level:

Evaluate  $\sum_{r=1}^{20} (3r - 1)$

**Solution 16**

$$\sum_{r=1}^{20} (3r - 1) = 2 + 5 + 8 + 11 + \dots + 59$$

$$a = 2 \quad l = 59 \quad n = 20$$

$$\begin{aligned} \sum_{r=1}^{20} (3r - 1) &= \frac{20}{2}(2 + 59) \\ &= 610 \end{aligned}$$

$$\sum_{r=1}^{20} (3r - 1) = 625$$

$$\sum_{r=1}^{20} (3r - 1) = 605$$

$$\sum_{r=1}^{20} (3r - 1) = 615$$

$$\sum_{r=1}^{20} (3r - 1) = 620$$

$$\sum_{r=1}^{20} (3r - 1) = 610$$

**Question 17**      Experience: 30      Order:      Level:

Evaluate  $\sum_{r=21}^{45} (2r - 25)$

**Solution 17**

$$\sum_{r=21}^{45} (2r - 25) = 17 + 19 + 21 + 23 + \dots + 65$$

$$a = 17 \quad l = 65 \quad n = 25$$

$$\begin{aligned} \sum_{r=21}^{45} (2r - 25) &= \frac{25}{2}(17 + 65) \\ &= 1025 \end{aligned}$$

$$\sum_{r=21}^{45} (2r - 25) = 1020$$

$$\sum_{r=21}^{45} (2r - 25) = 1015$$

$$\sum_{r=21}^{45} (2r - 25) = 1010$$

$$\sum_{r=21}^{45} (2r - 25) = 1030$$

$$\sum_{r=21}^{45} (2r - 25) = 1025$$

**Question 18**      Experience: 20      Order:      Level:

The first three terms of an arithmetic sequence are 3,5,7, find  $U_{10}$

**Solution 18**

$$a = 3 \quad n = 10 \quad d = 5 - 3 = 2$$

$$U_n = a + (n - 1)d$$

$$U_{10} = 3 + (10 - 1)2 = 21$$

$$U_{10} = 20$$

$$U_{10} = 17$$

$$U_{10} = 18$$

$$U_{10} = 19$$

$$U_{10} = 21$$

**Question 19**    Experience: 20    Order:    Level:

The first four terms of an arithmetic sequence are 5,9,13,17, find  $A_7$

**Solution 19**

$$a = 5 \quad n = 7 \quad d = 9 - 5 = 4$$

$$A_n = a + (n - 1)d$$

$$A_7 = 5 + (7 - 1)4 = 29$$

$$A_7 = 28$$

$$A_7 = 27$$

$$A_7 = 30$$

$$A_7 = 26$$

$$A_7 = 29$$

**Question 20**    Experience: 20    Order:    Level:

The first three terms of an arithmetic sequence are 22,19,16, find  $X_6$

**Solution 20**

$$a = 22 \quad n = 6 \quad d = 22 - 19 = 3$$

$$X_n = a + (n - 1)d$$

$$X_7 = 22 + (6 - 1)3 = 37$$

$$A_7 = 28$$

$$A_7 = 27$$

$$A_7 = 30$$

$$A_7 = 26$$

$$A_7 = 29$$

**Question 21**    Experience: 40    Order:    Level:

$a_n$  is an arithmetic sequence, given that  $a_3 = 13$  and  $a_6 = 19$ , find  $a_{11}$

**Solution 21**

$$a_n = a + (n - 1)d$$

$$a_3 = a + (3 - 1)d = a + 2d = 13 \quad (1)$$

$$a_6 = a + (6 - 1)d = a + 5d = 19 \quad (2)$$

$$(2) - (1) \quad a + 5d - (a + 2d) = 19 - 13$$

$$3d = 6$$

$$d = 2$$

$$\text{Sub into (1)} \quad a + 2(2) = 13$$

$$a = 9$$

$$a_{11} = 9 + (11 - 1)2 = 29$$

$$a_{11} = 25$$

$$a_{11} = 26$$

$$a_{11} = 27$$

$$a_{11} = 28$$

$$a_{11} = 29$$

**Question 22**      Experience: 40      Order:      Level:

$U_n$  is an arithmetic sequence, given that  $U_4 = 25$  and  $U_9 = 40$ , find  $U_{13}$

**Solution 22**

$$U_n = a + (n - 1)d$$

$$U_4 = a + (4 - 1)d = a + 3d = 25 \quad (1)$$

$$U_9 = a + (9 - 1)d = a + 8d = 40 \quad (2)$$

$$(2) - (1) \quad a + 8d - (a + 3d) = 40 - 25$$

$$5d = 15$$

$$d = 3$$

$$\text{Sub into (1)} \quad a + 3(3) = 25$$

$$a = 16$$

$$U_{13} = 51$$

$$U_{13} = 16 + (13 - 1)3 = 52$$

$$U_{13} = 50$$

$$U_{13} = 49$$

$$U_{13} = 53$$

$$U_{13} = 52$$

**Question 23**      Experience: 45      Order:      Level:

$X_n$  is an arithmetic sequence, given that  $X_{13} = 51$  and  $X_{19} = 33$ , find  $X_{10}$

**Solution 23**

$$X_n = a + (n - 1)d$$

$$X_{13} = a + (13 - 1)d = a + 12d = 51 \quad (1)$$

$$X_{19} = a + (19 - 1)d = a + 18d = 33 \quad (2)$$

$$(2) - (1) \quad a + 18d - (a + 12d) = 33 - 51$$

$$6d = -18$$

$$d = -3$$

$$\text{Sub into(1)} \quad a + 12(-3) = 51$$

$$a = 87$$

$$X_{10} = 63$$

$$X_{10} = 87 + (10 - 1)(-3) = 60$$

$$X_{10} = 62$$

$$X_{10} = 61$$

$$X_{10} = 59$$

$$X_{10} = 60$$

**Question 24** Experience: 45 Order: Level:

$u_n$  is an arithmetic sequence, given that  $u_3 = 5$  and  $u_7 = 13$ , for what value of  $n$  is  $a_n = 71$

**Solution 24**

$$u_n = a + (n - 1)d$$

$$u_3 = a + (3 - 1)d = a + 2d = 5 \quad (1)$$

$$u_7 = a + (7 - 1)d = a + 6d = 13 \quad (2)$$

$$(2) - (1) \quad a + 6d - (a + 2d) = 13 - 5$$

$$4d = 8$$

$$d = 2$$

$$\text{Sub into(1)} \quad a + 2(2) = 5$$

$$a = 1$$

$$u_n = 1 + (n - 1)2 = 71$$

$$n = 35$$

$$n - 1 = 35$$

$$n = 32$$

$$n = 36$$

$$n = 33$$

$$n = 34$$

$$n = 36$$

**Question 25** Experience: 30 Order: Level:

The first three terms of an arithmetic sequence are 11,14,17, find a  $n$  for which  $U_n = 83$

**Solution 25**

$$u_n = 83 \quad a = 11 \quad d = 3$$

$$u_n = a + (n - 1)d$$

$$83 = 11 + (n - 1)3$$

$$n - 1 = 24$$

$$n = 25$$

$$n = 24$$

$$n = 23$$

$$n = 22$$

$$n = 26$$

$$n = 25$$

**Question 26**      Experience: 45      Order:      Level:

$Y_n$  is an arithmetic sequence, given that  $Y_{15} = 51$  and  $X_{19} = 71$ , find  $Y_{26}$

**Solution 26**

$$Y_n = a + (n - 1)d$$

$$Y_{15} = a + (15 - 1)d = a + 14d = 51 \quad (1)$$

$$Y_{19} = a + (19 - 1)d = a + 18d = 71 \quad (2)$$

$$(2) - (1) \quad a + 18d - (a + 14d) = 71 - 51$$

$$4d = 20$$

$$d = 5$$

$$\text{Sub into (1)} \quad a + 14(5) = 51$$

$$a = -19$$

$$Y_{26} = 102$$

$$Y_{26} = 103$$

$$Y_{26} = 104$$

$$Y_{26} = 105$$

$$Y_{26} = 106$$

$$Y_{26} = -19 + (26 - 1)5 = 106$$

**Question 27**      Experience: 50      Order:      Level:

Kendrick decides to open up a savings account. He puts in £100 for the first month, £120 for the second month and an extra £20 for subsequent months till he's putting in £300 a month. Find the total amount he's saved in 2 years.

**Solution 27**

Sequence goes: 100,120,140,160,180,200...300,300,300,300...

$$U_n = a + (n - 1)d$$

$$U_n = 300 \quad a = 100 \quad d = 20$$

$$300 = 100 + (n - 1)20$$

$$n = 11$$

$$S_n = \frac{n}{2}(a + l)$$

$$n = 11 \quad a = 100 \quad l = 300$$

$$S_{11} = \frac{11}{2}(100 + 300)$$

$$S_{11} = 2200$$

Every term after is 300

$$\sum_{r=12}^{24} 300 = 13 \times 300$$

$$= 3900$$

$$\Rightarrow \text{Total days} = 2200 + 3900 = 6100$$

6105

6085

6090

6095

6100

**Question 28** Experience: 50 Order: Level:

Avery is playing with 340 sticks, she puts them in rows. The first row has 7 sticks, next row has 13 sticks, subsequent rows have 6 more sticks than the previous row. She has enough for  $k$  rows but not enough for  $k + 1$  rows. Find  $k$ .

**Solution 28**

Sequence goes: 7,13,19,25,31,37....

Not having enough for  $k+1$  rows means that  $S_k \leq 340$

$$S_n = \frac{n}{2}(2a + (n - 1)d)$$

$$S_k = \frac{k}{2}(2(7) + (k - 1)6)$$

$$S_k = k(7 + 3(k - 1))$$

$$S_k = k(3k + 4)$$

$$S_k = 3k^2 + 4k \quad (1)$$

$$S_k \leq 340$$

$$(1) \quad 3k^2 + 4k \leq 340$$

$$k = 9 \quad 3k^2 + 4k - 340 \leq 0 \quad P = -1020 \quad S = 4$$

$$k = 11 \left( k + \frac{34}{3} \right) (k - 10) \leq 0 \quad (34, -30) \quad \left( \frac{34}{3}, -10 \right)$$

© One Maths Limited  $k = 10$

$$k = 7$$

$$k = 8$$

$$k = 10$$

**Question 29** Experience: 50 Order: Level:

Griffin is training daily for a cycling marathon in 100 days. He cycles 10km on the first day, 11km on the second day and 1 more km then the previous day till he's cycling 40km a day. Calculate the total number of km he's cycled as training for the marathon.

**Solution 29**

Sequence goes: 10,11,12,13,14,15...40,40,40,40...

$$U_n = a + (n - 1)d$$

$$U_n = 40 \quad a = 10 \quad d = 1$$

$$40 = 10 + (n - 1)1$$

$$n = 31$$

$$S_n = \frac{n}{2}(a + l)$$

$$n = 31 \quad a = 10 \quad l = 40$$

$$S_{31} = \frac{31}{2}(10 + 40)$$

$$S_{31} = 775$$

Every term after is 40

$$\sum_{r=32}^{100} 40 = 69 \times 40$$

$$= 2760$$

$$\Rightarrow \text{Total days} = 775 + 2760 = 3535$$

3540

3520

3525

3530

3535

**Question 30** Experience: 50 Order: Level:

Heidi is training daily for a swimming competition in 60 days. She swims 10 laps on the first day, 12 laps on the second day and 2 more laps then the previous day till she's swimming 30 laps a day. Calculate the total number of laps she's swum as training for the competition.

**Solution 30**

Sequence goes: 10,12,14,16,18,20...30,30,30,30...



$$U_n = a + (n - 1)d$$

$$U_n = 30 \quad a = 10 \quad d = 2$$

$$30 = 10 + (n - 1)2$$

$$n = 11$$

$$S_n = \frac{n}{2}(a + l)$$

$$n = 11 \quad a = 10 \quad l = 30$$

$$S_{11} = \frac{11}{2}(10 + 30)$$

$$S_{11} = 220$$

Every term after is 30

$$\sum_{r=12}^{60} 30 = 49 \times 30$$

$$= 1470$$

$$\Rightarrow \text{Total days} = 220 + 1470 = 1690$$

1685

1705

1700

1695

1690

**Question 31**    Experience: 40    Order:    Level:

Find the sum of all numbers divisible by 3 between 2 and 200

**Solution 31**

$$200 \div 3 = 66 \text{ remainder } 2$$

$$\Rightarrow \text{last term} = 3 \times 66 = 198$$

$$S = 3 + 6 + 9 + \dots + 198$$

$$a = 3 \quad d = 3 \quad U_n = 198$$

$$U_n = a + (n - 1)d$$

$$198 = 3 + (n - 1)3$$

$$n = 66$$

$$S = \frac{n}{2}(a + l)$$

$$S = \frac{66}{2}(3 + 198)$$

$$S = 6633$$

$$S = 6642$$

$$S = 6639$$

$$S = 6636$$

$$S = 6630$$

$$S = 6633$$

**Question 32** Experience: 50 Order: Level:

James is playing with 324 sticks, she puts them in rows. The first row has 5 sticks, next row has 9 sticks, subsequent rows have 4 more sticks than the previous row. She has enough for  $k$  rows but not enough for  $k + 1$  rows. Find  $k$ .

**Solution 32**

Sequence goes: 5,9,13,17,21,25....

Not having enough for  $k+1$  rows means that  $S_k \leq 324$

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$S_k = \frac{k}{2}(2(5) + (k-1)4)$$

$$S_k = k(5 + 2k - 2)$$

$$S_k = k(2k + 3)$$

$$S_k = 2k^2 + 3k \quad (1)$$

$$S_k \leq 324$$

$$(1) \quad 2k^2 + 3k \leq 324$$

$$2k^2 + 3k - 324 \leq 0 \quad P = -648 \quad S = 3$$

$$\left(k + \frac{27}{2}\right)(k - 12) \leq 0 \quad (27, -24) \quad \left(\frac{27}{2}, -12\right)$$

$$k = 12$$

$$k = 11$$

$$k = 15$$

$$k = 14$$

$$k = 13$$

$$k = 12$$

### Lesson 3 Recurrence Relations

**Question 1** Experience: 50 Order: Level:

A sequence is defined by the recurrence relation  $X_{n+1} = \sqrt{k}X_n - 2$ ,  $X_1 = 2$ ,  $k > 0$ , given that  $X_3 = 2$  find the value of  $k$ .

**Solution 1**

$$X_2 = \sqrt{k}X_1 - 2$$

$$X_2 = 2\sqrt{k} - 2$$

$$X_3 = \sqrt{k}X_2 - 2$$

$$X_3 = \sqrt{k}(2\sqrt{k} - 2) - 2$$

$$X_3 = 2k - 2\sqrt{k} - 2 \quad \text{set } x = \sqrt{k}$$

$$X_3 = 2x^2 - 2x - 2 \quad X_3 = 2$$

$$2 = 2x^2 - 2x - 2$$

$$1 = x^2 - x - 1$$

$$0 = x^2 - x - 2 \quad S = -1 \quad P = -2$$

$$0 = (x - 2)(x + 1) \quad (-2, 1)$$

$$\sqrt{k} = 2$$

$$k = 4$$

$$k = 5$$

$$k = 3$$

$$k = 6$$

$$k = 7$$

$$k = 4$$

**Question 2** Experience: 50 Order: Level:

A sequence is defined by the recurrence relation  $U_{n+1} = aU_n + \frac{1}{b}$ ,  $U_1 = 3$ , given that  $U_2 = 7$  and  $U_3 = 15$  find the value of  $a$  and  $b$ .

**Solution 2**

$$U_2 = aU_1 + \frac{1}{b} \quad U_2 = 7$$

$$7 = 3a + \frac{1}{b} \quad (1)$$

$$U_3 = aU_2 + \frac{1}{b} \quad U_2 = 7, U_3 = 15$$

$$15 = 7a + \frac{1}{b} \quad (2)$$

$$(2) - (1) \quad 15 - 7 = 7a + \frac{1}{b} - \left(3a + \frac{1}{b}\right)$$

$$8 = 4a$$

$$a = 2$$

$$\text{Sub into (1)} \quad 7 = 3(2) + \frac{1}{b}$$

$$\frac{1}{b} = 1$$

$$b = 1$$

$$a = 2 \quad b = 3$$

$$a = 3 \quad b = 3$$

$$a = 3 \quad b = 1$$

$$a = 1 \quad b = 1$$

$$a = 2 \quad b = 1$$

**Question 3**    Experience: 70    Order:    Level:

A sequence is defined by the recurrence relation  $a_{n+1} = ka_n - 4, k > 0, a_1 = 5$ , given that  $\sum_{r=1}^3 a_r = 19$ , find the value of  $k$ .

**Solution 3**

$$a_2 = ka_1 - 4$$

$$a_2 = 5k - 4$$

$$a_3 = ka_2 - 4$$

$$a_3 = k(5k - 4) - 4$$

$$a_3 = 5k^2 - 4k - 4$$

$$\sum_{r=1}^3 a_r = a_1 + a_2 + a_3$$

$$\sum_{r=1}^3 a_r = (5) + (5k - 4) + (5k^2 - 4k - 4)$$

$$\sum_{r=1}^3 a_r = 5k^2 + k - 3 \quad \sum_{r=1}^3 a_r = 19$$

$$19 = 5k^2 + k - 3$$

$$0 = 5k^2 + k - 22 \quad S = 1 \quad P = -110$$

$$0 = \left(k + \frac{11}{5}\right)(k - 2) \quad (11, -10) \Rightarrow \left(\frac{11}{5}, -2\right)$$

$$k = 2$$

$$k = 3$$

$$k = 4$$

$$k = 1$$

$$k = 5$$

$$k = 2$$

**Question 4**    Experience: 60    Order:    Level:

A sequence is defined by the recurrence relation  $U_{n+1} = 5U_n - \frac{1}{k}, k > 0, U_1 = 2$ , given that  $\sum_{r=1}^4 U_r = 293$ , find the value of  $k$ .

**Solution 4**

$$U_2 = 5U_1 - \frac{1}{k}$$

$$U_2 = 5(2) - \frac{1}{k}$$

$$U_2 = 10 - \frac{1}{k}$$

$$U_3 = 5U_2 - \frac{1}{k}$$

$$U_3 = 5 \left( 10 - \frac{1}{k} \right) - \frac{1}{k}$$

$$U_3 = 50 - \frac{6}{k}$$

$$U_4 = 5U_3 - \frac{1}{k}$$

$$U_4 = 5 \left( 50 - \frac{6}{k} \right) - \frac{1}{k}$$

$$U_4 = 250 - \frac{31}{k}$$

$$\sum_{r=1}^4 U_r = U_1 + U_2 + U_3 + U_4$$

$$\sum_{r=1}^4 U_r = (2) + \left( 10 - \frac{1}{k} \right) + \left( 50 - \frac{6}{k} \right) + \left( 250 - \frac{31}{k} \right)$$

$$\sum_{r=1}^4 U_r = 312 - \frac{38}{k} \quad \sum_{r=1}^4 U_r = 293$$

$$312 - \frac{38}{k} = 293$$

$$19 = \frac{38}{k}$$

$$k = 2$$

$$k = 5$$

$$k = 4$$

$$k = 3$$

$$k = 1$$

$$k = 2$$

**Question 5**    Experience: 100    Order:    Level:

A sequence is defined by the recurrence relation  $X_{n+1} = \frac{k}{X_n} + 3$ ,  $X_1 = 1$ , given that  $2 \sum_{r=1}^3 X_r = 21$ , find the value of  $k$ .

**Solution 5**

$$X_2 = \frac{k}{X_1} + 3$$

$$X_2 = \frac{k}{1} + 3$$

$$X_2 = k + 3$$

$$X_3 = \frac{k}{X_2} + 3$$

$$X_3 = \frac{k}{k+3} + 3$$

$$\sum_{r=1}^3 X_r = X_1 + X_2 + X_3$$

$$\sum_{r=1}^3 X_r = (1) + (k+3) + \left( \frac{k}{k+3} + 3 \right)$$

$$\sum_{r=1}^3 X_r = k + 7 + \frac{k}{k+3} \quad 2 \sum_{r=1}^3 X_r = 21$$

$$21 = 2 \left( k + 7 + \frac{k}{k+3} \right)$$

$$21 = 2k + 14 + \frac{2k}{k+3}$$

$$7 = 2k + \frac{2k}{k+3}$$

$$7(k+3) = 2k(k+3) + 2k$$

$$7k + 21 = 2k^2 + 6k + 2k$$

$$0 = 2k^2 - k - 21 \quad S = -1 \quad P = -42$$

$$0 = \left( k + \frac{7}{2} \right) (k - 3) \quad (7, -6) \Rightarrow \left( \frac{7}{2}, -3 \right)$$

$$k = 3$$

$$k = 5$$

$$k = 2$$

$$k = 4$$

$$k = 1$$

$$k = 3$$

**Question 6**    Experience: 35    Order:    Level:

A sequence is defined by the recurrence relation  $a_{n+1} = a_n^2 - a_n$ , given that  $a_n$  is a positive sequence and that  $a_3 = 132$  find the value of  $a_1$ .

**Solution 6**

$$a_3 = a_2^2 - a_2$$

$$132 = a_2^2 - a_2$$

$$0 = a_2^2 - a_2 - 132 \quad S = 1 \quad P = -132$$

$$0 = (a_2 + 11)(a_2 - 12) \quad (11, -12)$$

$$a_2 = 12$$

$$a_2 = a_1^2 - a_1$$

$$12 = a_1^2 - a_1$$

$$0 = a_1^2 - a_1 - 12$$

$$0 = (a_1 - 4)(a_1 + 3)$$

$$a_1 = 4$$

$$a_1 = 6$$

$$a_1 = 5$$

$$a_1 = 3$$

$$a_1 = 7$$

$$a_1 = 4$$

**Question 7** Experience: 35 Order: Level:

A sequence is defined by the recurrence relation  $U_{n+1} = 5U_n - \frac{6}{U_n}$ , given that  $U_3 = 13$ ,  $U_2 > 0$ , find the value of  $U_2$ .

**Solution 7**

$$U_3 = 5U_2 - \frac{6}{U_2}$$

$$13 = 5U_2 - \frac{6}{U_2}$$

$$0 = 5U_2 - 13 - \frac{6}{U_2}$$

$$0 = 5(U_2)^2 - 13U_2 - 6 \quad S = -13 \quad P = -30$$

$$0 = \left(U_2 + \frac{2}{5}\right)(U_2 - 3) \quad (2, -15) \quad \left(\frac{2}{5}, -3\right)$$

$$U_2 = 3$$

$$U_2 = 4$$

$$U_2 = 5$$

$$U_2 = 2$$

$$U_2 = 1$$

$$U_2 = 3$$

**Question 8** Experience: 15 Order: Level:

A sequence is defined by the recurrence relation  $Y_{n+1} = 3Y_n - 5$ , given that  $Y_3 = 7$ , find the value of  $Y_1$ .

**Solution 8**

$$Y_3 = 3Y_2 - 5$$

$$7 = 3Y_2 - 5$$

$$Y_2 = 4$$

$$Y_2 = 3Y_1 - 5$$

$$4 = 3Y_1 - 5$$

$$Y_1 = 3$$

$$Y_1 = 5$$

$$Y_1 = 4$$

$$Y_1 = 1$$

$$Y_1 = 2$$

$$Y_1 = 3$$

**Question 9** Experience: 40 Order: Level:

A sequence is defined by the recurrence relation  $a_{n+1} = a_n - \frac{2a_n + 6}{a_n + 3}$ , given that  $a_2 = 5$ , find the value of  $a_1$ .

**Solution 9**

$$a_2 = a_1 - \frac{2a_1 + 6}{a_1 + 3}$$

$$5 = a_1 - \frac{2a_1 + 6}{a_1 + 3}$$

$$5(a_1 + 3) = a_1(a_1 + 3) - (2a_1 + 6)$$

$$5a_1 + 15 = (a_1)^2 + 3a_1 - 2a_1 - 6$$

$$0 = (a_1)^2 - 4a_1 - 21 \quad S = -4 \quad P = -21$$

$$0 = (a_1 + 3)(a_1 - 7) \quad (3, -7)$$

$$a_1 = 7$$

$$a_1 = 8$$

$$a_1 = 4$$

$$a_1 = 5$$

$$a_1 = 6$$

$$a_1 = 7$$

**Question 10** Experience: 25 Order: Level:

A sequence is defined by the recurrence relation  $X_{n+1} = 3(X_n)^2 - 11$ , given that  $X_1 = 2$ , find  $\sum_{r=1}^4 X_r$ .

**Solution 10**



$$X_2 = 3(X_1)^2 - 11$$

$$X_2 = 3(2)^2 - 11$$

$$X_2 = 1$$

$$X_3 = 3(X_2)^2 - 11$$

$$X_3 = 3(1)^2 - 11$$

$$X_3 = -8$$

$$X_4 = 3(X_3)^2 - 11$$

$$X_4 = 3(-8)^2 - 11$$

$$X_4 = 181$$

$$\sum_{r=1}^4 X_r = X_1 + X_2 + X_3 + X_4$$

$$\sum_{r=1}^4 X_r = (2) + (1) + (-8) + (181)$$

$$\sum_{r=1}^4 X_r = 176$$

$$\sum_{r=1}^4 X_r = 173$$

$$\sum_{r=1}^4 X_r = 170$$

$$\sum_{r=1}^4 X_r = 177$$

$$\sum_{r=1}^4 X_r = 172$$

$$\sum_{r=1}^4 X_r = 176$$

**Question 11**      Experience: 25      Order:      Level:

A sequence is defined by the recurrence relation  $U_{n+2} = 3U_{n+1} - U_n + 5$ , given that  $U_1 = 4$ ,  $U_2 = 2$ , find  $\sum_{r=1}^4 U_r$ .

**Solution 11**

$$U_3 = 3U_2 - U_1 + 5$$

$$U_3 = 3(2) - (4) + 5$$

$$U_3 = 7$$

$$U_4 = 3U_3 - U_2 + 5$$

$$U_4 = 3(7) - (2) + 5$$

$$U_4 = 24$$

$$\sum_{r=1}^4 U_r = U_1 + U_2 + U_3 + U_4$$

$$\sum_{r=1}^4 U_r = 4 + 2 + 7 + 24$$

$$\sum_{r=1}^4 U_r = 37$$

$$\sum_{r=1}^4 U_r = 36$$

$$\sum_{r=1}^4 U_r = 35$$

$$\sum_{r=1}^4 U_r = 38$$

$$\sum_{r=1}^4 U_r = 34$$

$$\sum_{r=1}^4 U_r = 37$$

**Question 12**      Experience: 25      Order:      Level:

A sequence is defined by the recurrence relation  $Y_{n+1} = 21 - 2Y_n$ , given that  $Y_1 = 5$ , find  $\sum_{r=2}^4 Y_r$ .

**Solution 12**

$$Y_2 = 21 - 2Y_1$$

$$Y_2 = 21 - 2(5)$$

$$Y_2 = 11$$

$$Y_3 = 21 - 2Y_2$$

$$Y_3 = 21 - 2(11)$$

$$Y_3 = -1$$

$$Y_4 = 21 - 2Y_3$$

$$Y_4 = 21 - 2(-1)$$

$$Y_4 = 23$$

$$\sum_{r=2}^4 Y_r = Y_2 + Y_3 + Y_4$$

$$\sum_{r=2}^4 Y_r = 11 + (-1) + 23$$

$$\sum_{r=2}^4 Y_r = 33$$

$$\sum_{r=2}^4 Y_r = 32$$

$$\sum_{r=2}^4 Y_r = 31$$

$$\sum_{r=2}^4 Y_r = 30$$

$$\sum_{r=2}^4 Y_r = 34$$

$$\sum_{r=2}^4 Y_r = 33$$

**Question 13**      Experience: 30      Order:      Level:

A sequence is defined by the recurrence relation  $X_{n+1} = 5 - X_n$ , given that  $X_1 = 7$ , find  $\sum_{r=1}^{20} X_r$ .

**Solution 13**

$$X_2 = 5 - X_1 = 5 - 7 = -2$$

$$X_3 = 5 - X_2 = 5 - (-2) = 7$$

$$X_4 = 5 - X_3 = 5 - 7 = -2$$

$$X_5 = 5 - X_4 = 5 - (-2) = 7$$

$$\sum_{r=1}^{20} X_r = X_1 + X_2 + X_3 + X_4 + \dots + X_{20}$$

$$\sum_{r=1}^{20} X_r = -2 + 7 + -2 + 7 + -2 + \dots + 7$$

$$\sum_{r=1}^{20} X_r = 10(-2) + 10(7)$$

$$\sum_{r=1}^{20} X_r = 50$$

$$\sum_{r=1}^{20} X_r = 20$$

$$\sum_{r=1}^{20} X_r = 60$$

$$\sum_{r=1}^{20} X_r = 30$$

$$\sum_{r=1}^{20} X_r = 40$$

$$\sum_{r=1}^{20} X_r = 50$$

**Question 14** Experience: 30 Order: Level:

A sequence is defined by the recurrence relation  $Y_{n+1} = 5 + 5Y_n - 2(Y_n)^3$ , given that  $Y_1 = 2$ , find  $Y_{1000}$ .

**Solution 14**

$$Y_2 = 5 + 5Y_1 - 2(Y_1)^3 = 5 + 5(2) - 2(2)^3 = -1$$

$$Y_3 = 5 + 5Y_2 - 2(Y_2)^3 = 5 + 5(-1) - 2(-1)^3 = 2$$

$$Y_4 = 5 + 5Y_3 - 2(Y_3)^3 = 5 + 5(2) - 2(2)^3 = -1$$

|       |       |       |       |       |       |
|-------|-------|-------|-------|-------|-------|
| $Y_1$ | $Y_2$ | $Y_3$ | $Y_4$ | $Y_5$ | $Y_6$ |
| -1    | 2     | -1    | 2     | -1    | 2     |

We can see that  $Y_2 = Y_4 = Y_6 = Y_8 = \dots = 2$

Every numbered term divisible by 2 is 2

Find a numbered term that is close to  $Y_{1000}$  that is divisible by 2

$$Y_2 = 2 \quad Y_4 = 2 \quad Y_{100} = 2 \quad Y_{1000} = 2$$

$$Y_{1000} = 1$$

$$Y_{1000} = 0$$

$$Y_{1000} = 3$$

$$Y_{1000} = 4$$

$$Y_{1000} = 2$$

**Question 15**      Experience: 15      Order:      Level:

Given  $\sum_{r=1}^n x_r = 5n^2 - 3$ , find  $\sum_{r=1}^7 x_r$

**Solution 15**

$$\sum_{r=1}^7 x_r = 5(7)^2 - 3 = 242$$

$$\sum_{r=1}^7 x_r = 239$$

$$\sum_{r=1}^7 x_r = 240$$

$$\sum_{r=1}^7 x_r = 243$$

$$\sum_{r=1}^7 x_r = 241$$

$$\sum_{r=1}^7 x_r = 242$$

**Question 16**      Experience: 40      Order:      Level:

A sequence is defined by the recurrence relation  $U_{n+1} = \frac{13 - 5U_n}{7 - 3U_n}$ , given that  $U_1 = 1$ , find  $U_{50}$ .

**Solution 16**

$$U_2 = \frac{13 - 5U_1}{7 - 3U_1} = \frac{13 - 5(1)}{7 - 3(1)} = \frac{8}{4} = 2$$

$$U_3 = \frac{13 - 5U_2}{7 - 3U_2} = \frac{13 - 5(2)}{7 - 3(2)} = \frac{3}{1} = 3$$

$$U_4 = \frac{13 - 5U_3}{7 - 3U_3} = \frac{13 - 5(3)}{7 - 3(3)} = \frac{-2}{-2} = 1$$

$$U_5 = \frac{13 - 5U_4}{7 - 3U_4} = \frac{13 - 5(1)}{7 - 3(1)} = \frac{8}{4} = 2$$

|       |       |       |       |       |       |
|-------|-------|-------|-------|-------|-------|
| $U_1$ | $U_2$ | $U_3$ | $U_4$ | $U_5$ | $U_6$ |
| 1     | 2     | 3     | 1     | 2     | 3     |

We can see that  $U_3 = U_6 = U_9 = U_{12} = \dots = 3$

Every numbered term divisible by 3 is 3

Find a numbered term that is close to  $U_{50}$  that is divisible by 3

$$U_3 = 3 \quad U_9 = 3 \quad U_{30} = 3 \quad U_{51} = 3$$

$U_{51} = 3 \Rightarrow U_{50} = 2$  since 2 is the term before 3 in the sequence

i.e. 1, 2, 3, 1, 2, 3, 1,

$$U_{50} = 1$$

$$U_{50} = 3$$

$$U_{50} = 4$$

$$U_{50} = 5$$

$$U_{50} = 2$$

**Question 17**      Experience: 30      Order:      Level:

Given  $\sum_{r=1}^n a_r = 2n^3 + 5$ , find  $a_2$

**Solution 17**

$$\sum_{r=1}^n a_r = 2n^3 + 5$$

$$a_2 = \sum_{r=1}^2 a_r - \sum_{r=1}^1 a_r$$

$$a_2 = 2(2)^3 + 5 - (2(1)^3 + 5)$$

$$a_2 = 21 - 7$$

$$a_2 = 14$$

$$a_2 = 13$$

$$a_2 = 12$$

$$a_2 = 11$$

$$a_2 = 15$$

$$a_2 = 14$$

**Question 18**      Experience: 15      Order:      Level:

Given  $\sum_{r=1}^n U_r = 6n^2 + 11$ , find  $U_1$

**Solution 18**

$$\sum_{r=1}^1 U_r = U_1 = 6(1)^2 + 11 = 17$$

$$U_1 = 16$$

$$U_1 = 15$$

$$U_1 = 14$$

$$U_1 = 18$$

$$U_1 = 17$$

**Question 19**      Experience: 15      Order:      Level:

Given  $\sum_{r=1}^n u_r = n^3 + 4$ , find  $\sum_{r=1}^5 u_r$

**Solution 19**

$$\sum_{r=1}^5 u_r = (5)^3 + 4 = 129$$

$$\sum_{r=1}^5 u_r = 130$$

$$\sum_{r=1}^5 u_r = 126$$

$$\sum_{r=1}^5 u_r = 127$$

$$\sum_{r=1}^5 u_r = 128$$

$$\sum_{r=1}^5 u_r = 129$$

**Question 20**      Experience: 30      Order:      Level:

Given  $\sum_{r=1}^n Y_r = 3n^3 - 2$ , find  $Y_3$

**Solution 20**

$$Y_3 = \sum_{r=1}^3 - \sum_{r=1}^2$$

$$Y_3 = 3(3)^3 - 2 - (3(2)^3 - 2)$$

$$Y_3 = 57$$

$$Y_3 = 54$$

$$Y_3 = 55$$

$$Y_3 = 52$$

$$Y_3 = 56$$

$$Y_3 = 57$$

**Question 21**      Experience: 30      Order:      Level:

Given  $\sum_{r=1}^n U_r = 3n + 7$ , find  $U_5$

**Solution 21**

$$U_5 = \sum_{r=1}^5 U_r - \sum_{r=1}^4 U_r$$

$$U_5 = 3(5) + 7 - (3(4) + 7)$$

$$U_5 = 3$$

$$U_5 = 2$$

$$U_5 = 1$$

$$U_5 = 4$$

$$U_5 = 5$$

$$U_5 = 3$$

**Question 22** Experience: 45 Order: Level:

A sequence is defined by the recurrence relation  $U_{n+1} = kU_n - 4$ ,  $U_1 = 3$ ,  $k > 0$ , given that  $U_3 = 0$  find the value of  $k$

**Solution 22**

$$U_2 = kU_1 - 4$$

$$U_2 = 3k - 4$$

$$U_3 = kU_2 - 4$$

$$U_3 = k(3k - 4) - 4$$

$$U_3 = 3k^2 - 4k - 4 \quad U_3 = 0$$

$$0 = 3k^2 - 4k - 4 \quad S = -4 \quad P = -12$$

$$0 = \left(k + \frac{2}{3}\right)(k - 2) \quad (2, -6) \Rightarrow \left(\frac{2}{3}, -2\right)$$

$$k = 2$$

$$k = 3$$

$$k = 4$$

$$k = 1$$

$$k = 5$$

$$k = 2$$

**Question 23** Experience: 60 Order: Level:

A sequence is defined by the recurrence relation  $a_{n+1} = \frac{a_n}{k} + 3$ ,  $a_1 = 3$ ,  $k > 0$ , given that  $a_3 = 9$  find the value of  $k$

**Solution 23**



$$a_2 = \frac{a_1}{k} + 3$$

$$a_2 = \frac{3}{k} + 3$$

$$a_3 = \frac{a_2}{k} + 3$$

$$a_3 = \frac{\left(\frac{3}{k} + 3\right)}{k} + 3$$

$$a_3 = \frac{3}{k^2} + \frac{3}{k} + 3 \quad a_3 = 9$$

$$9 = \frac{3}{k^2} + \frac{3}{k} + 3$$

$$6 - \frac{3}{k^2} - \frac{3}{k} = 0$$

$$6k^2 - 3k - 3 = 0$$

$$2k^2 - k - 1 = 0 \quad S = -1 \quad P = -2$$

$$\left(x + \frac{1}{2}\right)(k - 1) = 0 \quad (1, -2) \Rightarrow \left(\frac{1}{2}, -1\right)$$

$$k = 1$$

$$k = 2$$

$$k = 3$$

$$k = 4$$

$$k = 5$$

$$k = 1$$

**Question 24**      Experience: 60      Order:      Level:

A sequence is defined by the recurrence relation  $u_{n+1} = \sqrt{a} \left(u_n - \frac{1}{b}\right)$ ,  $5u_1 = 4$ , given that  $u_2 = 7$  and  $u_3 = 13$  find the value of  $a$  and  $b$ .

**Solution 24**

$$u_2 = \sqrt{a} \left( u_1 - \frac{1}{b} \right)$$

$$7 = \sqrt{a} \left( 4 - \frac{1}{b} \right) \quad (1)$$

$$7 = 4\sqrt{a} - \frac{\sqrt{a}}{b} \quad (2)$$

$$u_3 = \sqrt{a} \left( u_2 - \frac{1}{b} \right)$$

$$13 = \sqrt{a} \left( 7 - \frac{1}{b} \right)$$

$$13 = 7\sqrt{a} - \frac{\sqrt{a}}{b} \quad (3)$$

$$(3) - (2) \quad 13 - 7 = 7\sqrt{a} - \frac{\sqrt{a}}{b} - \left( 4\sqrt{a} - \frac{\sqrt{a}}{b} \right)$$

$$6 = 3\sqrt{a}$$

$$2 = \sqrt{a}$$

$$a = 4$$

$$\text{Sub into (1)} \quad 7 = \sqrt{4} \left( 4 - \frac{1}{b} \right)$$

$$\frac{7}{2} = 4 - \frac{1}{b}$$

$$-\frac{1}{2} = -\frac{1}{b}$$

$$b = 2$$

$$a = 3 \quad b = 2$$

$$a = 4 \quad b = 3$$

$$a = 3 \quad b = 3$$

$$a = 2 \quad b = 3$$

$$a = 4 \quad b = 2$$

**Question 25**      Experience: 30      Order:      Level:

A sequence is defined by the recurrence relation  $a_{n+1} = 3 - a_n$ , given that  $a_1 = 1$ , find  $\sum_{r=1}^{100} a_r$ .

**Solution 25**

$$a_2 = 3 - a_1 = 3 - 1 = 2$$

$$a_3 = 3 - a_2 = 3 - 2 = 1$$

$$a_4 = 3 - a_3 = 3 - 1 = 2$$

$$a_5 = 3 - a_4 = 3 - 2 = 1$$

$$\sum_{r=1}^{100} a_r = a_1 + a_2 + a_3 + a_4 + \dots + a_{100}$$

$$\sum_{r=1}^{100} a_r = 1 + 2 + 1 + 2 + 1 + 2 + \dots + 2$$

$$\sum_{r=1}^{100} a_r = 50(2) + 50(1)$$

$$\sum_{r=1}^{100} a_r = 150$$

$$\sum_{r=1}^{100} a_r = 100$$

$$\sum_{r=1}^{100} a_r = 200$$

$$\sum_{r=1}^{100} a_r = 50$$

$$\sum_{r=1}^{100} a_r = 250$$

$$\sum_{r=1}^{100} a_r = 150$$

**Question 26** Experience: 40 Order: Level:

A sequence is defined by the recurrence relation  $A_{n+1} = \frac{4A_n - 16}{3A_n - 8}$ , given that  $A_1 = 0$ , find  $A_{100}$ .

**Solution 26**

$$A_2 = \frac{4A_1 - 16}{3A_1 - 8} = \frac{4(0) - 16}{3(0) - 8} = \frac{-16}{-8} = 2$$

$$A_3 = \frac{4A_2 - 16}{3A_2 - 8} = \frac{4(2) - 16}{3(2) - 8} = \frac{-8}{-2} = 4$$

$$A_4 = \frac{4A_1 - 16}{3A_1 - 8} = \frac{4(4) - 16}{3(4) - 8} = 0$$

$$A_5 = \frac{4A_1 - 16}{3A_1 - 8} = \frac{4(0) - 16}{3(0) - 8} = \frac{-16}{-8} = 2$$

|       |       |       |       |       |       |
|-------|-------|-------|-------|-------|-------|
| $a_1$ | $a_2$ | $a_3$ | $a_4$ | $a_5$ | $a_6$ |
| 0     | 2     | 4     | 0     | 2     | 4     |

We can see that  $a_3 = a_6 = a_9 = a_{12} = \dots = 4$

Every numbered term divisible by 3 is 4

Find a numbered term that is close to  $a_{100}$  that is divisible by 3

$$a_3 = 4 \quad a_9 = 4 \quad a_{30} = 4 \quad a_{99} = 4$$

$a_{99} = 4 \Rightarrow a_{100} = 0$  since 0 is the next term after 4 in the sequence

i.e. 0, 2, 4, 0, 2, 4, 0

$$a_{100} = 1$$

$$a_{100} = 2$$

$$a_{100} = 3$$

$$a_{100} = 4$$

$$a_{100} = 0$$

#### Lesson 4 Arithmetic Sequence 2

**Question 1**    Experience: 50    Order:    Level:

$U_n$  is an arithmetic sequence with  $S_n$  being the sum of the first  $n$  terms of the sequence. Given that  $U_4 = 11$  and  $U_7 = 23$ , find  $S_{11}$

**Solution 1**

$$U_n = a + (n - 1)d$$

$$U_4 = a + (4 - 1)d = a + 3d = 11 \quad (1)$$

$$U_7 = a + (7 - 1)d = a + 6d = 23 \quad (2)$$

$$(2) - (1) \quad a + 6d - (a + 3d) = 23 - 11$$

$$3d = 12$$

$$d = 4$$

$$\text{Sub into (1)} \quad a + 3(4) = 11$$

$$a = -1$$

$$S_n = \frac{n}{2}(2(a) + (n - 1)d)$$

$$S_{11} = \frac{11}{2}(2(-1) + (11 - 1)4) = 209$$

$$S_{11} = 208$$

$$S_{11} = 205$$

$$S_{11} = 206$$

$$S_{11} = 207$$

$$S_{11} = 209$$

**Question 2**    Experience: 50    Order:    Level:

$U_n$  is an arithmetic sequence with  $S_n$  being the sum of the first  $n$  terms of the sequence. Given that  $U_3 = -5$  and  $U_5 = -11$ , find  $S_7$

**Solution 2**

$$U_n = a + (n - 1)d$$

$$U_3 = a + (3 - 1)d = a + 2d = -5 \quad (1)$$

$$U_5 = a + (5 - 1)d = a + 4d = -11 \quad (2)$$

$$(2) - (1) \quad a + 4d - (a + 2d) = -11 - (-5)$$

$$2d = -6$$

$$d = -3$$

$$\text{Sub into (1)} \quad a + 2(-3) = -5$$

$$a = 1$$

$$U_7 = 1 + (7 - 1)(-3) = -17$$

$$S_7 = \frac{7}{2}(1 + (-17)) = -56$$

$$S_7 = -52$$

$$S_7 = -53$$

$$S_7 = -54$$

$$S_7 = -55$$

$$S_7 = -56$$

**Question 3** Experience: 45 Order: Level:

The first three terms of an arithmetic sequence are 60, 58, 56..., there exists a  $k^{\text{th}}$  term which = 0, find the value of  $k$ , hence of otherwise find the maximum value of  $S_n$

**Solution 3**

$$U_n = a + (n - 1)d$$

$$U_k = 60 + (k - 1)(-2) = 0$$

$$k - 1 = 30$$

$$k = 31$$

maximum value of  $S_n = S_k$  as any term after  $U_k$  is negative

$$S_n = \frac{n}{2}(a + d)$$

$$k = 28 \quad S_k = 935$$

$$k = 29 \quad S_k = 915$$

$$k = 30 \quad S_k = 920$$

$$k = 32 \quad S_k = 925$$

$$k = 31 \quad S_k = 930$$

$$S_k = \frac{31}{2}(60 + 0)$$

$$S_k = 930$$

**Question 4** Experience: 50 Order: Level:

$U_n$  is an arithmetic sequence with  $S_n$  being the sum of the first  $n$  terms of the sequence. Given that  $S_5 = 85$  and  $S_8 = 184$ , find  $U_6$

**Solution 4**

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$S_5 = \frac{5}{2}(2a + (5-1)d) = 85$$

$$2a + 4d = 34 \quad (1)$$

$$S_8 = \frac{8}{2}(2a + (8-1)d) = 184$$

$$2a + 7d = 46 \quad (2)$$

$$(2) - (1) \quad 2a + 7d - (2a + 4d) = 46 - 34$$

$$3d = 12$$

$$d = 4$$

$$\text{Sub into (1)} \quad 2a + 4(4) = 34$$

$$a = 9$$

$$U_6 = a + (6-1)d = 9 + 5(4) = 29$$

$$U_6 = 31$$

$$U_6 = 30$$

$$U_6 = 27$$

$$U_6 = 28$$

$$U_6 = 29$$

**Question 5** Experience: 50 Order: Level:

$U_n$  is an arithmetic sequence with  $S_n$  being the sum of the first  $n$  terms of the sequence. Given that  $U_5 = 19$  and  $S_{10} = 170$ , find  $U_4$

**Solution 5**

$$U_n = a + (n-1)d$$

$$U_5 = a + 4d = 19 \quad (1)$$

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$S_{10} = \frac{10}{2}(2a + (10-1)d) = 170$$

$$2a + 9d = 34 \quad (2)$$

$$(2) - 2(1) \quad 2a + 9d - 2(a + 4d) = 34 - 38$$

$$d = -4$$

$$d = -4$$

$$\text{Sub into (1)} \quad a + 4(-4) = 19$$

$$a = 35$$

$$U_4 = a + (4-1)d = 35 + (3)(-4) = 23$$

$$U_4 = 19$$

$$U_4 = 20$$

$$U_4 = 21$$

$$U_4 = 22$$

$$U_4 = 23$$

**Question 6** Experience: 50 Order: Level:

$U_n$  is an arithmetic sequence with  $S_n$  being the sum of the first  $n$  terms of the sequence. Given that  $U_4 = 8$  and  $S_{12} = 0$ , find  $S_9$

**Solution 6**

$$U_n = a + (n - 1)d$$

$$U_4 = a + 3d = 8 \quad (1)$$

$$S_n = \frac{n}{2}(2a + (n - 1)d)$$

$$S_{12} = \frac{12}{2}(2a + (12 - 1)d) = 0$$

$$2a + 11d = 0 \quad (2)$$

$$(2) - 2(1) \quad 2a + 11d - 2(a + 3d) = 0 - 16$$

$$8d = -16$$

$$d = -2$$

$$\text{Sub into (1)} \quad a + 3(-2) = 8$$

$$a = 14$$

$$S_9 = \frac{9}{2}(2(14) + (9 - 1)(-2)) = 54$$

$$S_9 = 55$$

$$S_9 = 51$$

$$S_9 = 52$$

$$S_9 = 53$$

$$S_9 = 54$$

**Question 7** Experience: 50 Order: Level:

$U_n$  is an arithmetic sequence with  $S_n$  being the sum of the first  $n$  terms of the sequence. Given that  $U_3 = 4$  and  $U_7 = 0$ , find  $S_{10}$

**Solution 7**

$$U_n = a + (n - 1)d$$

$$U_3 = a + 2d = 4 \quad (1)$$

$$U_7 = a + 6d = 0 \quad (2)$$

$$(2) - (1) \quad a + 6d - (a + 2d) = 0 - 4$$

$$4d = -4$$

$$d = -1$$

$$\text{Sub into (1)} \quad a + 2(-1) = 4$$

$$a = 6$$

$$S_n = \frac{n}{2}(2a + (n - 1)d)$$

$$S_{10} = \frac{10}{2}(2(6) + (10 - 1)(-1)) = 15$$

$$S_{10} = 14$$

$$S_{10} = 13$$

$$S_{10} = 12$$

$$S_{10} = 16$$

$$S_{10} = 15$$

**Question 8**    Experience: 50    Order:    Level:

$U_n$  is an arithmetic sequence with  $S_n$  being the sum of the first  $n$  terms of the sequence. Given that  $U_4 = 10$  and  $S_6 = 57$ , find  $S_{11}$

**Solution 8**

$$U_n = a + (n - 1)d$$

$$U_4 = a + 3d = 10 \quad (1)$$

$$S_n = \frac{n}{2}(2a + (n - 1)d)$$

$$S_6 = \frac{6}{2}(2a + (5 - 1)d) = 57$$

$$a + 2d = \frac{19}{2} \quad (2)$$

$$(1) - (2) \quad a + 3d - (a + 2d) = 10 - \frac{19}{2}$$

$$d = \frac{1}{2}$$

$$\text{Sub into (1)} \quad a + 3\left(\frac{1}{2}\right) = 10$$

$$a = \frac{17}{2}$$

$$S_{11} = \frac{11}{2} \left( 2\left(\frac{17}{2}\right) + (11 - 1)\left(\frac{1}{2}\right) \right) = 121$$

$$S_{10} = 14$$

$$S_{10} = 13$$

$$S_{10} = 12$$



$$S_{10} = 16$$

$$S_{10} = 15$$

**Question 9** Experience: 30 Order: Level:

Three consecutive terms in an arithmetic sequence are  $3k + 2, 2k + 5, 4k + 5$ , find the value of  $k$

**Solution 9**

$$2k + 5 - (3k + 2) = d = 4k + 5 - (2k + 5)$$

$$-k + 3 = 2k$$

$$k = 5$$

$$3k = 3$$

$$k = 4$$

$$k = 3$$

$$k = 1$$

$$k = 2$$

$$k = 1$$

**Question 10** Experience: 30 Order: Level:

Three consecutive terms in an arithmetic sequence are  $k^2 + 3, -k, k - 1$ , find the possible values of  $k$

**Solution 10**

$$-k - (k^2 + 3) = d = k - 1 - (-k)$$

$$-k - k^2 - 3 = 2k - 1$$

$$k = -2, -3$$

$$0 = k^2 + 3k + 2$$

$$k = -1, -2$$

$$0 = (k + 2)(k + 1)$$

$$k = -3, -1$$

$$k = -1, -2$$

$$k = -2, -1$$

**Question 11** Experience: 30 Order: Level:

Three consecutive terms in an arithmetic sequence are  $k + 16, 3k + 12, 7k - 2$ , find the value of  $k$

**Solution 11**

$$3k + 12 - (k + 16) = d = 7k - 2 - (3k + 12)$$

$$2k - 4 = 4k - 14$$

$$k = 6$$

$$2k = 10$$

$$k = 2$$

$$k = 3$$

$$k = 5$$

$$k = 4$$

$$k = 5$$

**Question 12** Experience: 45 Order: Level:

The first three terms in an arithmetic sequence are  $2k, k + 9, 3k$ , find the smallest  $n$  such that  $S_n > 117$

**Solution 12**

$$k + 9 - 2k = d = 3k - (k + 9)$$

$$-k + 9 = 2k - 9$$

$$3k = 18$$

$$k = 6$$

$$\Rightarrow U_1 = 12 \quad U_2 = 15 \quad U_3 = 18$$

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$\frac{n}{2}(2(12) + (n-1)3) > 117$$

$$n(24 + 3n - 3) > 234$$

$$3n^2 + 21n - 234 > 0$$

$$n^2 + 7n - 78 > 0 \quad P = -78 \quad S = 7$$

$$(n + 13)(n - 6) > 0 \quad (13, -6)$$

$$n = 6$$

$$n = 7$$

$$n = 3$$

$$n = 4$$

$$n = 5$$

$$n = 6$$

**Question 13**    Experience: 40    Order:    Level:

The first three terms of an arithmetic sequence are 99, 96, 93..., there exists a  $k^{\text{th}}$  term which = 0, find the value of  $k$ , hence of otherwise find the maximum value of  $S_n$

**Solution 13**

$$U_n = a + (n-1)d$$

$$U_k = 99 + (k-1)(-3) = 0$$

$$k - 1 = 33$$

$$k = 34$$

maximum value of  $S_n = S_k$  as any term after  $U_k$  is negative

$$S_n = \frac{n}{2}(a + l)$$

$$S_k = \frac{34}{2}(99 + 0)$$

$$S_k = 1683$$

$$k = 33 \quad S_k = 1689$$

$$k = 32 \quad S_k = 1686$$

$$k = 35 \quad S_k = 1677$$

$$k = 36 \quad S_k = 1680$$

$$k = 34 \quad S_k = 1683$$

**Question 14** Experience: 35 Order: Level:

The first three terms in an arithmetic sequence are 5, 7, 9, find the smallest  $n$  such that  $S_n > 252$

**Solution 14**

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$S_n = \frac{n}{2}(2(5) + (n-1)2) > 252$$

$$n(5 + n - 1) > 252$$

$$n^2 + 4n - 252 > 0 \quad P = -252 \quad S = 4$$

$$(n + 18)(n - 14) > 0 \quad (18, -14)$$

$$n = 14$$

$$n = 15$$

$$n = 11$$

$$n = 12$$

$$n = 13$$

$$n = 14$$

**Question 15** Experience: 35 Order: Level:

The first three terms in an arithmetic sequence are 9, 12, 15, find the smallest  $n$  such that  $S_n > 750$

**Solution 15**

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$S_n = \frac{n}{2}(2(9) + (n-1)3) > 750$$

$$n(18 + 3n - 3) > 1500$$

$$3n(5 + n) > 1500$$

$$n(5 + n) > 500$$

$$n^2 + 5n - 500 > 0 \quad P = -500 \quad S = 5$$

$$(n + 25)(n - 20) > 0 \quad (25, -20)$$

$$n = 20$$

$$n = 21$$

$$n = 17$$

$$n = 18$$

$$n = 19$$

$$n = 20$$

**Question 16** Experience: 35 Order: Level:

The first three terms in an arithmetic sequence are 12, 16, 20, 24, find the smallest  $n$  such that  $S_n > 672$

**Solution 16**

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$S_n = \frac{n}{2}(2(12) + (n-1)4) > 672$$

$$n(12 + 2n - 2) > 672$$

$$2n^2 + 10n - 672 > 0$$

$$n^2 + 5n - 336 > 0 \quad P = -336 \quad S = 5$$

$$(n + 21)(n - 16) > 0 \quad (21, -16)$$

$$n = 16$$

$$n = 15$$

$$n = 19$$

$$n = 18$$

$$n = 17$$

$$n = 16$$

**Question 17** Experience: 50 Order: Level:

Judith is playing with 294 sticks, she puts them in rows. The first row has 8 sticks, next row has 10 sticks, subsequent rows have 2 more sticks than the previous row. She has enough for  $k$  rows but not enough for  $k + 1$  rows. Find  $k$ .

**Solution 17**

Sequence goes: 8,10,12,14,18,20....

Not having enough for  $k+1$  rows means that  $S_k \leq 294$

$$S_n = \frac{n}{2}(2a + (n-1)d)$$

$$S_k = \frac{k}{2}(2(8) + (k-1)2)$$

$$S_k = k(8 + k - 1)$$

$$S_k = k(k + 7)$$

$$S_k = k^2 + 7k \quad (1)$$

$$S_k \leq 294$$

$$(1) \quad k^2 + 7k \leq 294$$

$$k^2 + 7k - 294 \leq 0 \quad P = 294 \quad S = 7$$

$$k = 11$$

$$k = 15 \quad (k + 21)(k - 14) \leq 0 \quad (21, -14)$$

$$k = 12 \quad k = 14$$

$$k = 13$$

$$k = 14$$

**Question 18** Experience: 50 Order: Level:

$U_n$  is an arithmetic sequence with  $S_n$  being the sum of the first  $n$  terms of the sequence. Given that  $S_{11} = 0$  and  $U_2 = 8$ , find  $U_6$

**Solution 18**

$$U_n = a + (n - 1)d$$

$$U_2 = a + (2 - 1)d = a + d = 8 \quad (1)$$

$$S_n = \frac{n}{2}(2a + (n - 1)d)$$

$$S_{11} = \frac{11}{2}(2a + (11 - 1)d) = 0$$

$$S_{11} = a + 5d = 0 \quad (2)$$

$$(2) - (1) \quad a + 5d - (a + d) = 0 - 8$$

$$4d = -8$$

$$d = -2$$

$$\text{Sub into (1)} \quad a + (-2) = 8$$

$$a = 10$$

$$U_6 = 1 + (7 - 1)(-2) = -11$$

$$U_6 = -13$$

$$U_6 = -12$$

$$U_6 = -9$$

$$U_6 = -10$$

$$U_6 = -11$$

**Question 19** Experience: 45 Order: Level:

The first three terms of an arithmetic sequence are 44, 41, 38..., there exists a  $k^{\text{th}}$  term which is the smallest positive term in the sequence, find the value of  $k$ , hence of otherwise find the maximum value of  $S_n$

**Solution 19**

$$U_n = a + (n - 1)d$$

$$U_k = 44 + (k - 1)(-3) = 0$$

$$k - 1 = \frac{44}{3}$$

$$k = \frac{44}{3} + 1 = 15.6$$

$$k = 15$$

maximum value of  $S_n = S_k$  as any term after  $U_k$  is negative

$$k = 14 \quad S_k = 2658 \quad S_n = \frac{n}{2}(2a + (n - 1)d)$$

$$k = 13 \quad S_k = 2656$$

$$k = 12 \quad S_k = 2654 \quad S_k = \frac{34}{2}(2(99) + (15 - 1)(-3))$$

$$k = 16 \quad S_k = 2650$$

$$k = 15 \quad S_k = 2652 \quad S_k = 2652$$

**Question 20** Experience: 50 Order: Level:

At the start of the year 2000, Tony the farmer has  $50m^2$  of land, he buys  $7m^2$  of land at the end of each year. At the beginning of this year, Tony owns  $141m^2$  of land. What year is it?

### Solution 20

Sequence goes from the start of every year: 50,57,64,71,78,85....

$$U_n = a + (n - 1)d$$

$$U_n = 141 \quad a = 50 \quad d = 7$$

$$141 = 50 + (n - 1)7$$

$$n - 1 = 13$$

$$\text{Year} = 2015$$

$$\text{Year} = 2011 \quad n = 14$$

$$\text{Year} = 2012 = 2000 + 14 = 2014$$

$$\text{Year} = 2013$$

$$\text{Year} = 2014$$

## Unit 2 Core 2

### Chapter 1 Logarithms

#### Lesson 1 Basic logarithms

**Question 1**    Experience: 10    Order: a2    Level: a2

Express  $\log_{x+5} 10 = 4$  in power form

**Solution 1**

$$\log_{x+5} 10 = 4$$

$$(x + 5)^4 = 10$$

$$4^{x+5} = 10$$

$$(x + 5)^{10} = 4$$

$$10^{x+5} = 4$$

$$(x + 5)^{10} = 4$$

$$(x + 5)^4 = 10$$

**Question 2**    Experience: 10    Order: a2    Level: a2

Express  $\log_{a+b} 6 = c$  in power form

**Solution 2**

$$\log_{a+b} 6 = c$$

$$(a + b)^c = 6$$

$$(a + b)^6 = c$$

$$6^c = a + b$$

$$(a + b)^c = 6$$

$$6^{a+b} = 6$$

$$(a + b)^c = 6$$

**Question 3**    Experience: 10    Order: a2    Level: a2

Express  $\log_{xy} 3 = 2$  in power form

**Solution 3**

$$\log_{xy} 3 = 2$$

$$(xy)^2 = 3$$

$$2^{xy} = 3$$

$$3^2 = xy$$

$$xy^3 = 2$$

$$(3)^{xy} = 2$$

$$(xy)^2 = 3$$

**Question 4**    Experience: 10    Order: b1    Level: b1

Express  $a^b = c$  in log form

**Solution 4**

$$a^b = c$$

$$\log_a c = b$$

$$\log_c a = b$$

$$\log_b c = a$$

$$\log_b a = c$$

$$\log_a b = c$$

$$\log_a c = b$$

**Question 5**    Experience: 10    Order: b1    Level: b1

Express  $5^2 = 25$  in log form

**Solution 5**

$$5^2 = 25$$

$$\log_5 25 = 2$$

$$\log_5 2 = 25$$

$$\log_{25} 2 = 5$$

$$\log_{25} 5 = 2$$

$$\log_2 25 = 5$$

$$\log_5 25 = 2$$

**Question 6**    Experience: 10    Order: b2    Level: b2

Express  $(xy)^5 = 20$  in log form

**Solution 6**

$$(xy)^5 = 20$$

$$\log_{xy} 20 = 5$$

$$\log_5 20 = xy$$

$$\log_{xy} 5 = 20$$

$$\log_{20} 5 = xy$$

$$\log_{20} xy = 5$$

$$\log_{xy} 20 = 5$$

**Question 7** Experience: 15 Order: c2 Level: c2

Express  $\log_2(x^2y) - \log_2 x$  as a single logarithm

**Solution 7**

$$\log_2(x^2y) - \log_2 x$$

$$= \log_2((x^2y) \div x)$$

$$= \log_2 xy$$

$$2 \log_{x^2y} 1$$

$$\log_{x^2y} 2$$

$$\log_2 x^2y$$

$$2x \log_2 y$$

$$\log_2 xy$$

**Question 8** Experience: 10 Order: b1 Level: b1

Express  $a^{bc} = 6$  in log form

**Solution 8**

$$a^{bc} = 6$$

$$\log_a 6 = bc$$

$$\log_6 ab = c$$

$$\log_{bc} a = 6$$

$$\log_{bc} 6 = a$$

$$\log_a bc = 6$$

$$\log_a 6 = bc$$

**Question 9** Experience: 10 Order: b2 Level: b2

Express  $(a + b)^4 = 15$  in log form

**Solution 9**

$$(a + b)^4 = 15$$

$$\log_{(a+b)} 15 = 4$$

$$\log_4 15 = a + b$$

$$\log_{15}(a + b) = 4$$

$$\log_{15} 4 = a + b$$

$$\log_4(a + b) = 15$$

$$\log_{(a+b)} 15 = 4$$

**Question 10** Experience: 10 Order: b2 Level: b2

Express  $(x + 4)^4 = 5$  in log form

**Solution 10**



$$(x + 4)^4 = 5$$

$$\log_{(x+4)} 5 = 4$$

$$\log_4(x + 4) = 5$$

$$\log_5 4 = x + 4$$

$$\log_{(x+4)} 4 = 5$$

$$\log_5(x + 4) = 5$$

$$\log_{(x+4)} 5 = 4$$

**Question 11** Experience: 15 Order: c1 Level: c1

Express  $\log_4(x + y) + \log_4 6$  as a single logarithm

**Solution 11**

$$\log_4(x + y) + \log_4 6$$

$$= \log_4((x + y) \times 6)$$

$$= \log_4 6(x + y)$$

$$4 \log_{(x+y)} 6$$

$$\log_{(x+y)} 24$$

$$4 \log_6 x + y$$

$$6 \log_4 x + y$$

$$\log_4 6(x + y)$$

**Question 12** Experience: 10 Order: a1 Level: a1

Express  $\log_x 9 = 2$  in power form

**Solution 12**

$$\log_x 9 = 2$$

$$x^2 = 9$$

$$x^9 = 2$$

$$x^2 = 2$$

$$x^9 = 9$$

$$x^2 = 7$$

$$x^2 = 9$$

**Question 13** Experience: 15 Order: c1 Level: c1

Express  $3 \log_3(a + b) + \log_3 4$  as a single logarithm

**Solution 13**

$$3 \log_3(a + b) + \log_3 4$$

$$3 \log_{a+b} (a + b)^3 + \log_3 4$$

$$\log_{a+b}^{(12)} ((a + b)^3 \times 4)$$

$$12 \log_3 a + b$$

$$4 \log_3 (a + b)^3$$

$$\log_3 4(a+b)^3$$

**Question 14** Experience: 15 Order: c2 Level: c2

Express  $\log_4(a^2 - b^2) - 2\log_4(a + b)$  as a single logarithm

**Solution 14**

$$\log_4 \frac{a^2 - b^2}{(a+b)^2} - 2\log_4(a+b)$$

$$\log_4 \frac{(a+b)(a-b)}{(a+b)^2} - \log_4(a+b)^2$$

$$\log_4 \frac{a-b}{a+b} - \log_4((a+b)^2 \div (a+b)^2)$$

$$\log_4 \frac{a-b}{a+b} - \log_4 \left( \frac{(a+b)(a-b)}{(a+b)^2} \right)$$

$$\log_4 \frac{a-b}{a+b} - \log_4 \frac{a-b}{a+b}$$

$$\log_4 1$$

**Question 15** Experience: 15 Order: c2 Level: c2

Express  $\log_x(4a - 6b) + \log_x \frac{1}{2}$  as a single logarithm

**Solution 15**

$$\log_x(4a - 6b) + \log_x \frac{1}{2}$$

$$= \log_x \frac{1}{2}(4a - 6b)$$

$$= \log_x(2a - 3b)$$

$$\log_{(4a-6b)} \frac{1}{2}^x$$

$$\log_x(4a - 6b)$$

$$\frac{1}{2} \log_{(4a-6b)} x$$

$$\frac{1}{2} \log_x(2a - 3b)$$

$$\log_x(2a - 3b)$$

**Question 16** Experience: 15 Order: d1 Level: d1

Express  $\log_4(6a) - \log_4(2a)$  as a single logarithm

**Solution 16**

$$\log_4(6a) - \log_4(2a)$$

$$= \log_4(6a \div 2a)$$

$$= \log_4 3$$

$$\log_a 2$$

$$\log_a 3$$

$$\log_4 3a$$

$$\log_4 12a^2$$

$$\log_4 3$$

**Question 17** Experience: 15 Order: d1 Level: d1

Express  $\log_{10}(15) - \log_{10}(3)$  as a single logarithm

**Solution 17**

$$\begin{aligned}\log_{10}(15) - \log_{10}(3) \\&= \log_{10}(15 \div 3) \\&= \log_{10} 5\end{aligned}$$

$$\log_{10} 3$$

$$\log_5 45$$

$$\log_{10} 45$$

$$\log_5 10$$

$$\log_{10} 5$$

**Question 18**      Experience: 15      Order: d2      Level: d2

Express  $3 \log_y(5) + \log_y(4)$  as a single logarithm

**Solution 18**

$$\begin{aligned}3 \log_y(5) + \log_y(4) \\&= \log_y 5^3 + \log_y 4 \\&= \log_y(5^3 \times 4) \\&= \log_y 500\end{aligned}$$

$$\log_y 100$$

$$\log_y 8000$$

$$4 \log_y 125$$

$$4 \log_y 50$$

$$\log_y 500$$

**Question 19**      Experience: 15      Order: d2      Level: d2

Express  $3 \log_a(4) - 4 \log_a(2)$  as a single logarithm

**Solution 19**

$$\begin{aligned}\log_a(4^3) - \log_a(2^4) \\&= \log_a(64) - \log_a(16) \\&= \log_y(64 \div 16) \\&= \log_y 4\end{aligned}$$

$$\log_4 a^2$$

$$\log_4 16$$

$$\log_4 64$$

$$\log_y 16$$

$$\log_y 4$$

**Question 20**      Experience: 10      Order: a1      Level: a1

Express  $\log_3 7 = a + b^2$  in power form

**Solution 20**

$$\log_3 7 = a + b^2$$

$$3^{a+b^2} = 7$$

$$7^{a+b^2} = 3$$

$$3^7 = 7a + b^2$$

$$(a + b^2)^3 = 7$$

$$3^7 = a + b^2$$

$$3^{a+b^2} = 7$$

**Question 21** Experience: 30 Order: d3 Level: d3

Express  $4 \log_9 5 - 2 \log_3(15)$  as a single logarithm

**Solution 21**

$$\begin{aligned} & 4 \log_9 5 - 2 \log_3(9) \\ &= 4 \left( \frac{\log_3 5}{\log_3 9} \right) - 2 \log_3(15) \\ &= \left( \frac{4 \log_3 5}{2} \right) - \log_3(15^2) \\ &= 2 \log_3 5 - \log_3(15^2) \\ &= \log_3(5^2 \div 15^2) \\ &= \log_3 \frac{25}{225} \\ &= \log_3 \frac{1}{9} \end{aligned}$$

$$\log_9 225$$

$$\log_9 25$$

$$\log_9 \frac{1}{3}$$

$$\log_3 9$$

$$\log_3 \frac{1}{9}$$

**Question 22** Experience: 15 Order: c1 Level: c1

Express  $\log_a 4 + \log_a 5$  as a single logarithm

**Solution 22**

$$\begin{aligned} & \log_a 4 + \log_a 5 \\ &= \log_a (4 \times 5) \\ &= \log_a 20 \end{aligned}$$

$$\log_4 5a$$

$$a \log_4 5$$

$$4 \log_a 5$$

$$5 \log_a 4$$

$$\log_a 20$$

**Question 23** Experience: 30 Order: d3 Level: d3

Express  $2 \log_{16} 8 - 4 \log_4(2)$  as a single logarithm

## Solution 23

$$\begin{aligned} & 2 \log_{16} 8 - 4 \log_4(2) \\ &= 2 \left( \frac{\log_4 8}{\log_4 16} \right) - 4 \log_4(2) \\ &= \left( \frac{2 \log_4 8}{2} \right) - \log_4(16) \\ &= \log_4(8 \div 16) \\ &= \log_4 \frac{1}{2} \end{aligned}$$

$$\log_8 4$$

$$\log_8 2$$

$$\log_8 \frac{1}{2}$$

$$\log_4 \frac{2}{2}$$

$$\log_4 \frac{1}{2}$$

## OCR A Level Maths

To be added .....

## AQA A Level Maths

To be added .....

## MEI A Level Maths

To be added .....

## Unit 1 C 1

## Chapter 1 asdfasfd

## Edexcel A Level Further Maths

To be added .....

## MEI A Level Further Maths

To be added .....