

CSDS 455: Homework 13

Shaochen (Henry) ZHONG, sxz517

Due and submitted on 10/07/2020
Fall 2020, Dr. Connamacher

Problem 1

I consulted <https://www.youtube.com/watch?v=otky1bBhwgM> for this problem.

Proof. We will have the following diagram for a base graph:




Figure 1: Base graph for *Problem 1*

For better clarity, we have $u_1, u_2, \dots, u_i \in N(x)$ being the adjacent vertices of x excluding y . Known that $\chi'(G - xy) = \Delta + 1$, every vertex in $G - xy$ must be missing¹ at least one color from the $\Delta + 1$ colors, we use dashed lines to represent such colors. We also use solid lines to represent the color of edges under a $\Delta + 1$ edge-coloring of $G - xy$. We will also use some arbitrary actual color names to help understanding.

We want to show that other than having an 2-color alternating path between x and y , we will always have $\chi'(G) = \Delta + 1$, thus a contradiction.

Proposition 1. x and y can't miss the same color.


This observation is almost trivial as otherwise we may just color edge xy with this color, then we have a $\Delta + 1$ coloring of G – a contradiction.

¹ “Missing” in this context means not having an edge of a certain color. “Vertex v is missing color c ” means there is no edge of color c connected to v


Proposition 2. x must include every missing color of $N(x)$.

Say we have a set S which include all the missing colors of $N(x)$, such S must be a subset of the colors connected x . This is because if we have a vertex $u_i \in N(x)$ which has a missing color of **purple** which is not included in the connected colors of x , we have two cases.

First, u_i is u_1 , where edge u_1x has the color which y misses, then we may recolor the graph as following. Which will make x and y both missing **orange**; we can therefore color xy with **orange** and have a $\Delta + 1$ coloring of G – a contradiction.



Second, we may have u_ix not having the color which y misses. We can then recolor u_ix to be **purple**, and recolor the original **blue** u_jx (u_j is another neighbor vertex of x which is missing **green**) to be **green**, and recolor another neighbor vertex of x which is missing the color of u_jx ... Take $u_3 = u_i$ as an example, we will have:




We know this recoloring method works since $N(x)$ is finite, so we will always able to recolor u_1x (**orange**, which y misses) to the missing color of u_1 (**blue**), then recolor the **blue** edge u_jx to its missing color, then we will eventually reaches u_i and recolor u_ix to its missing color **purple**. Then we will have x and y both missing **orange** and thus a contradiction.

Now we have showed that every missing color of $N(X)$ must show on an edge of ux for $u \in N(x)$. Be familiar with this recoloring method as we will use it later.

Proposition 3. If x misses **red**. Every $u \in N(x)$ must have an **red** edge connected to u .

Because otherwise, by using the recoloring method introduced in **Proposition 2**, we may have x and y both missing **orange** and thus a contradiction. The following is an example assuming u_3 missing **red**, but it can be any $u \in N(x)$.




Combining the finding of **Proposition 1, 2, 3**, we will have a graph like [Figure 1].

Proposition 4. If x misses **red** and y misses **orange**, the maximal **orange-red** alternating path from x can't end on a vertex $\notin N(x)$ and not y .

If we consider a subgraph G' with only **orange** and **red** edges by the coloring of $G - xy$, we know that every vertex in G' will have a degree of 0, 1, or 2. Now we take a **orange-red** alternating walk P from x , until it reaches a vertex $v \in V(G)$ where we can't extend the walk any farther (which implies $d(v) = 1$). We let $v \notin N(x)$ and $v \neq y$ in this case.

We can't have this kind of v as we may simply interchange the **orange-red** coloring of P . We may then have x and y both missing **orange** and thus a contradiction. So we know a **orange-red** alternating walk P from x can't end on a vertex outside of $N(x)$ and y .




Proposition 4. If x misses **red** and y misses **orange**, the maximal **orange-red** alternating path from x can't end on a vertex $\in N(x)$.

It is trivial to tell a **orange-red** alternating walk P from x can't end on u_1 (where xu_1 is the first edge of P). P may either end on a u_i where u_i misses **orange** (like u_3 in the below graph); or P will visit another $u_j \in N(x)$ which has **orange** edge connected to it, but in this case this P will not end on $N(x)$.

We first show the case of P ending on a vertex $u_i \in N(x)$ where u_i misses the same color as y (**orange**). In this case we interchange the color in P as following and we have x and y both missing **orange** – a contradiction.

In the case of P ending on a vertex $u_j \in N(x)$ where u_j has **orange** edge connected to it. We can tell P will not end on $N(x)$ as:

1. $u_{j-1}u_j \in E(P)$ is **orange**, then we may extend this P farther to include the **red** edge connected to u_j .
2. $u_{j-1}u_j \in E(P)$ is **red**, in this case we may extend this P farther by including the **orange** edge connected to u_j .




Either of the case will make P end on a vertex v outside of $N(x)$ and y . According to **Proposition 2**, this will give us a contradiction.

Conclusion: P must end on y .

By **Proposition 3, 4**, we know that a **orange-red** alternating walk P from x can't end on $N(x)$, can't end on $G - y$, and obviously it can't end on x itself as x misses **red**. So the only left option it to end on y .

In the following diagram we will show by having P ends on y , we can't make a contradiction by doing the recoloring operation.



It is clear that although an interchange recoloring of P is possible, we only swapped the missing color between x and y , so we can't color xy with an existing color and no contradiction is made.

The statement in question is therefore proven. □

Problem 2