

# Corrigendum to “Effects of training with elastic resistance versus conventional resistance on muscular strength: A systematic review and meta-analysis”

SAGE Open Medicine

Volume 8: 2

© The Author(s) 2020

Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/2050312120961220

journals.sagepub.com/home/smo



Lopes JSS, Machado AF, Micheletti JK, et al. Effects of training with elastic resistance versus conventional resistance on muscular strength: A systematic review and meta-analysis. *SAGE Open Medicine* 2019; Feb 19; 7. doi: 10.1177/2050312119831116

The authors would like to draw attention to the following errors in the above paper:

The standardized mean difference (SMD) in the Abstract should have been stated as **-0.11** instead of -0.011.

The sentence in the Abstract which reads “The results of the meta-analysis demonstrated no superiority between the methods analyzed for upper limb (SMD = -0.011, 95% CI -0.40, 0.19,  $p = 0.48$ ) or lower limb muscular strength (SMD=0.09; 95% CI -0.18, 0.35;  $p=0.52$ )” should have read **“The results of the meta-analysis demonstrated no superiority between the methods analyzed for lower limb (SMD = -0.11, 95% CI -0.40, 0.19,  $p = 0.48$ ) or upper limb muscular strength (SMD=0.09; 95% CI -0.18, 0.35;  $p = 0.52$ )”.**

The legends of Figures 2 and 3 were mistakenly swapped. The forest plot of Figure 2 output contains data for lower limb outcomes and should have been labelled **“lower limbs”** instead of “upper limbs”. **The opposite is true of Figure 3.**

For the sake of clarity, it is reiterated that the authors used a fixed-effect model for the meta-analysis **“through the I2 value”** (the I2 value being mentioned in the Methods section).

In the Results section, the sentences which read “The search carried out in the databases identified a total of 365 articles, of which 23 were considered eligible. Of these, 10 studies were excluded as they did not compare training between elastic devices and conventional machines; three did not use muscular strength as an outcome; and two did not perform training with elastic devices. Thus, eight articles (Table 1) corresponded to the inclusion criteria, comprising a total of 224 individuals aged between 15 and 88 years” should have read **“The search carried out in the databases identified a total of 365 articles, of which 29 were considered eligible. Of these, 17 studies were excluded as they did not compare training between elastic devices and**

**conventional machines; three did not use muscular strength as an outcome; and two did not perform training with elastic devices. Thus, seven articles (Table 1) corresponded to the inclusion criteria, comprising a total of 224 individuals aged between 15 and 88 years”.**

The PRISMA checklist in Figure 1 should have noted **seven studies (n = 7)** rather than eight.

In the Results sub-section “Methodological quality of included studies”, the sentence “One study<sup>3</sup> scored 8; four<sup>1,6</sup> scored 7; one scored 6, and two<sup>2</sup> scored 5 on the scale, thus classifying the articles as “moderate quality” according to the classification used in the review and meta-analysis study by Machado et al<sup>15</sup> (Table 2)” should have read **“One study<sup>3</sup> scored 8; three<sup>1,7,8</sup> scored 7; one scored 6<sup>4</sup>, and two<sup>2,6</sup> scored 5 on the scale, thus classifying the articles as “moderate quality” according to the classification used in the review and meta-analysis study by Machado et al<sup>15</sup> (Table 2)”.**

In the Results sub-section entitled “Effects of elastic resistance training and conventional resistance training on muscular strength”, the sentence “The results of the meta-analysis for the upper and lower limbs, respectively, showed that there is no superiority between training performed with elastic resistance and training with weight machines and/or free weights on strength gain (Upper limbs: SMD = -0.11, 95% CI -0.40, 0.19;  $p = 0.48$ ; Lower limbs: SMD = 0.09; 95% CI -0.18, 0.35;  $p = 0.52$ ) (Figs 2 and 3) should have read **“The results of the meta-analysis for the upper and lower limbs, respectively, showed that there is no superiority between training performed with elastic resistance and training with weight machines and/or free weights on strength gain (Lower limbs: SMD = -0.11, 95% CI -0.40, 0.19;  $p = 0.48$ ; Upper limbs: SMD = 0.09; 95% CI -0.18, 0.35;  $p = 0.52$ ) (Figs 2 and 3)”.**

In the Discussion section, the sentence which read “The findings are in accordance with systematic reviews with meta-analysis that demonstrated positive effects on muscular strength gain from the use of elastic resistance when compared with a control group in the elderly<sup>16</sup> and individuals with osteoarthritis<sup>17</sup> and fibromyalgia<sup>18</sup>” should



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons

Attribution-NonCommercial 4.0 License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

have read “The findings are in accordance with systematic reviews with meta-analysis that demonstrated positive effects on muscular strength gain from the use of elastic resistance when compared with a control group in the elderly <sup>16</sup>. Other studies reviewed all types of resistance training, including elastic resistance in specific

populations, which included individuals with osteoarthritis and fibromyalgia, and found positive outcomes on this method <sup>17,18</sup>”.

The authors would like to thank Dr Phil Page (Baton Rouge, Louisiana, USA) for highlighting the errors.