

108. Silder A, Thelen DG, Heiderscheit BC. Effects of prior hamstring strain injury on strength, flexibility and running mechanics. *Clin Biomech (Bristol, Avon)* 2010;25(7):681-6.
109. Barreira P, Drust B, Robinson MA et al. Asymmetry after hamstring injury in English Premier League: issue resolved or perhaps not? *Int J Sports Med* 2015;36(6):455-9.
110. Opar DA, Williams MD, Timmins RG et al. Knee flexor strength and bicep femoris electromyographical activity is lower in previously strained hamstrings. *J Electromyogr Kinesiol* 2013;23(3):696-703.
111. Opar DA, Williams MD, Timmins RG et al. Rate of torque and electromyographic development during anticipated eccentric contraction is lower in previously strained hamstrings. *Am J Sports Med* 2013;41(1):116-25.
112. Sole G, Milosavljevic S, Nicholson HD et al. Selective strength loss and decreased muscle activity in hamstring injury. *J Orthop Sports Phys Ther* 2011;41(5):354-63.
113. Silder A, Heiderscheit BC, Thelen DG et al. MR observations of long-term musculotendon remodelling following a hamstring strain injury. *Skeletal Radiol* 2008;37(12):1101-9.
114. Timmins RG, Shield AJ, Williams MD et al. Biceps femoris long head architecture: a reliability and retrospective injury study. *Med Sci Sports Exerc* 2015;47(5):905-13.
115. Silder A, Reeder SB, Thelen DG. The influence of prior hamstring injury on lengthening muscle tissue mechanics. *J Biomech* 2010;43(12):2254-60.
116. Koulouris G, Connell DA, Brukner P et al. Magnetic resonance imaging parameters for assessing risk of recurrent hamstring injuries in elite athletes. *Am J Sports Med* 2007;35(9):1500-6.
117. Thorborg K. What are the most important risk factors for hamstring muscle injury? *Clin J Sport Med* 2014;24(2):160-1.
118. Opar DA, Williams MD, Timmins RG et al. Eccentric hamstring strength and hamstring injury risk in Australian footballers. *Med Sci Sports Exerc* 2015;47(4):857-65.
119. Timmins RG, Bourne MN, Shield AJ et al. Short biceps femoris fascicles and eccentric knee flexor weakness increase the risk of hamstring injury in elite football (soccer): a prospective cohort study. *Br J Sports Med* 2015 Dec 16. pii: bjsports-2015-095362. doi: 10.1136/bjsports-2015-095362. [Epub ahead of print]
120. Evangelidis PE, Massey GJ, Pain MT et al. Biceps femoris aponeurosis size: a potential risk factor for strain injury? *Med Sci Sports Exerc* 2015;47(7):1383-9.
121. Fiorentino NM, Blemker SS. Musculotendon variability influences tissue strains experienced by the biceps femoris long head muscle during high-speed running. *J Biomech* 2014;47(13):3325-33.
122. Fiorentino NM, Epstein FH, Blemker SS. Activation and aponeurosis morphology affect in vivo muscle tissue strains near the myotendinous junction. *J Biomech* 2012;45(4):647-52.
123. Gabbe BJ, Bennell KL, Finch CF et al. Predictors of hamstring injury at the elite level of Australian football. *Scand J Med Sci Sports* 2006;16(1):7-13.
124. Bradley PS, Portas MD. The relationship between preseason range of motion and muscle strain injury in elite soccer players. *J Strength Cond Res* 2007;21(4):1155-9.
125. Goldman EF, Jones DE. Interventions for preventing hamstring injuries. *Cochrane Database Syst Rev* 2010(1):CD006782.
126. Goode AP, Reiman MP, Harris L et al. Eccentric training for prevention of hamstring injuries may depend on intervention compliance: a systematic review and meta-analysis. *Br J Sports Med* 2015;49(6):349-56.
127. Askling C, Karlsson J, Thorstensson A. Hamstring injury occurrence in elite soccer players after preseason strength training with eccentric overload. *Scand J Med Sci Sports* 2003;13(4):244-50.
128. Gabbe BJ, Branson R, Bennell KL. A pilot randomised controlled trial of eccentric exercise to prevent hamstring injuries in community-level Australian Football. *J Sci Med Sport* 2006; 9(1-2):103-9.
129. Engebretsen AH, Myklebust G, Holme I et al. Prevention of injuries among male soccer players: a prospective, randomised intervention study targeting players with previous injuries or reduced function. *Am J Sports Med* 2008;36(6):1052-60.
130. Kraemer R, Knobloch K. A soccer-specific balance training program for hamstring muscle and patellar and achilles tendon injuries: an intervention study in premier league female soccer. *Am J Sports Med* 2009;37(7):1384-93.
131. Soderman K, Werner S, Pietila T et al. Balance board training: prevention of traumatic injuries of the lower extremities in female soccer players? a prospective randomised intervention study. *Knee Surg Sports Traumatol Arthrosc* 2000;8(6):356-63.
132. Emery CA, Rose MS, McAllister JR et al. A prevention strategy to reduce the incidence of injury in high school basketball: a cluster randomised controlled trial. *Clin J Sport Med* 2007;17(1):17-24.
133. Orchard J. Understanding some of the risks for soft tissue injury—a Malcolm Blight legacy. *J Sci Med Sport* 2002;5(2):v-vii.
134. Nedelec M, McCall A, Carling C et al. The influence of soccer playing actions on the recovery kinetics after a soccer match. *J Strength Cond Res* 2014;28(6):1517-23.
135. McCall A, Nedelec M, Carling C et al. Reliability and sensitivity of a simple isometric posterior lower limb muscle test in professional football players. *J Sports Sci* 2015;33(12):1298-304.
136. Schache AG, Crossley KM, Macindoe IG et al. Can a clinical test of hamstring strength identify football players at risk of hamstring strain? *Knee Surg Sports Traumatol Arthrosc* 2011;19(1):38-41.
137. Thorborg K, Bandholm T, Hölmich P. Hip- and knee-strength assessments using a hand-held dynamometer with external belt-fixation are inter-tester reliable. *Knee Surg Sports Traumatol Arthrosc* 2013;21:550-5.
138. Wollin M, Purdam C, Drew MK. Reliability of externally fixed dynamometry hamstring strength testing in elite youth football players. *J Sci Med Sport* 2016;19(1):93-6.

139. Kornberg C, Lew P. The effect of stretching neural structures on grade one hamstring injuries. *J Orthop Sports Phys Ther* 1989;10(12):481-7.
140. Huguenin L, Brukner PD, McCrory P et al. Effect of dry needling of gluteal muscles on straight leg raise: a randomised, placebo controlled, double blind trial. *Br J Sports Med* 2005;39(2):84-90.
141. Bogduk N. *Clinical Anatomy of the Lumbar Spine and Sacrum*. 4th ed. Edinburgh: Elsevier Churchill Livingstone, 2004.
142. Barash HL, Galante JO, Lambert CN et al. Spondylolisthesis and tight hamstrings. *J Bone Joint Surg Am* 1970;52(7):1319-28.
143. O'Sullivan PB, Twomey L, Allison GT. Dynamic stabilisation of the lumbar spine. *Clin Rev Phys Rehab Med* 1997;9(3&4): 315-30.
144. Misaggi B, Gallazzi M, Colombo M et al. Articular facets syndrome: diagnostic grading and treatment options. *Eur Spine J* 2009;18 Suppl 1(1):49-51.
145. Koulouris G, Connell D. Evaluation of the hamstring muscle complex following acute injury. *Skeletal Radiol* 2003;32(10):582-9.
146. Subbu R, Benjamin-Laing H, Haddad F. Timing of surgery for complete proximal hamstring avulsion injuries successful clinical outcomes at 6 weeks, 6 months and after 6 months of injury. *Am J Sports Med* 2015;43(2):385-91.
147. Wood DG, Packham I, Trikha SP et al. Avulsion of the proximal hamstring origin. *J Bone Joint Surg Am* 2008;90(11): 2365-74.
148. Harris JD, Griesser MJ, Best TM et al. Treatment of proximal hamstring ruptures—a systematic review. *Int J Sports Med* 2011;32(7):490-5.
149. Askling CM, Koulouris G, Werner S et al. Total proximal hamstring ruptures. Clinical and MRI aspects including guidelines for postoperative rehabilitation. *Knee Surg Sports Arthrosc* 21:515-533, 2013.
150. Zissen MH, Wallace G, Stevens KJ et al. High hamstring tendinopathy: MRI and ultrasound imaging and therapeutic efficacy of percutaneous corticosteroid injection. *AJR Am J Roentgenol* 2010;195(4):993-8.
151. Jayaseelan DJ, Moats N, Ricardo CR. Rehabilitation of proximal hamstring tendinopathy utilising eccentric training, lumbopelvic stabilisation and trigger point dry needling: 2 case reports. *J Orthop Sports Phys Ther* 2014;44(3):198-205.
152. Cacchio A, Rompe JD, Furia JP et al. Shockwave therapy for the treatment of chronic proximal hamstring tendinopathy in professional athletes. *Am J Sports Med* 2011;39(1):146-53.
153. Cacchio A, De Paulis F, Maffulli N. Development and validation of a new visa questionnaire (VISA-H) for patients with proximal hamstring tendinopathy. *Br J Sports Med* 2014;48(6):448-52.
154. Puranen J, Orava S. The hamstring syndrome. A new diagnosis of gluteal sciatic pain. *Am J Sports Med* 1988;16(5):517-21.
155. McCrory PM, Bell SN, Bradshaw C. Nerve entrapment of the lower leg, ankle and foot in sport. *Sports Med* 2002;32(6):371-91.
156. Orava S, Rantanen J, Kujala UM. Fasciotomy of the posterior femoral muscle compartment in athletes. *Int J Sports Med* 1998;19(1):71-5.
157. Abraham P, Saumet JL, Chevalier JM. External iliac artery endofibrosis in athletes. *Sports Med* 1997;24(4):221-6.

CHAPTER 35

1. Myklebust G, Bahr R. Return to play guidelines after anterior cruciate ligament surgery. *Br J Sports Med* 2005;39(3):127-31.
2. Bachmann LM, Habereith S, Steurer J et al. The accuracy of the Ottawa knee rule to rule out knee fractures: a systematic review. *Ann Intern Med* 2004;140(2):121-4.
3. Eck C, Bekerom M, Fu F et al. Methods to diagnose acute anterior cruciate ligament rupture: a meta-analysis of physical examinations with and without anaesthesia. *Knee Surg Sports Traumatol Arthrosc* 2013;21(8):1895-903.
4. Kopkow C, Freiberg A, Kirschner S et al. Physical examination tests for the diagnosis of posterior cruciate ligament rupture: a systematic review. *J Orthop Sports Phys Ther* 2013;43(11): 804-13.
5. Smith BE, Thacker D, Crewesmith A et al. Special tests for assessing meniscal tears within the knee: a systematic review and meta-analysis. *Evid Based Med* 2015;20(3):88-97.
6. Hegedus EJ, Cook C, Hasselblad V et al. Physical examination tests for assessing a torn meniscus in the knee: a systematic review with meta-analysis. *J Orthop Sports Phys Ther* 2007;37(9):541-50.
7. Kastelein M, Wagemakers HP, Luijsterburg PA et al. Assessing medial collateral ligament knee lesions in general practice. *Am J Med* 2008;121(11):982-8.e2.
8. Ebrahimzadeh MH, Makhmalbaf H, Golhasani-Keshtan F et al. The International Knee Documentation Committee (IKDC) Subjective Short Form: a validity and reliability study. *Knee Surg Sports Traumatol Arthrosc* 2015;23(11):3163-7.
9. Padua R, Bondi R, Ceccarelli E et al. Italian version of the International Knee Documentation Committee Subjective Knee Form: cross-cultural adaptation and validation. *Arthroscopy* 2004;20(8):819-23.
10. van Meer BL, Meuffels DE, Vissers MM et al. Knee injury and Osteoarthritis Outcome Score or International Knee Documentation Committee Subjective Knee Form: which questionnaire is most useful to monitor patients with an anterior cruciate ligament rupture in the short term? *Arthroscopy* 2013;29(4):701-15.
11. Crawford K, Briggs KK, Rodkey WG et al. Reliability, validity and responsiveness of the IKDC score for meniscus injuries of the knee. *Arthroscopy* 2007;23(8):839-44.
12. Metsavaht L, Leporace G, Riberto M et al. Translation and cross-cultural adaptation of the Brazilian version of the International Knee Documentation Committee Subjective Knee Form: validity and reproducibility. *Am J Sports Med* 2010;38(9):1894-9.

13. Çelik D, Coşkunsu D, Kiliçoğlu Ö et al. Translation and cross-cultural adaptation of the International Knee Documentation Committee Subjective Knee Form into Turkish. *J Orth Sports Phys Ther* 2014;44(11):899-909.
14. Kim JG, Ha JK, Lee JY et al. Translation and validation of the Korean version of the International Knee Documentation Committee Subjective Knee Form. *Knee Surg Rel Res* 2013;25(3):106-11.
15. Briggs KK, Lysholm J, Tegner Y et al. The reliability, validity, and responsiveness of the Lysholm Score and Tegner Activity Scale for anterior cruciate ligament injuries of the knee: 25 years later. *Am J Sports Med* 2009;37(5):890-7.
16. Haverkamp D, Siersevelt IN, Breugem SJM et al. Translation and validation of the Dutch version of the International Knee Documentation Committee Subjective Knee Form. *Am J Sports Med* 2006;34(10):1680-4.
17. Agel J, LaPrade RF. Assessment of differences between the modified Cincinnati and International Knee Documentation Committee patient outcome scores: a prospective study. *Am J Sports Med* 2009;37(11):2151-7.
18. Greco NJ, Anderson AF, Mann BJ et al. Responsiveness of the International Knee Documentation Committee Subjective Knee Form in comparison to the Western Ontario and McMaster Universities Osteoarthritis Index, Modified Cincinnati Knee Rating System and Short Form 36 in patients with focal articular cartilage defects. *Am J Sports Med* 2010;38(5):891-902.
19. Irrgang JJ, Anderson AF, Boland AL et al. Responsiveness of the International Knee Documentation Committee Subjective Knee Form. *Am J Sports Med* 2006;34(10):1567-73.
20. Collins NJ, Misra D, Felson DT et al. Measures of knee function: International Knee Documentation Committee (IKDC) Subjective Knee Evaluation Form, Knee Injury and Osteoarthritis Outcome Score (KOOS), Knee Injury and Osteoarthritis Outcome Score Physical Function Short Form (KOOS-PS), Knee Outcome Survey Activities of Daily Living Scale (KOS-ADL), Lysholm Knee Scoring Scale, Oxford Knee Score (OKS), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), Activity Rating Scale (ARS) and Tegner Activity Score (TAS). *Arthritis Care Res* 2011;63 Suppl 11:S208-28.
21. Collins NJ, Prinsen CAC, Christensen R et al. Knee Injury and Osteoarthritis Outcome Score (KOOS): systematic review and meta-analysis of measurement properties. *Osteoarth Cart* 2016; 24(8):1317-29.
22. Tegner Y, Lysholm J. Rating systems in the evaluation of knee ligament injuries. *Clin Orthop Relat Res* 1985;198:43-9.
23. Briggs KK, Kocher MS, Rodkey WG et al. Reliability, validity, and responsiveness of the Lysholm knee score and Tegner activity scale for patients with meniscal injury of the knee. *J Bone Joint Surg Am* 2006;88(4):698-705.
24. Kocher MS, Steadman JR, Briggs KK et al. Reliability, validity and responsiveness of the Lysholm knee scale for various chondral disorders of the knee. *J Bone Joint Surg Am* 2004; 86-a(6):1139-45.
25. Paxton EW, Fithian DC, Stone ML et al. The reliability and validity of knee-specific and general health instruments in assessing acute patellar dislocation outcomes. *Am J Sports Med* 2003;31(4):487-92.
26. Fithian DC, Paxton EW, Stone ML et al. Prospective trial of a treatment algorithm for the management of the anterior cruciate ligament injured knee. *Am J Sports Med* 2005;33(3):335-46.
27. Marx RG, Stump TJ, Jones EC et al. Development and evaluation of an activity rating scale for disorders of the knee. *Am J Sports Med* 2001;29(2):213-8.
28. Stiell IG, Greenberg GH, Wells GA et al. Prospective validation of a decision rule for the use of radiography in acute knee injuries. *JAMA* 1996;275(8):611-5.
29. Englund M, Guermazi A, Gale D et al. Incidental meniscal findings on knee MRI in middle-aged and elderly persons. *N Engl J Med* 2008;359(11):1108-15.
30. Frobell RB, Roos HP, Roos EM et al. The acutely ACL injured knee assessed by MRI: are large volume traumatic bone marrow lesions a sign of severe compression injury? *Osteoarth Cart* 2008;16(7):829-36.
31. Johnson DL, Urban WP, Caborn DNM et al. Articular cartilage changes seen with magnetic resonance imaging-detected bone bruises associated with acute anterior cruciate ligament rupture. *Am J Sports Med* 1998;26(3):409-14.
32. DeHaven KE, Bronstein RD. Arthroscopic medial meniscus repair in the athlete. *Clin Sports Med* 1997;16:69-86.
33. Pyne S. Current progress in meniscal repair and postoperative rehabilitation. *Curr Sports Med Reports* 2002;1:265-71.
34. Herrlin S, Hallander M, Wange P et al. Arthroscopic or conservative treatment of degenerative medial meniscal tears: a prospective randomised trial. *Knee Surg Sports Traumatol Arthrosc* 2007;15(4):393-401.
35. Sihvonen R, Paavola M, Malmivaara A et al. Arthroscopic partial meniscectomy versus sham surgery for a degenerative meniscal tear. *N Engl J Med* 2013;369(26):2515-24.
36. Kise NJ, Risberg MA, Stensrud S et al. Exercise therapy versus arthroscopic partial meniscectomy for degenerative meniscal tear in middle-aged patients: randomised controlled trial with two-year follow-up. *BMJ* 2016;354:i3740.
37. Suter LG, Fraenkel L, Losina E et al. Medical decision making in patients with knee pain, meniscal tear and osteoarthritis. *Arthritis Rheum* 2009;61(11):1531-8.
38. Rodeo SA. Instructional course lectures, the American Academy of Orthopaedic Surgeons—arthroscopic meniscal repair with use of the outside-in technique. *J Bone Joint Surg Am* 2000;82:127-41.
39. Cannon WD, Jr., Vittori JM. The incidence of healing in arthroscopic meniscal repairs in anterior cruciate ligament-reconstructed knees versus stable knees. *Am J Sports Med* 1992;20(2):176-81.
40. Nawabi DH, Cro S, Hamid IP et al. Return to play after lateral meniscectomy compared with medial meniscectomy in elite professional soccer players. *Am J Sports Med* 2014 Sept;42(9):2193-8.

41. Arendt E. *OKU Orthopaedic Knowledge Update: Sports Medicine 2*. 2nd ed. Rosemont, IL: American Academy of Orthopaedic Surgeons, 1999.
42. Indelicato PA. Isolated medial collateral injuries in the knee. *J Am Acad Orthop Surg* 1995;3(1):9-14.
43. Frobell RB, Lohmander LS, Roos HP. Acute rotational trauma to the knee: poor agreement between clinical assessment and magnetic resonance imaging findings. *Scand J Med Sci Sports* 2007;17(2):109-14.
44. Spindler KP, Wright RW. Clinical practice. Anterior cruciate ligament tear. *N Engl J Med* 2008;359(20):2135-42.
45. Gottlob CA, Baker CL, Jr., Pellissier JM et al. Cost effectiveness of anterior cruciate ligament reconstruction in young adults. *Clin Orthop Relat Res* 1999;367:272-82.
46. Arendt E, Dick R. Knee injuries patterns among men and women in collegiate basketball and soccer. NCAA data and review of literature. *Am J Sports Med* 1995;23:694-701.
47. Arendt EA, Agel J, Dick R. Anterior cruciate ligament injury patterns among collegiate men and women. *J Athl Train* 1999;34:86-92.
48. Garrick JG, Requa RK. Anterior cruciate ligament injuries in men and women: how common are they? In: Gryphon LY, ed. *Prevention of Noncontact ACL Injuries*. Rosemont, IL: American Academy of Orthopaedic Surgeons, 2001:1-10.
49. Agel J, Arendt E, Bershadsky B. Anterior cruciate ligament injury in national collegiate athletic association basketball and soccer: a 13-year review. *Am J Sports Med* 2005;33(4):524-30.
50. Beynnon BD, Johnson RJ, Abate JA et al. Treatment of anterior cruciate ligament injuries, part 2. *Am J Sports Med* 2005;33(11):1751-67.
51. Alentorn-Geli E, Myer GD, Silvers HJ et al. Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 1: Mechanisms of injury and underlying risk factors. *Knee Surg Sports Traumatol Arthrosc* 2009;17(7):705-29.
52. John R, Dhillon MS, Sharma S et al. Is there a genetic predisposition to anterior cruciate ligament tear? A systematic review. *Am J Sports Med* 2016 Feb 3. pii: 0363546515624467. [Epub ahead of print]
53. Barber FA. What is the terrible triad? *Arthroscopy* 1992;8(1):19-22.
54. Lubowitz JH, Hwang M, Piefer J et al. Anterior cruciate ligament femoral footprint anatomy: systematic review of the 21st century literature. *Arthroscopy* 2014;30:539-41.
55. Sonnerly-Cottet B, Colombet P. Partial tears of the anterior cruciate ligament. *Orth Traumatol* 2016;102:S59-67.
56. McLean SG, Huang X, Su A et al. Sagittal plane biomechanics cannot injure the ACL during sidestep cutting. *Clin Biomech (Bristol, Avon)* 2004;19(8):828-38.
57. McLean SG, Huang X, van den Bogert AJ. Association between lower extremity posture at contact and peak knee valgus moment during sidestepping: implications for ACL injury. *Clin Biomech (Bristol, Avon)* 2005;20(8):863-70.
58. McLean SG, Walker K, Ford KR et al. Evaluation of a two-dimensional analysis method as a screening and evaluation tool for anterior cruciate ligament injury. *Br J Sports Med* 2005;39(6):355-62.
59. Olsen OE, Myklebust G, Engebretsen L et al. Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am J Sports Med* 2004;32(4):1002-12.
60. Teitz CC. Video analysis of ACL injuries. In: Gryphon LY, ed. *Prevention of Noncontact ACL Injuries*. Rosemont, IL: American Academy of Orthopaedic Surgeons, 2001:93-96.
61. Ireland ML. Anterior cruciate ligament injuries in young female athletes. *Your Patient & Fitness* 1996;10(5):26-30.
62. Ettlinger CF, Johnson RJ, Shealy JE. A method to help reduce the risk of serious knee sprains incurred in alpine skiing. *Am J Sports Med* 1995;23:531-7.
63. Logan MC, Williams A, Lavelle J et al. What really happens during the Lachman Test? A dynamic MRI analysis of tibiofemoral motion. *Am J Sports Med* 2004;32(2):369-75.
64. Segond P. Recherches cliniques et expérimentales sur les épanchements sanguins du genou par entorse. *Progres Med* 1879; 7: 297-9, 319-21, 340-1.
65. Claes S, Vereecke E, Maes et al. Anatomy of the anterolateral ligament of the knee. *J Anatomy* 2013;223(4):321-8.
66. Hanypsiak BT, Spindler KP, Rothrock CR et al. Twelve-year follow-up on anterior cruciate ligament reconstruction: long-term outcomes of prospectively studied osseous and articular injuries. *Am J Sports Med* 2008;36(4):671-7.
67. Frobell RB, Roos EM, Roos HP et al. A randomized trial of treatment for acute anterior cruciate ligament tears. *N Engl J Med* 2010;363(4):331-42.
68. Frobell R, Roos H, Roos E et al. Treatment for acute anterior cruciate ligament tear: five-year outcome of randomised trial. *BMJ* 2013;346:232.
69. Dunn WR, Lyman S, Lincoln AE et al. The effect of anterior cruciate ligament reconstruction on the risk of knee reinjury. *Am J Sports Med* 2004;32:1906-14.
70. Kostogiannis I, Ageberg E, Neuman P et al. Clinically assessed knee joint laxity as a predictor for reconstruction after an anterior cruciate ligament injury: a prospective study of 100 patients treated with activity modification and rehabilitation. *Am J Sports Med* 2008;36(8):1528-33.
71. Eitzen I, Moksnes H, Snyder-Mackler L et al. Functional tests should be accentuated more in the decision for ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2010;18(11):1517-25.
72. Eggerding V, Meuffels DE, Bierma-Zeinstra SM et al. Factors related to the need for surgical reconstruction after anterior cruciate ligament rupture: a systematic review of the literature. *J Orthop Sports Phys Ther* 2015;45(1):37-44.
73. Meuffels DE, Favejee MM, Vissers MM et al. Ten-year follow-up study comparing conservative versus operative treatment of anterior cruciate ligament ruptures. A matched-pair analysis of high level athletes. *Br J Sports Med* 2009;43(5):347-51.

74. Rodriguez-Merchan EC. Evidence-based ACL reconstruction. *Arch Bone Jt Surg* 2015;3(1):9-12.
75. Xie X, Liu X, Chen Z et al. A meta-analysis of bone-patellar tendon-bone autograft versus four-strand hamstring tendon autograft for anterior cruciate ligament reconstruction. *Knee* 2015;22(2):100-10.
76. Zeng C, Gao SG, Li H et al. Autograft versus allograft in anterior cruciate ligament reconstruction: a meta-analysis of randomized controlled trials and systematic review of overlapping systematic reviews. *Arthroscopy* 2016;32(1):153-63.e18.
77. Chen G, Wang S. Comparison of single-bundle versus double-bundle anterior cruciate ligament reconstruction after a minimum of 3-year follow-up: a meta-analysis of randomized controlled trials. *Int J Clin Exp Med* 2015;8(9):14604-14.
78. Desai N, Bjornsson H, Musahl V et al. Anatomic single- versus double-bundle ACL reconstruction: a meta-analysis. *Knee Surg Sports Traumatol Arthrosc* 2014;22(5):1009-23.
79. Shelbourne KD, Patel DV. Timing of surgery in anterior cruciate ligament-injured knees. *Knee Surg Sports Traumatol Arthrosc* 1995;3:148-56.
80. Andernord D, Karlsson J, Musahl V et al. Timing of surgery of the anterior cruciate ligament. *Arthroscopy* 2013;29(11):1863-71.
81. Filardo G, de Caro F, Andriolo L et al. Do cartilage lesions affect the clinical outcome of anterior cruciate ligament reconstruction? A systematic review. *Knee Surg Sports Traumatol Arthrosc* 2016 Apr 4. [Epub ahead of print].
82. Shelbourne KD, Nitz P. Accelerated rehabilitation after anterior cruciate ligament reconstruction. *Am J Sports Med* 1990;18(3):292-9.
83. van Grinsven S, van Cingel RE, Holla CJ et al. Evidence-based rehabilitation following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc* 2010;18(8):1128-44.
84. Liu-Ambrose T, Taunton JE, MacIntyre D et al. The effects of proprioceptive or strength training on the neuromuscular function of the ACL reconstructed knee. *Scan J Med Sci Sports* 2003;13:115-23.
85. Cooper RL, Taylor NF, Feller JA. A randomised controlled trial of proprioceptive and balance training after surgical reconstruction of the anterior cruciate ligament. *Res Sports Med* 2005;13:217-30.
86. Logerstedt DS, Snyder-Mackler L, Ritter RC et al. Knee stability and movement coordination impairments: knee ligament sprain: clinical practice guidelines linked to the international classification of functioning, disability and health from the Orthopaedic Section of the American Physical Therapy Association. *J Orth Sports Phys Ther* 2010;40(4):A1-A37.
87. Wright RW, Haas AK, Anderson J et al. Anterior cruciate ligament reconstruction rehabilitation: MOON guidelines. *Sports Health* 2015;7(3):239-43.
88. Beynnon BD, Uh BS, Johnson RJ et al. Rehabilitation after anterior cruciate ligament reconstruction: a prospective, randomized, double-blind comparison of programs administered over 2 different time intervals. *Am J Sports Med* 2005;33(3):347-59.
89. Shelbourne KD, Klootwyk TE, DeCarlo MS. Rehabilitation program for anterior cruciate ligament reconstruction. *Sports Med Arthrosc Rev* 1997;5:77-82.
90. Grindem H, Granan LP, Risberg MA et al. How does a combined preoperative and postoperative rehabilitation programme influence the outcome of ACL reconstruction 2 years after surgery? A comparison between patients in the Delaware-Oslo ACL Cohort and the Norwegian National Knee Ligament Registry. *Br J Sports Med* 2014;49(6):385-9.
91. Failla MJ, Logerstedt DS, Grindem H et al. Does extended preoperative rehabilitation influence outcomes 2 years after ACL reconstruction? A comparative effectiveness study between the MOON and Delaware-Oslo ACL cohorts. *Am J Sports Med* 2016 Jul 14. pii: 0363546516652594. [Epub ahead of print]
92. Powers CM, Ho KY, Chen YJ et al. Patellofemoral joint stress during weight-bearing and non-weight-bearing quadriceps exercises. *J Orthop Sports Phys Ther* 2014;44(5):320-7.
93. Beynnon BD, Fleming BC, Johnson RJ et al. Anterior cruciate ligament strain behavior during rehabilitation exercises in vivo. *Am J Sports Med* 1995;23(1):24-34.
94. Culvenor AG, Collins NJ, Vicenzino B et al. Predictors and effects of patellofemoral pain following hamstring-tendon ACL reconstruction. *J Sci Med Sport* 2016;19(7):518-23.
95. Frobell RB. Change in cartilage thickness, posttraumatic bone marrow lesions and joint fluid volumes after acute ACL disruption: a two-year prospective MRI study of sixty-one subjects. *J Bone Joint Surg Am* 2011;93(12):1096-103.
96. Culvenor AG, Collins NJ, Guermazi A et al. Early knee osteoarthritis is evident one year following anterior cruciate ligament reconstruction: a magnetic resonance imaging evaluation. *Arthritis Rheumatol* 2015;67(4):946-55.
97. Eakin CL. Knee arthrofibrosis: prevention and management of a potentially devastating condition. *Phys Sportsmed* 2001;29(3):31-42.
98. Papalia R, Franceschi F, D'Adamio S et al. Hamstring tendon regeneration after harvest for anterior cruciate ligament reconstruction: a systematic review. *Arthroscopy* 2015;31(6):1169-83.
99. Suijkerbuijk MA, Reijman M, Lodewijks SJ et al. Hamstring tendon regeneration after harvesting: a systematic review. *Am J Sports Med* 2015;43(10):2591-8.
100. Petersen W, Taheri P, Forkel P et al. Return to play following ACL reconstruction: a systematic review about strength deficits. *Arch Orthop Trauma Surg* 2014;134(10):1417-28.
101. Howells BE, Ardern CL, Webster KE. Is postural control restored following anterior cruciate ligament reconstruction? A systematic review. *Knee Surg Sports Traumatol Arthrosc* 2011;19(7):1168-77.
102. Relph N, Herrington L, Tyson S. The effects of ACL injury on knee proprioception: a meta-analysis. *Physiotherapy* 2014;100(3):187-95.
103. Gokeler A, Benjaminse A, van Eck CF et al. Return of normal gait as an outcome measurement in ACL reconstructed

- patients. A systematic review. *Int J Sports Phys Ther* 2013;8(4):441-51.
104. Ardern CL, Taylor NF, Feller JA et al. Psychological responses matter in returning to preinjury level of sport after anterior cruciate ligament reconstruction surgery. *Am J Sports Med* 2013;41(7):1549-58.
 105. Ardern CL, Taylor NF, Feller JA et al. Fear of re-injury in people who have returned to sport following anterior cruciate ligament reconstruction surgery. *J Sci Med Sport* 2012;15(6):488-95.
 106. Everhart JS, Best TM, Flanigan DC. Psychological predictors of anterior cruciate ligament reconstruction outcomes: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2015;23(3):752-62.
 107. te Wierike SC, van der Sluis A, van den Akker-Scheek I et al. Psychosocial factors influencing the recovery of athletes with anterior cruciate ligament injury: a systematic review. *Scand J Med Sci Sports* 2013;23(5):527-40.
 108. Noyes FR, Barber SD, Mooar LA. A rationale for assessing sports activity levels and limitations in knee disorders. *Clin Orthop* 1989;246:238-49.
 109. Hefti F, Muller W, Jakob RP et al. Evaluation of knee ligament injuries with the IKDC form. *Knee Surg Sports Traumatol Arthrosc* 1993;1(3-4):226-34.
 110. Roos EM. Outcome after anterior cruciate ligament reconstruction—a comparison of patients' and surgeons' assessments. *Scand J Med Sci Sports* 2001;11(5):287-91.
 111. Magnusson RA, Verlage M, Flanigan DC et al. Patient-reported outcomes and their predictors at minimum 10 years after anterior cruciate ligament reconstruction: a systematic review of prospectively collected data. *Orthop J Sports Med* 2015;3(3):2325967115573706.
 112. Ageberg E, Forssblad M, Herbertsson P et al. Sex differences in patient-reported outcomes after anterior cruciate ligament reconstruction: data from the Swedish knee ligament register. *Am J Sports Med* 2010;38(7):1334-42.
 113. Paradowski PT, Bergman S, Sundén-Lundius A et al. Knee complaints vary with age and gender in the adult population. Population-based reference data for the Knee injury and Osteoarthritis Outcome Score (KOOS). *BMC Musculoskelet Disord* 2006;7:38.
 114. Filbay SR, Culvenor AG, Ackerman IN et al. Quality of life in anterior cruciate ligament-deficient individuals: a systematic review and meta-analysis. *Br J Sports Med* 2015;49(16):1033-41.
 115. Corry IS, Webb JM, Clingeleffer AJ et al. Arthroscopic reconstruction of the anterior cruciate ligament. A comparison of patellar tendon autograft and four-strand tendon autograft. *Am J Sports Med* 1999;27:444-54.
 116. Feller JA, Webster KE. A randomized comparison of patellar tendon and hamstring tendon anterior cruciate ligament reconstruction. *Am J Sports Med* 2003;31(4):564-73.
 117. Gobbi A, Ty B, Mahajan S et al. Quadrupled bone-semi-tendinosus anterior cruciate ligament reconstruction: a clinical investigation in a group of athletes. *Arthroscopy* 2003;19:691-9.
 118. Siegel MG, Barber-Westin SD. Arthroscopic-assisted outpatient anterior cruciate ligament reconstruction using the semitendinosus and gracilis tendons. *Arthroscopy* 1998;14:268-77.
 119. Biau DJMD, Tournoux CMD, Katsahian SMD et al. ACL reconstruction: a meta-analysis of functional scores. *Clin Orthop Relat Res* 2007;458:180-7.
 120. Walden M, Hägglund M, Magnusson H et al. Anterior cruciate ligament injury in elite football: a prospective three-cohort study. *Knee Surg Sports Traumatol Arthrosc* 2011;19(1):11-9.
 121. Shah VM, Andrews JR, Fleisig GS et al. Return to play after anterior cruciate ligament reconstruction in National Football League athletes. *Am J Sports Med* 2010;38(11):2233-9.
 122. Ardern CL, Taylor NF, Feller JA et al. Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *Br J Sports Med* 2014;48(21):1543-52.
 123. Czuppon S, Racette BA, Klein SE et al. Variables associated with return to sport following anterior cruciate ligament reconstruction: a systematic review. *Br J Sports Med* 2014;48(5):356-64.
 124. Nagelli CV, Hewett TE. Should return to sport be delayed until 2 years after anterior cruciate ligament reconstruction? Biological and functional considerations. *Sports Med* 2016 Jul 11. [Epub ahead of print]
 125. Myklebust G, Holm I, Maehlum S et al. Clinical, functional and radiologic outcome in team handball players 6 to 11 years after anterior cruciate ligament injury: a follow-up study. *Am J Sports Med* 2003;31(6):981-9.
 126. Roos H, Ornell M, Gardsell P et al. Soccer after anterior cruciate ligament injury: an incompatible combination? A national survey of incidence and risk factors and a 7 year follow up of 310 players. *Acta Orthop Scand* 1995;66:107-12.
 127. Kostogiannis I, Ageberg E, Neuman P et al. Activity level and subjective knee function 15 years after anterior cruciate ligament injury: a prospective, longitudinal study of nonreconstructed patients. *Am J Sports Med* 2007;35(7):1135-43.
 128. Fink C, Hoser C, Hackl W et al. Long-term outcome of operative or non-operative treatment of anterior cruciate ligament rupture: is sports activity a determining variable? *Int J Sports Med* 2001;22:304-9.
 129. von Porat A, Roos EM, Roos H. High prevalence of osteoarthritis 14 years after an anterior cruciate ligament tear in male soccer players: a study of radiographic and patient relevant outcomes. (A). *Ann Rheum Dis* 2004;63(3):269-73.
 130. Myklebust G, Bahr R. Return to play guidelines after anterior cruciate ligament surgery. *Br J Sports Med* 2005;39(3):127-31.
 131. Undheim MB, Cosgrave C, King E et al. Isokinetic muscle strength and readiness to return to sport following anterior

- cruciate ligament reconstruction: is there an association? A systematic review and a protocol recommendation. *Br J Sports Med* 2015;49(20):1305-10.
132. Paterno MV, Schmitt LC, Ford KR et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med* 2010;38:1968-78.
 133. Barber-Westin SD, Noyes FR. Objective criteria for return to athletics after anterior cruciate ligament reconstruction and subsequent reinjury rates: a systematic review. *Phys Sportsmed* 2011;39(3):100-10.
 134. Kyritsis P, Bahr R, Landreau P et al. Likelihood of ACL graft rupture: not meeting six clinical discharge criteria before return to sport is associated with a four times greater risk of rupture. *Br J Sports Med* 2016;50(15):946-51.
 135. Munro AG, Herrington LC. Between-session reliability of four hop tests and the agility T test. *J Strength Cond Res* 2011;25:1470-7.
 136. Crawford SN, Waterman BR, Lubowitz JH. Long-term failure of anterior cruciate ligament reconstruction. *Arthroscopy* 2013;29(9):1566-71.
 137. Swärd P, Kostogiannis I, Roos H. Risk factors for a contralateral anterior cruciate ligament injury. *Knee Surg Sports Traumatol Arthrosc* 2010;18(3):277-91.
 138. Salmon L, Russell V, Musgrove T et al. Incidence and risk factors for graft rupture and contralateral rupture after anterior cruciate ligament reconstruction. *Arthroscopy* 2005;21(8):948-57.
 139. Webster KE, Feller JA. Exploring the high reinjury rate in younger patients undergoing anterior cruciate ligament reconstruction. *Am J Sports Med* 2016 Jul 7. pii: 0363546516651845. [Epub ahead of print]
 140. Grassi A, Ardern CL, Marcheggiani Muccioli GM et al. Does revision ACL reconstruction measure up to primary surgery? A meta-analysis comparing patient-reported and clinician-reported outcomes and radiographic results. *Br J Sports Med* 2016;50(12):716-24.
 141. Andriolo L, Filardo G, Kon E et al. Revision anterior cruciate ligament reconstruction: clinical outcome and evidence for return to sport. *Knee Surg Sports Traumatol Arthrosc* 2015;23(10):2825-45.
 142. Lohmander LS, Englund PM, Dahl LL et al. The long-term consequence of anterior cruciate ligament and meniscus injuries. *Am J Sports Med* 2007;35(10):1756-69.
 143. Carey J, Dunn W, Dahm D, et al. A systematic review of anterior cruciate ligament reconstruction with autograft compared with allograft. *J Bone Joint Surg Am* 2009;91: 2242-50.
 144. Frobell RB, Roos HP, Roos EM et al. Treatment for acute anterior cruciate ligament tear: five-year outcome of randomised trial. *BMJ (Online)* 2013;346:f232.
 145. Culvenor AG, Collins NJ, Guermazi A et al. Early knee osteoarthritis is evident one year following anterior cruciate ligament reconstruction: a magnetic resonance imaging evaluation. *Arth Rheumatol* 2015;67(4):946-55.
 146. Neuman P, Englund M, Kostogiannis I et al. Prevalence of tibiofemoral osteoarthritis 15 years after nonoperative treatment of anterior cruciate ligament injury: a prospective cohort study. *Am J Sports Med* 2008;36:1717-25.
 147. Grindem H, Eitzen I, Moksnes H et al. A pair-matched comparison of return to pivoting sports at 1 year in anterior cruciate ligament-injured patients after a non-operative versus operative treatment course. *Am J Sports Med* 2012;40:2509-16.
 148. Frobell RB, Le Graverand MP, Buck R et al. The acutely ACL injured knee assessed by MRI: changes in joint fluid, bone marrow lesions and cartilage during the first year. *Osteoarth Cart* 2009;17(2):161-7.
 149. Røtterud JH, Sivertsen EA, Forssblad M et al. Effect of meniscal and focal cartilage lesions on patient-reported outcome after anterior cruciate ligament reconstruction: a nationwide cohort study from Norway and Sweden of 8476 patients with 2-year follow-up. *Am J Sports Med* 2013;41(3):535-43.
 150. Spindler KP, Schils JP, Bergfeld JA et al. Prospective study of osseous, articular, and meniscal lesions in recent anterior cruciate ligament tears by magnetic resonance imaging and arthroscopy. *Am J Sports Med* 1993;21(4):551-7.
 151. Buckwalter JA. Articular cartilage injuries. *Clin Orthop Relat Res* 2002;402:21-37.
 152. Martin JA, Brown T, Heiner A et al. Post-traumatic osteoarthritis: the role of accelerated chondrocyte senescence. *Biorheology* 2004;41:479-91.
 153. Andriacchi TP, Mundermann A, Smith RL et al. A framework for the *in vivo* pathomechanics of osteoarthritis at the knee. *Ann Biomed Eng* 2004;32(3):447-57.
 154. Chaudhari AMW, Briant PL, Beville SL et al. Knee kinematics, cartilage morphology and osteoarthritis after ACL injury. *Med Sci Sports Exerc* 2008;40(2):215-22.
 155. Hart HF, Culvenor AG, Collins NJ et al. Knee kinematics and joint moments during gait following anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Br J Sports Med* 2016;50(10):597-612.
 156. Keays SL, Bullock-Saxton JE, Keays AC et al. A 6-year follow-up of the effect of graft site on strength, stability, range of motion, function and joint degeneration after anterior cruciate ligament reconstruction. *Am J Sports Med* 2007;35(5):729-39.
 157. Hewett TE, Myer GD, Ford KR et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes. *Am J Sports Med* 2005;33(4):492-501.
 158. Hart HF, Collins NJ, Ackland DC et al. Gait characteristics of people with lateral knee osteoarthritis after ACL reconstruction. *Med Sci Sports Exerc* 2015;47(11):2406-15.
 159. Sward P, Kostogiannis I, Neuman P et al. Differences in the radiological characteristics between post-traumatic and non-traumatic knee osteoarthritis. *Scand J Med Sci Sports* 2010;20:731-9.

160. Hewett TE, Stroupe AL, Nance TA et al. Plyometric training in female athletes: decreased impact forces and increased hamstring torques. *Am J Sports Med* 1996;24(6):765-73.
161. Culvenor AG, Cook JL, Collins NJ et al. Is patellofemoral joint osteoarthritis an under-recognised outcome of anterior cruciate ligament reconstruction? A narrative literature review. *Br J Sports Med* 2013;47(2):66-70.
162. Culvenor AG, Crossley KM. Patellofemoral osteoarthritis: are we missing an important source of symptoms after anterior cruciate ligament reconstruction? *J Orth Sports Physl Ther* 2016;46(4):232-4.
163. Culvenor AG, Collins NJ, Guermazi A et al. Early patellofemoral osteoarthritis features one year after anterior cruciate ligament reconstruction: symptoms and quality of life at three years. *Arth Care Res* 2016;68(6):784-92.
164. Culvenor AG, Perraton L, Guermazi A et al. Knee kinematics and kinetics are associated with early patellofemoral osteoarthritis following anterior cruciate ligament reconstruction. *Osteoarth Cart* 2016;24(9):1548-53.
165. Culvenor AG, Schache AG, Vicenzino B et al. Are knee biomechanics different in those with and without patellofemoral osteoarthritis after anterior cruciate ligament reconstruction? *Arth Care Res* 2014;66(10):1566-70.
166. Culvenor AG, Collins NJ, Vicenzino B et al. Predictors and effects of patellofemoral pain following hamstring-tendon ACL reconstruction. *J Sci Med Sport* 2016;19(7):518-23.
167. Barber-Westin SD, Noyes F, Andrews M. A rigorous comparison between sexes of results and complications after anterior cruciate reconstruction. *Am J Sports Med* 1997;25:514-25.
168. Noojin FK, Barret GR, Hartzog CW et al. Clinical comparison of intra-articular anterior cruciate ligament reconstruction using autogenous semitendinosus and gracilis tendons in men versus women. *Am J Sports Med* 2000;28:783-9.
169. Ferrari JD, Bach BRJ, Bush-Joseph CA et al. Anterior cruciate ligament reconstruction in men and women: an outcome analysis comparing gender. *Arthroscopy* 2001;17:588-96.
170. Gobbi A, Domzalski M, Pascual J. Comparison of anterior cruciate ligament reconstruction in male and female athletes using the patellar tendon and hamstring autografts. *Knee Surg Sports Traumatol Arthrosc* 2004;12:534-9.
171. Salmon LJ, Refshauge KM, Russell VJ et al. Gender differences in outcome after anterior cruciate ligament reconstruction with hamstring tendon autograft. *Am J Sports Med* 2006;34(4):621-9.
172. Ryan J, Magnussen RA, Cox CL et al. ACL reconstruction: do outcomes differ by sex? A systematic review. *J Bone Joint Surg Am* 2014;96(6):507-12.
173. Kaeding CC, Aros B, Pedroza A et al. Allograft versus autograft anterior cruciate ligament reconstruction: predictors of failure from a MOON prospective longitudinal cohort. *Sports Health* 2011;3:73-81.
174. Shelbourne KD, Gray T, Haro M. Incidence of subsequent injury to either knee within 5 years after anterior cruciate ligament reconstruction with patellar tendon autograft. *Am J Sports Med* 2009;37(2):246-51.
175. Andernord D, Desai N, Björnsson H et al. Patient predictors of early revision surgery after anterior cruciate ligament reconstruction a cohort study of 16 930 patients with 2-year follow-up. *Am J Sports Med* 2014;43:121-7.
176. Mall NA, Frank RM, Saltzman BM et al. Results after anterior cruciate ligament reconstruction in patients older than 40 years: how do they compare with younger patients? A systematic review and comparison with younger populations. *Sports Health* 2016;8(2):177-81.
177. Pujol N, Colombet P, Cucurulo T et al. Natural history of partial anterior cruciate ligament tears: a systematic literature review. *Orthop Traumatol Surg Res* 2012;98(8 Suppl):S160-4.
178. DeFranco MJ, Bach BRJ. A comprehensive review of partial anterior cruciate ligament tears. *J Bone Joint Surg Am* 2009;91(1):198-208.
179. Kannus P, Jarvinen M, Johnson R et al. Function of the quadriceps and hamstrings muscles in knees with chronic partial deficiency of the anterior cruciate ligament. Isometric and isokinetic evaluation. *Am J Sports Med* 1992;20:162-8.
180. Ahmad CS, Cohen ZA, Levine WN et al. Codominance of the individual posterior cruciate ligament bundles. An analysis of bundle lengths and orientation. *Am J Sports Med* 2003;31(2):221-5.
181. LaPrade CM, Civitarese DM, Rasmussen MT et al. Emerging updates on the posterior cruciate ligament: a review of the current literature. *Am J Sports Med* 2015;43(12):3077-92.
182. Becker EH, Watson JD, Dreese JC. Investigation of multiligamentous knee injury patterns with associated injuries presenting at a level I trauma center. *J Orthop Trauma* 2013;27(4):226-31.
183. Fanelli GC, Edson CJ. Posterior cruciate ligament injuries in trauma patients, part II. *Arthroscopy* 1995;11(5):526-9.
184. Hamada M, Shino K, Mitsuoka T et al. Chondral injury associated with acute isolated posterior cruciate ligament injury. *Arthroscopy* 2000;16:59-63.
185. Bergfeld JA, McAllister DR, Parker RD et al. The effects of tibial rotation on posterior translation in knees in which the posterior cruciate ligament has been cut. *J Bone Joint Surg Am* 2001;83-a(9):1339-43.
186. Hewett TE, Noyes FR, Lee MD. Diagnosis of complete and partial posterior cruciate ligament ruptures. Stress radiography compared with KT-1000 arthrometer and posterior drawer testing. *Am J Sports Med* 1997;25:648-55.
187. Sekiya JK, Whiddon DR, Zehms CT et al. A clinically relevant assessment of posterior cruciate ligament and posterolateral corner injuries: evaluation of isolated and combined deficiency. *J Bone Joint Surg Am* 2008;90(8):1621-7.
188. Gross ML, Groiver JS, Bassett LW et al. Magnetic resonance imaging of the posterior cruciate ligament: clinical use to improve diagnostic accuracy. *Am J Sports Med* 1992;20:732-7.
189. Esmaili Jah AA, Keyhani S, Zarei R et al. Accuracy of MRI in comparison with clinical and arthroscopic findings in

- ligamentous and meniscal injuries of the knee. *Acta Orthop Belg* 2005;71(2):189-96.
190. Servant CT, Ramos JP, Thomas NP. The accuracy of magnetic resonance imaging in diagnosing chronic posterior cruciate ligament injury. *Knee* 2004;11(4):265-70.
 191. Rodriguez WJ, Vinson EN, Helms CA et al. MRI appearance of posterior cruciate ligament tears. *AJR Am J Roentgenol* 2008;191(4):1031.
 192. Mair SD, Schlegel TF, Gill TJ et al. Incidence and location of bone bruises after acute posterior cruciate ligament injury. *Am J Sports Med* 2004;32(7):1681-7.
 193. LaPrade RF, Smith SD, Wilson KJ et al. Quantification of functional brace forces for posterior cruciate ligament injuries on the knee joint: an in vivo investigation. *Knee Surg Sports Traumatol Arthrosc* 2015;23(10):3070-6.
 194. Shelbourne KD, Davis TJ, Patel DV. The natural history of acute, isolated, nonoperatively treated posterior cruciate ligament injuries. A prospective study. *Am J Sports Med* 1999;27(3):276-83.
 195. Shelbourne KD, Muthukaruppan Y. Subjective results of nonoperatively treated, acute, isolated posterior cruciate ligament injuries. *Arthroscopy* 2005;21(4):457-61.
 196. Parolie JM, Bergfeld JA. Long-term results of nonoperative treatment of isolated posterior cruciate ligament injuries in the athlete. *Am J Sports Med* 1986;14(1):35-8.
 197. Montgomery SR, Johnson JS, McAllister DR et al. Surgical management of PCL injuries: indications, techniques and outcomes. *Curr Rev Musculoskel Med* 2013;6(2):115-23.
 198. Boynton MD, Tietjens BR. Long-term followup of the untreated isolated posterior cruciate ligament-deficient knee. *Am J Sports Med* 1996;24(3):306-10.
 199. Dandy DJ, Pusey RJ (1982) The long-term results of unrepaired tears of the posterior cruciate ligament. *J Bone Joint Surg [Br]* 64-B(1):92-94.
 200. Shelbourne KD, Clark M, Grey T. Minimum 10-year follow-up of patients after an acute, isolated posterior cruciate ligament injury treated nonoperatively. *Am J Sports Med* 2013;41(7):1526-33.
 201. Cooper DE, Stewart D. Posterior cruciate ligament reconstruction using single-bundle patella tendon graft with tibial inlay fixation: 2- to 10-year follow-up. *Am J Sports Med* 2004;32(2):346-60.
 202. Wijdicks CA, Kennedy NI, Goldsmith MT et al. Kinematic analysis of the posterior cruciate ligament, part 2: a comparison of anatomic single- versus double-bundle reconstruction. *Am J Sports Med* 2013;41(12):2839-48.
 203. Hermans S, Corten K, Bellemans J. Long-term results of isolated anterolateral bundle reconstructions of the posterior cruciate ligament: a 6- to 12-year follow-up study. *Am J Sports Med* 2009;37(8):1499-507.
 204. Jackson WF, van der Tempel WM, Salmon LJ et al. Endoscopically-assisted single-bundle posterior cruciate ligament reconstruction: results at minimum ten-year follow-up. *J Bone Joint Surg Br* 2008;90(10):1328-33.
 205. Hammoud S, Reinhardt KR, Marx RG. Outcomes of posterior cruciate ligament treatment: a review of the evidence. *Sports Med Arthrosc* 2010;18(4):280-91.
 206. Pierce CM, O'Brien L, Gryphon LW et al. Posterior cruciate ligament tears: functional and postoperative rehabilitation. *Knee Surg Sports Traumatol Arthrosc* 2013;21(5):1071-84.
 207. Bushnell BD, Bitting SS, Crain JM et al. Treatment of magnetic resonance imaging-documented isolated grade III lateral collateral ligament injuries in National Football League athletes. *Am J Sports Med* 2010;38(1):86-91.
 208. LaPrade RF, Wentorf FA, Fritts H et al. A prospective magnetic resonance imaging study of the incidence of posterolateral and multiple ligament injuries in acute knee injuries presenting with a hemarthrosis. *Arthroscopy* 2007;23(12):1341-7.
 209. Chahla J, Moatshe G, Dean CS et al. Posterolateral corner of the knee: current concepts. *Arch Bone Jt Surg* 2016;4(2):97-103.
 210. LaPrade RF, Ly TV, Wentorf FA et al. The posterolateral attachments of the knee: a qualitative and quantitative morphologic analysis of the fibular collateral ligament, popliteus tendon, popliteofibular ligament and lateral gastrocnemius tendon. *Am J Sports Med* 2003;31(6):654-60.
 211. Cooper JM, McAndrews PT, LaPrade RF. Posterolateral corner injuries of the knee: anatomy, diagnosis and treatment. *Sports Med Arthrosc* 2006;14(4):213-20.
 212. LaPrade RF, Terry GC. Injuries to the posterolateral aspect of the knee. Association of anatomic injury patterns with clinical instability. *Am J Sports Med* 1997;25(4):433-8.
 213. Crespo B, James EW, Metsavaht L et al. Injuries to posterolateral corner of the knee: a comprehensive review from anatomy to surgical treatment. *Rev Bras Ortop* 2015;50(4):363-70.
 214. Kannus P. Nonoperative treatment of grade II and III sprains of the lateral ligament compartment of the knee. *Am J Sports Med* 1989;17(1):83-8.
 215. Geeslin AG, LaPrade RF. Outcomes of treatment of acute grade-III isolated and combined posterolateral knee injuries: a prospective case series and surgical technique. *J Bone Joint Surg Am* 2011;93(18):1672-83.
 216. Harner CD, Vogrin TM, Hoher J et al. Biomechanical analysis of a posterior cruciate ligament reconstruction. Deficiency of the posterolateral structures as a cause of graft failure. *Am J Sports Med* 2000;28(1):32-9.
 217. Geeslin AG, Moulton SG, LaPrade RF. A systematic review of the outcomes of posterolateral corner knee injuries, part 1: surgical treatment of acute injuries. *Am J Sports Med* 2016;44(5):1336-42.
 218. Stannard JP, Brown SL, Farris RC et al. The posterolateral corner of the knee: repair versus reconstruction. *Am J Sports Med* 2005;33(6):881-8.
 219. Levy BA, Dajani KA, Morgan JA et al. Repair versus reconstruction of the fibular collateral ligament and

- posterolateral corner in the multiligament-injured knee. *Am J Sports Med* 2010;38(4):804-9.
220. Moulton SG, Geeslin AG, LaPrade RF. A systematic review of the outcomes of posterolateral corner knee injuries, part 2: surgical treatment of chronic injuries. *Am J Sports Med* 2016;44(6):1616-23.
 221. Herman B, Litchfield R, Getgood A. Role of osteotomy in posterolateral instability of the knee. *J Knee Surg* 2015;28(6):441-9.
 222. Flanigan DC, Harris JD, Trinh TQ et al. Prevalence of chondral defects in athletes' knees: systematic review. *Med Sci Sports Exerc* 2010;42(10):1795-801.
 223. Harris JD, Brophy RH, Siston RA et al. Treatment of chondral defects in the athlete's knee. *Arthroscopy* 2010;26(6):841-52.
 224. Behery O, Siston RA, Harris JD et al. Treatment of cartilage defects of the knee: expanding on the existing algorithm. *Clin J Sport Med* 2014;24(1):21-32.
 225. Scolaro J, Bernstein J, Ahn J. In brief: patellar fractures. *Clin Orthop Relat Res* 2011;469(4):1213-5.
 226. Duthon VB. Acute traumatic patellar dislocation. *Orthop Traumatol Surg Res* 2015;101(1 Suppl):S59-67.
 227. Desio SM, Burks RT, Bachus KN. Soft tissue restraints to lateral patellar translation in the human knee. *Am J Sports Med* 1998;26(1):59-65.
 228. O'Reilly MA, O'Reilly PM, Bell J. Sonographic appearances of medialretinacular complex injury in transient patellar dislocation. *Clin Radiol* 2003;58(8):636-41.
 229. Sillanpaa P, Mattila VM, Iivonen T et al. Incidence and risk factors of acute traumatic primary patellar dislocation. *Med Sci Sports Exerc* 2008;40(4):606-11.
 230. Nomura E, Horiuchi Y, Inoue M. Correlation of MR imaging findings and open exploration of medial patellofemoral ligament injuries in acute patellar dislocations. *Knee* 2002;9(2):139-43.
 231. Stefancin JJ, Parker RD. First-time traumatic patellar dislocation: a systematic review. *Clin Orthop Relat Res* 2007;455:93-101.
 232. Hing CB, Smith TO, Donell S et al. Surgical versus non-surgical interventions for treating patellar dislocation. *Cochrane Database Syst Rev* 2011;11:CD008106.
 233. Dye SF, Vaupel GL, Dye CS. Conscious neurosensory mapping of the internal structures of the human knee without intraarticular anesthesia. *Am J Sports Med* 1998;26(6):773-7.
 3. van Middelkoop M, van Linschoten R, Berger M et al. Knee complaints seen in general practice: active sport participants versus non-sport participants. *BMC Musculoskel Dis* 2008;9(1):36.
 4. Rio E, Kidgell D, Moseley GL et al. Elevated corticospinal excitability in patellar tendinopathy compared with other anterior knee pain or no pain. *Scand J Med Sci Sports* 2016;26(9):1072-9.
 5. Torry MR, Decker MJ, Viola RW et al. Intra-articular knee joint effusion induces quadriceps avoidance gait patterns. *Clinical Biomech* 2000;15(3):147-59.
 6. Hodges PW, Mellor R, Crossley K et al. Pain induced by injection of hypertonic saline into the infrapatellar fat pad and effect on coordination of the quadriceps muscles. *Arthritis Care Res* 2009;61(1):70-7.
 7. Elias JJ, Kilambi S, Cosgarea AJ. Computational assessment of the influence of vastus medialis obliquus function on patellofemoral pressures: model evaluation. *J Biomech* 2010;43(4):612-17.
 8. Purdam CR, Jonsson P, Alfredson H et al. A pilot study of the eccentric decline squat in the management of painful chronic patellar tendinopathy. *Br J Sports Med* 2004;38(4):395-7.
 9. Cook C, Mabry L, Reiman MP et al. Best tests/clinical findings for screening and diagnosis of patellofemoral pain syndrome: a systematic review. *Physiotherapy* 2012;98(2):93-100.
 10. Cook C, Hegedus E, Hawkins R et al. Diagnostic accuracy and association to disability of clinical test findings associated with patellofemoral pain syndrome. *Physiother Canada* 2010;62(1):17-24.
 11. Naslund J, Naslund UB, Odenbring S et al. Comparison of symptoms and clinical findings in subgroups of individuals with patellofemoral pain. *Physiother Theory Prac* 2006;22(3):105-18.
 12. Sweitzer BA, Cook C, Steadman JR et al. The inter-rater reliability and diagnostic accuracy of patellar mobility tests in patients with anterior knee pain. *Phys Sportsmed* 2010;38(3):90-6.
 13. Nunes GS, Stapait EL, Kirsten MH et al. Clinical test for diagnosis of patellofemoral pain syndrome: systematic review with meta-analysis. *Phys Ther Sport* 2013;14(1):54-9.
 14. Chatman AB, Hyams SP, Neel JM et al. The patient-specific functional scale—measurement properties in patients with knee dysfunction. *Phys Ther* 1997;77(8):820-9.
 15. Kujala UM, Jaakkola LH, Koskinen SK et al. Scoring of patellofemoral disorders. *Arthroscopy* 1993;9(2):159-163.
 16. Green A, Liles C, Rushton A et al. Measurement properties of patient-reported outcome measures (PROMS) in patellofemoral pain syndrome: a systematic review. *Manual Ther* 2014;19(6):517-26.
 17. Crossley KM, Bennell KL, Green S et al. Analysis of outcome measures for persons with patellofemoral pain: which are reliable and valid? *Arch Phys Med Rehabil* 2004;85(5):815-22.
 18. Elias DA, White LM. Imaging of patellofemoral disorders. *Clinical Radiol* 2004;59(7):543-57.
 19. Crossley KM, Callaghan M, van Linschoten R. Patellofemoral pain: practice pointer. *BMJ* 2015;351:h3939.
 20. Warden SJ, Kiss ZS, Malara FA et al. Comparative accuracy of magnetic resonance imaging and ultrasonography in confirming

CHAPTER 36

1. Taunton JE, Ryan MB, Clement DB et al. A retrospective case-control analysis of 2002 running injuries. *Br J Sports Med* 2002;36:95-101.
2. Baquie P, Brukner P. Injuries presenting to an Australian Sports Medicine Centre: a 12-month study. *Clin J Sport Med* 1997;7(1):28-31.

- clinically diagnosed patellar tendinopathy. *Am J Sports Med* 2007;35(3):427-36.
21. Cook JL, Khan KM, Kiss ZS et al. Reproducibility and clinical utility of tendon palpation to detect patellar tendinopathy in young basketball players. *Br J Sports Med* 2001;35:65-9.
 22. Cook JL, Khan KM, Kiss ZS et al. Asymptomatic hypoechoic regions on patellar tendon ultrasound: a 4-year clinical and ultrasound followup of 46 tendons. *Scand J Med Sci Sports* 2001;11:321-7.
 23. Shalaby M, Almekinders LC. Patellar tendinitis: the significance of magnetic resonance imaging findings. *Am J Sports Med* 1999;27(3):345-9.
 24. Crossley KM, Stefanik J, Selfe J et al. Patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester Part 1: terminology, definitions, clinical examination, natural history, patellofemoral osteoarthritis and patient-reported outcome measures. *Br J Sports Med* 2016;50(14):839-43.
 25. Dye SF, Vaupel GL, Dye CS. Conscious neurosensory mapping of the internal structures of the human knee without intraarticular anaesthesia. *Am J Sports Med* 1998;26(6):773-7.
 26. Insall J, Falvo KA, Wise DW. Chondromalacia patellae. A prospective study. *J Bone Joint Surg Am* 1976;58(1):1-8.
 27. Fulkerson JP. Diagnosis and treatment of patients with patellofemoral pain. *Am J Sports Med* 2002;30(3):447-56.
 28. Fulkerson JP, Gossling HR. Anatomy of the knee joint lateral retinaculum. *Clin Orthop Rel Res* 1980;153:183-8.
 29. Fulkerson JP, Tennant R, Jaivin JS et al. Histologic evidence of retinacular nerve injury associated with patellofemoral malalignment. *Clin Orthop Rel Res* 1985;197:196-205.
 30. Sanchis-Alfonso V, Rosello-Sastre E, Monteagudo-Castro C et al. Quantitative analysis of nerve changes in the lateral retinaculum in patients with isolated symptomatic patellofemoral malalignment. *Am J Sports Med* 1998;26(5):703-9.
 31. Sanchis-Alfonso V, Rosellà-Sastre E. Immunohistochemical analysis for neural markers of the lateral retinaculum in patients with isolated symptomatic patellofemoral malalignment. *Am J Sports Med* 2000;28(5):725-31.
 32. Witoński D, Wągrowska-Danielewicz M. Distribution of substance-P nerve fibers in the knee joint in patients with anterior knee pain syndrome. A preliminary report. *Knee Surg Sports Traumatol Arthrosc* 1999;7(3):177-83.
 33. Duri ZAA, Aichroth PM, Dowd G. The fat pad. *Am J Knee Surg* 1996;9(2):55-66.
 34. Tsirbas A, Paterson RS, Keene GCR. Fat pad impingement: a missed cause of patello femoral pain? *Aus J Sci Med Sport* 1991;23:24-6.
 35. Fulkerson JP. *Disorders of the patellofemoral joint*. 3rd ed. Baltimore: Williams & Wilkins, 1997.
 36. Matthews LS, Sonstegard DA, Henke JA. Load bearing characteristics of the patello-femoral joint. *Acta Orthop* 1977;48(5):511-16.
 37. Dye SF, Vaupel GL. The pathophysiology of patellofemoral pain. *Sports Med Arthrosc Rev* 1994;2:203-10.
 38. Rathleff MS, Roos EM, Olesen JL et al. Lower mechanical pressure pain thresholds in female adolescents with patellofemoral pain syndrome. *J Orth Sports Phys Ther* 2013;43(6): 414-421.
 39. Graven-Nielsen T, Arendt-Nielsen L. Assessment of mechanisms in localized and widespread musculoskeletal pain. *Nat Rev Rheumatol* 2010;6(10):599-606.
 40. Rathleff MS, Roos EM, Olesen JL et al. Self-reported recovery is associated with improvement in localised hyperalgesia among adolescent females with patellofemoral pain—results from a cluster randomised trial. *Clin J Pain* 2016;32(5):428-34.
 41. Nijs J, Kosek E, Vanoosterwijck J et al. Dysfunctional endogenous analgesia during exercise in patients with chronic pain: to exercise or not to exercise? *Pain Phys* 2012;15 (3 Suppl):ES205-13.
 42. Kornaat P, Bloem J, Ceulemans R et al. Osteoarthritis of the knee: association between clinical features and MR imaging findings. *Radiology* 2006;239(3):811-17.
 43. Duncan R, Peat G, Thomas E et al. Incidence, progression and sequence of development of radiographic knee osteoarthritis in a symptomatic population. *Ann Rheum Dis* 2011;70(11):1944-8.
 44. Hinman RS, Lentzos J, Vicenzino B et al. Patellofemoral osteoarthritis is common in middle-aged people with chronic patellofemoral pain. *Arthr Care Res* 2014;16(8):1252-7.
 45. Crossley KM. Is patellofemoral osteoarthritis a common sequela of patellofemoral pain? *Br J Sports Med* 2014;48(6):409-10.
 46. Utting MR, Davies G, Newman JH. Is anterior knee pain a predisposing factor to patellofemoral osteoarthritis? *Knee* 2005;12(5):362-5.
 47. Schiphof D, van Middelkoop M, de Klerk BM et al. Crepitus is a first indication of patellofemoral osteoarthritis (and not tibiofemoral osteoarthritis). *Osteoarthritis Cart* 2014;22(5):631-8.
 48. Culvenor AG, Cook JL, Collins NJ et al. Is patellofemoral joint osteoarthritis an under-recognised outcome of anterior cruciate ligament reconstruction? A narrative literature review. *Br J Sports Med* 2013;47(2):66-70.
 49. Frobell RB, Roos HP, Roos EM et al. Treatment for acute anterior cruciate ligament tear: five year outcome of randomised trial. *BMJ* 2013;346:f232.
 50. Culvenor AG, Collins NJ, Guermazi A et al. Early knee osteoarthritis is evident one year following anterior cruciate ligament reconstruction: a magnetic resonance imaging evaluation. *Arthritis Rheumatol* 2015; 67(4):946-55.
 51. Culvenor AG, Lai CCH, Gabbe BJ et al. Patellofemoral osteoarthritis is prevalent and associated with worse symptoms and function after hamstring tendon autograft ACL reconstruction. *Br J Sports Med* 2014;48(6):435-9.
 52. Li RT, Lorenz S, Xu Y et al. Predictors of radiographic knee osteoarthritis after anterior cruciate ligament reconstruction. *Am J Sports Med* 2011;39(12):2595-603.
 53. Englund M, Lohmander LS. Patellofemoral osteoarthritis coexistent with tibiofemoral osteoarthritis in a meniscectomy population. *Ann Rheum Dis* 2005;64:1721-6.

54. Myer GD, Ford KR, Di Stasi SL et al. High knee abduction moments are common risk factors for patellofemoral pain (PFP) and anterior cruciate ligament injury (ACL) in girls: is PFP itself a predictor for subsequent ACL injury? *Br J Sports Med* 2014;49:118-22.
55. Maenpaa H, Lehto MUK. Patellofemoral osteoarthritis after patellar dislocation. *Clin Orthop Rel Res* 1997;339:156-62.
56. Nomura E, Inoue M. Cartilage lesions of the patella in recurrent patellar dislocation. *Am J Sports Med* 2004;32(2):498-502.
57. Wluka AE, Ding C, Jones G et al. The clinical correlates of articular cartilage defects in symptomatic knee osteoarthritis: a prospective study. *Rheumatology* 2005;44:1311-16.
58. Schiphof D, van Middelkoop M, de Klerk BM et al. Crepitus is a first indication of patellofemoral osteoarthritis (and not of tibiofemoral osteoarthritis). *Osteoarthritis Cart* 2014;22(5):631-8.
59. Hunter DJ, Arden N, Conaghan PG et al. Definition of osteoarthritis on MRI: results of a Delphi exercise. *Osteoarthritis Cart* 2011;19:963-9.
60. Powers CM. The influence of altered lower-extremity kinematics on patellofemoral joint dysfunction: a theoretical perspective. *J Orth Sports Phys Ther* 2003;33(11):639-46.
61. Grelsamer RP, Weinstein CH. Applied biomechanics of the patella. *Clin Orthop Rel Res* 2001;389:9-14.
62. Rathleff MS, Rathleff CR, Crossley KM et al. Is hip strength a risk factor for patellofemoral pain? A systematic review and meta-analysis. *Br J Sports Med* 2014;48(14):1088.
63. Nakagawa TH, Moriya ÉTU, Maciel CD et al. Trunk, pelvis, hip and knee kinematics, hip strength and gluteal muscle activation during a single-leg squat in males and females with and without patellofemoral pain syndrome. *J Orth Sports Phys Ther* 2012;42(6):491-501.
64. Noehren B, Pohl MB, Sanchez Z et al. Proximal and distal kinematics in female runners with patellofemoral pain. *Clinical Biomech* 2012;27(4):366-71.
65. Barton CJ, Levinger P, Menz HB et al. Kinematic gait characteristics associated with patellofemoral pain syndrome: a systematic review. *Gait Posture* 2009;30(4):405-16.
66. Boling MC, Padua DA, Marshall SW et al. A prospective investigation of biomechanical risk factors for patellofemoral pain syndrome. *Am J Sports Med* 2009;37(11):2108-16.
67. Barton CJ, Levinger P, Crossley KM et al. The relationship between rearfoot, tibial and hip kinematics in individuals with patellofemoral pain syndrome. *Clinical Biomech* 2012;27(7):702-5.
68. Dierks TA, Manal KT, Hamill J et al. Proximal and distal influences on hip and knee kinematics in runners with patellofemoral pain during a prolonged run. *J Orth Sports Phys Ther* 2008;38(8):448-56.
69. Myer GD, Ford KR, Barber Foss KD et al. The incidence and potential pathomechanics of patellofemoral pain in female athletes. *Clin Biomech* 2010;25(7):700-7.
70. Willson JD, Davis IS. Lower extremity mechanics of females with and without patellofemoral pain across activities with progressively greater task demands. *Clin Biomech* 2008;23(2):203-11.
71. Teng HL, Powers CM. Sagittal plane trunk posture influences patellofemoral joint stress during running. *J Orth Sports Phys Ther* 2014;44(10):785-92.
72. Dos Reis AC, Ferrari Correa OC, Bley AS et al. Kinematic and kinetic analysis of the single-leg triple hop test in women with and without patellofemoral pain. *J Orth Sports Phys Ther* 2015;45(10):799-807.
73. Nakagawa TH, Maciel CD, Serrão FV. Trunk biomechanics and its association with hip and knee kinematics in patients with and without patellofemoral pain. *Man Ther* 2015;20(1):189-93.
74. McKenzie K, Galea V, Wessel J et al. Lower extremity kinematics of females with patellofemoral pain syndrome while stair stepping. *J Orth Sports Phys Ther* 2010;40(10):625-32.
75. Souza RB, Draper CE, Fredericson M et al. Femur rotation and patellofemoral joint kinematics: a weight-bearing magnetic resonance imaging analysis. *J Orth Sports Phys Ther* 2010;40(5):277-85.
76. Dowling GJ, Murley GS, Munteanu SE et al. Dynamic foot function as a risk factor for lower limb overuse injury: a systematic review. *J Foot Ankle Res* 2014;7(1):53.
77. Neal BS, Griffiths IB, Dowling GJ et al. Foot posture as a risk factor for lower limb overuse injury: a systematic review and meta-analysis. *J Foot Ankle Res* 2014;7(1):55.
78. McPoil TG, Warren M, Vicenzino B et al. Variations in foot posture and mobility between individuals with patellofemoral pain and those in a control group. *J Am Pod Med Assoc* 2011;101(4):289-96.
79. Heino Brechter J, Powers CM. Patellofemoral joint stress during stair ascent and descent in persons with and without patellofemoral pain. *Gait Post* 2002;16:115-23.
80. Heino Brechter J, Powers CM. Patellofemoral stress during walking in persons with and without patellofemoral pain. *Med Sci Sports Exerc* 2002;34(10):1582-93.
81. Crossley KM, Cowan SM, Bennell KL et al. Knee flexion during stair ambulation is altered in individuals with patellofemoral pain. *J Orthop Res* 2004;22(2):267-74.
82. Bley AS, Correa JCF, Reis ACD et al. Propulsion phase of the single leg triple hop test in women with patellofemoral pain syndrome: a biomechanical study. *PLoS ONE* 2014;9(5):e97606.
83. Pollard CD, Sigward SM, Powers CM. Limited hip and knee flexion during landing is associated with increased frontal plane knee motion and moments. *Clin Biomech* 2010;25(2):142-6.
84. Amin S, Baker K, Niu J et al. Quadriceps strength and the risk of cartilage loss and symptom progression in knee osteoarthritis. *Arth Rheum* 2009;60(1):189-98.
85. Watson CJ, Propps M, Galt W et al. Reliability of McConnell's classification of patellar orientation in symptomatic and asymptomatic subjects. *J Orth Sports Phys Ther* 1999;29(7):386-93.
86. McConnell J. Reliability of McConnell's classification of patellar orientation in symptomatic and asymptomatic subjects—invited commentary. *J Orth Sports Phys Ther* 1999;29(7):388-91.

87. Powers CM, Perry J, Hsu A et al. Are patellofemoral pain and quadriceps femoris muscle torque associated with locomotor function? *Phys Ther* 1997;77(10):1063-75.
88. Thomee R, Renstrom P, Karlsson J et al. Patellofemoral pain syndrome in young women. II. Muscle function in patients and healthy controls. *Scand J Sports Med* 1995;5:245-51.
89. Lankhorst NE, Bierma-Zeinstra SMA, van Middelkoop M. Factors associated with patellofemoral pain syndrome: a systematic review. *Br J Sports Med* 2013;47(4):193-206.
90. Giles LS, Webster KE, McClelland JA et al. Does quadriceps atrophy exist in individuals with patellofemoral pain? A systematic literature review with meta-analysis. *J Orth Sports Phys Ther* 2013;43(11):766-76.
91. Lankhorst NE, Bierma-Zeinstra SMA, van Middelkoop M. Risk factors for patellofemoral pain syndrome: a systematic review. *J Orth Sports Phys Ther* 2012;42(2):81-A12.
92. Boucher J, King M, Lefebvre R et al. Quadriceps femoris muscle activity in patellofemoral pain syndrome. *Am J Sports Med* 1992;20(5):527-32.
93. Mariano P, Caruso I. An electromyographic investigation of subluxation of the patella. *J Bone Joint Surg* 1979;61:169-71.
94. Morrish GM, Woledge RC. A comparison of the activation of muscles moving the patella in normal subjects and in patients with chronic patellofemoral problems. *Scand J Rehabil Med* 1997;29(1):43-8.
95. Petschnig R, Baron R, Engel A et al. Objectivation of the effects of knee problems on vastus medialis and vastus lateralis with EMG and dynamometry. *PM&R* 1991;2:50-4.
96. Miller JP. Vastus medialis obliquus and vastus lateralis activity in patients with and without patellofemoral pain syndrome. *J Sport Rehabil* 1997;6:1-10.
97. Powers CM, Landel RF, Perry J. Timing and intensity of vastus muscle activity during functional activities in subjects with and without patellofemoral pain. *Phys Ther* 1996;76:946-55.
98. Sheehy P, Burdett RC, Irrgang JJ et al. An electromyographic study of vastus medialis oblique and vastus lateralis activity while ascending and descending steps. *J Orth Sports Phys Ther* 1998;27:423-9.
99. Coqueiro K, Bevilacqua-Grossi D, Berzin F et al. Analysis on the activation of the VMO and VLL muscles during semisquat exercises with and without hip adduction in individuals with patellofemoral pain syndrome. *J Electromyog Kinesiol* 2005;15:596-603.
100. Souza DR, Gross M. Comparison of vastus medialis obliquus: vastus lateralis muscle integrated electromyographic ratios between healthy subjects and patients with patellofemoral pain. *Phys Ther* 1991;71(4):310-20.
101. Cavazzuti L, Merlo A, Orlandi F et al. Delayed onset of electromyographic activity of vastus medialis obliquus relative to vastus lateralis in subjects with patellofemoral pain syndrome. *Gait & Posture* 2010;32:290-5.
102. Sakai N, Lou Z-P, Rand JA et al. The influence of weakness in the vastus medialis oblique muscle on the patellofemoral joint: an in vitro biomechanical study. *Clin Biomech* 2000;15:335-9.
103. Neptune RR, Wright IC, van dBAJ. The influence of orthotic devices and vastus medialis strength and timing on patellofemoral loads during running. *Clin Biomech* 2000;15(8):611-8.
104. Cowan SM, Hodges PW, Bennell KL et al. Altered vastii recruitment when people with patellofemoral pain syndrome complete a postural task. *Arch Phys Med Rehabil* 2002;83(7):989-95.
105. Cesarelli M, Bifulco P, Bracale M. Study of the control strategy of the quadriceps in anterior knee pain. *IEEE Trans Rehabil Eng* 2000;8:330-41.
106. Cowan SM, Bennell KL, Hodges PW et al. Delayed onset of electromyographic activity of vastus medialis obliquus relative to vastus lateralis in subjects with patellofemoral pain syndrome. *Arch Phys Med Rehabil* 2001;82(2):183-9.
107. Giles LS, Webster KE, McClelland JA et al. Atrophy of the quadriceps is not isolated to the vastus medialis oblique in individuals with patellofemoral pain. *J Orth Sports Phys Ther* 2015;45(8):613-19.
108. Barton CJ, Lack S, Hemmings S et al. The 'Best Practice Guide to Conservative Management of Patellofemoral Pain': incorporating level 1 evidence with expert clinical reasoning. *Br J Sports Med* 2015;49(14):923-34.
109. Thomee R. A comprehensive treatment approach for patellofemoral pain syndrome in young women. *Phys Ther* 1997;77(12):1690-703.
110. Crossley K, Bennell K, Green S et al. Physical therapy for patellofemoral pain: a randomised, double-blinded, placebo-controlled study. *Am J Sports Med* 2002;30(6):857-65.
111. Collins N, Crossley K, Beller E et al. Foot orthoses and physiotherapy in the treatment of patellofemoral pain syndrome: randomised clinical trial. *BMJ* 2008;337:a1735.
112. Collins NJ, Bisset LM, Crossley KM et al. Efficacy of nonsurgical interventions for anterior knee pain: systematic review and meta-analysis of randomised trials. *Sports Med* 2012;42(1):31-49.
113. Crossley KM, van Middelkoop M, Callaghan MJ et al. Patellofemoral pain consensus statement from the 4th International Patellofemoral Pain Research Retreat, Manchester Part 2: recommended physical interventions (exercise, taping, bracing, foot orthoses and combined interventions). *Br J Sports Med* 2016;50(14):844-52.
114. Cowan SM, Bennell KL, Hodges PW. Therapeutic patellar taping changes the timing of vasti muscle activation in people with patellofemoral pain syndrome. *Clin J Sport Med* 2002;12:339-47.
115. Lack S, Barton C, Sohan OG et al. Proximal muscle rehabilitation is effective for patellofemoral pain: a systematic review with meta-analysis. *Br J Sports Med* 2015;49(21):1365-76.
116. Noehren B, Scholz J, Davis I. The effect of real-time gait retraining on hip kinematics, pain and function in subjects

- with patellofemoral pain syndrome. *Br J Sports Med* 2011;45(9): 691-6.
117. Garber CE, Blissmer B, Deschenes MR et al. Quantity and quality of exercise for developing and maintaining cardiorespiratory, musculoskeletal and neuromotor fitness in apparently healthy adults: guidance for prescribing exercise. *Med Sci Sports Exerc* 2011;43(7):1334-59.
 118. Barton C, Balachandar V, Lack S et al. Patellar taping for patellofemoral pain: a systematic review and meta-analysis to evaluate clinical outcomes and biomechanical mechanisms. *Br J Sports Med* 2014;48(6):417-24.
 119. Crossley KM, Cowan SM, Bennell KL et al. Patellar taping: is clinical success supported by scientific evidence? *Man Ther* 2000;5(3):142-50.
 120. Callaghan MJ. Patellar taping, the theory versus the evidence: a review. *Phys Ther Rev* 1997;2:181-3.
 121. Collins NJ, Bisset LM, Crossley KM et al. Efficacy of nonsurgical interventions for anterior knee pain: systematic review and meta-analysis of randomized trials. *Sports Med* 2012; 42(1):31-49.
 122. Franettovich M, Chapman A, Blanch P et al. Continual use of augmented low-Dye taping increases arch height in standing but does not influence neuromotor control of gait. *Gait & Posture* 2010;31(2):247-50.
 123. Vicenzino B, Collins N, Cleland J et al. A clinical prediction rule for identifying patients with patellofemoral pain who are likely to benefit from foot orthoses: a preliminary determination. *Br J Sports Med* 2010;44(12):862-6.
 124. Vicenzino B. Foot orthotics in the treatment of lower limb conditions: a musculoskeletal physiotherapy perspective. *Man Ther* 2004;9(4):185-96.
 125. Barton CJ, Menz HB, Levinger P et al. Greater peak rearfoot eversion predicts foot orthoses efficacy in individuals with patellofemoral pain syndrome. *Br J Sports Med* 2011;45(9): 697-701.
 126. Powers CM, Ward SR, Chan L-D et al. The effect of bracing on patella alignment and patellofemoral joint contact area. *Med Sci Sports Exerc* 2004;36(7):1226-32.
 127. Powers CM, Ward SR, Chen Y et al. The effect of bracing on patellofemoral joint stress during free and fast walking. *Am J Sports Med* 2004;32(1):224-31.
 128. Lun VMY, Wiley Preston J, Meeuwisse WH et al. Effectiveness of patellar bracing for treatment of patellofemoral pain syndrome. *Clin J Sport Med* 2005;15(4):235-40.
 129. Barton CJ, Munteanu SE, Menz HB et al. The efficacy of foot orthoses in the treatment of individuals with patellofemoral pain syndrome: a systematic review. *Sports Med* 2010;40(5):377-95.
 130. van der Heijden RA, Lankhorst NE, van Linschoten R et al. Exercise for treating patellofemoral pain syndrome. *Cochrane Database Syst Rev* 2015;1(CD010387).
 131. Mills K, Blanch P, Dev P et al. A randomised control trial of short term efficacy of in-shoe foot orthoses compared with a wait and see policy for anterior knee pain and the role of foot mobility. *Br J Sports Med* 2012;46(4):247-52.
 132. Swart NM, van Linschoten R, Bierma-Zeinstra SMA et al. The additional effect of orthotic devices on exercise therapy for patients with patellofemoral pain syndrome: a systematic review. *Br J Sports Med* 2012;46(8):570-7.
 133. Collins NJ, Bisset LM, Crossley KM et al. Efficacy of nonsurgical interventions for anterior knee pain systematic review and meta-analysis of randomized trials. *Sports Med* 2012;42(1):31-49.
 134. Bolgia LA, Boling MC. An update for the conservative management of patellofemoral pain syndrome: a systematic review of the literature from 2000 to 2010. *Int J Sports Phys Ther* 2011;6(2):112-25.
 135. Lack S, Barton C, Sohan OG et al. Proximal muscle rehabilitation is effective for patellofemoral pain: a systematic review with meta-analysis. *Br J Sports Med* 2015;49(21): 1365-76.
 136. Kooiker L, van De Port IG, Weir A et al. Effects of physical therapist-guided quadriceps-strengthening exercises for the treatment of patellofemoral pain syndrome: a systematic review. *J Orthop Sports Phys Ther* 2014;44(6):391-B1.
 137. Clijsen R, Fuchs J, Taeymans J. Effectiveness of exercise therapy in treatment of patients with patellofemoral pain syndrome: systematic review and meta-analysis. *Phys Ther* 2014;94(12):1697-708.
 138. Hossain M, Alexander P, Burls A et al. Foot orthoses for patellofemoral pain in adults. *Cochrane Database Syst Rev* 2011(1):CD008402.
 139. Kalron A, Bar-Sela S. A systematic review of the effectiveness of kinesio taping—fact or fashion? *Eur J Phys Rehabil Med* 2013;49:699-709.
 140. Callaghan MJ, Selfe J. Patellar taping for patellofemoral pain syndrome in adults. *Cochrane Database Syst Rev* 2012;4:CD006717.
 141. Lake DA, Wofford NH. Effect of therapeutic modalities on patients with patellofemoral pain syndrome: a systematic review. *Sports Health* 2011;3(2):182-9.
 142. Gecha SR, Torg Js. Clinical prognosticators for the efficacy of retinacular release surgery to treat patellofemoral pain. *Clin Orthop Rel Res* 1990;253:203-8.
 143. Crossley KM, Vicenzino B, Lentzos J et al. Exercise, education, manual-therapy and taping compared to education for patellofemoral osteoarthritis: A blinded, randomised clinical trial. *Osteoarthritis Cartilage* 2015;23(9):1457-64.
 144. Callaghan MJ, Parkes MJ, Hutchinson CE et al. A randomised trial of a brace for patellofemoral osteoarthritis targeting knee pain and bone marrow lesions. *Ann Rheum Dis* 2015; 74(6): 1164-70.
 145. Callaghan MJ, Parkes MJ, Hutchinson CE et al. A randomised trial of a brace for patellofemoral osteoarthritis targeting knee pain and bone marrow lesions. *Ann Rheum Dis* 2015;74(6): 1164-70.
 146. McWalter EJ, Hunter DJ, Harvey WF et al. The effect of a patellar brace on three-dimensional patellar kinematics in

- patients with lateral patellofemoral osteoarthritis. *Osteoarthritis Cartilage* 2011.
147. Frye JL, Ramey LN, Hart JM. The effects of exercise on decreasing pain and increasing function in patients with patellofemoral pain syndrome: a systematic review. *Sports Health* 2012;4(3):205-10.
 148. Peat G, Duncan RC, Wood LRJ et al. Clinical features of symptomatic patellofemoral joint osteoarthritis. *Arthritis Res Ther* 2012;14(2).
 149. Farrokhi S, Piva SR, Gil AB et al. Association of severity of coexisting patellofemoral disease with increased impairments and functional limitations in patients with knee osteoarthritis. *Arthritis Care Res* 2013;65(4):544-51.
 150. Fok LA, Schache AG, Crossley KM et al. Patellofemoral joint loading during stair ambulation in people with patellofemoral osteoarthritis. *Arthritis Rheum* 2013;65(8):2059-69.
 151. Amin S, Baker K, Niu J et al. Quadriceps strength and the risk of cartilage loss and symptom progression in knee osteoarthritis. *Arthritis Rheum* 2009;60(1):189-98.
 152. Stefanik JJ, Guermazi A, Zhu Y et al. Quadriceps weakness, patella alta, and structural features of patellofemoral osteoarthritis. *Arthritis Care Res* 2011;63(10):1391-97.
 153. Powers CM. The influence of altered lower-extremity kinematics on patellofemoral joint dysfunction: a theoretical perspective. *J Orthop Sports Phys Ther* 2003;33(11):639-46.
 154. Powers CM. The influence of abnormal hip mechanics on knee injury: a biomechanical perspective. *J Orthop Sports Phys Ther* 2010;40(2):42-51.
 155. Farrokhi S, Keyak JH, Powers CM. Individuals with patellofemoral pain exhibit greater patellofemoral joint stress: a finite element analysis study. *Osteoarthritis Cartilage* 2011;19(3):287-94.
 156. Nakagawa TH, Serrao FV, Maciel CD et al. Hip and knee kinematics are associated with pain and self-reported functional status in males and females with patellofemoral pain. *Int J Sports Med* 2013; 34(11):997-1002.
 157. Long-Rossi F, Salsich GB. Pain and hip lateral rotator muscle strength contribute to functional status in females with patellofemoral pain. *Physiother Res Int* 2010;15(1):57-64.
 158. Nakagawa TH, Baldon Rde M, Muniz TB et al. Relationship among eccentric hip and knee torques, symptom severity and functional capacity in females with patellofemoral pain syndrome. *Phys Ther Sport* 2011;12(3):133-9.
 159. Farrokhi S, O'Connell M, Fitzgerald GK. Altered gait biomechanics and increased knee-specific impairments in patients with coexisting tibiofemoral and patellofemoral osteoarthritis. *Gait Posture* 2015;41(1):81-5.
 160. Teng HL, MacLeod TD, Kumar D et al. Individuals with isolated patellofemoral joint osteoarthritis exhibit higher mechanical loading at the knee during the second half of the stance phase. *Clin Biomech (Bristol, Avon)* 2015;30(4):383-90.
 161. Crossley KM, Dorn TW, Ozturk H et al. Altered hip muscle forces during gait in people with patellofemoral osteoarthritis. *Osteoarthritis and Cartilage* 2012;20(11): 1243-49.
 162. Teng HL, MacLeod TD, Link TM et al. Higher knee flexion moment during the second half of the stance phase of gait is associated with magnetic resonance imaging progression of patellofemoral joint osteoarthritis. *J Orthop Sports Phys Ther* 2015:1-32.
 163. Rathleff MS, Skuldbol SK, Rasch MN et al. Care-seeking behaviour of adolescents with knee pain: a population-based study among 504 adolescents. *BMC Musculoskelet Disord* 2013;14:225.
 164. Rathleff MS, Roos EM, Olesen JL et al. Exercise during school hours when added to patient education improves outcome for 2 years in adolescent patellofemoral pain: a cluster randomised trial. *Br J Sports Med* 2015;49(6):406-12.
 165. Rathleff MS, Vicenzino B, Middelkoop M et al. Patellofemoral pain in adolescence and adulthood: same same, but different? *Sports Med* 2015; 45(11):1489-95.
 166. Rathleff CR, Olesen JL, Roos EM et al. Half of 12-15-year-olds with knee pain still have pain after one year. *Dan Med J* 2013;60(11):A4725.
 167. El-Metwally A, Salminen JJ, Auvinen A et al. Lower limb pain in a preadolescent population: prognosis and risk factors for chronicity-a prospective 1- and 4-year follow-up study. *Pediatrics* 2005;116(3):673-81.
 168. Schöttle PB, Fucentese SF, Romero J. Clinical and radiological outcome of medial patellofemoral ligament reconstruction with a semitendinosus autograft for patella instability. *Knee Surg Sports Traumatol Arthrosc* 2005;13(7):516-21.
 169. Smith TO, Donell S, Song F et al. Surgical versus non-surgical interventions for treating patellar dislocation. *Cochrane Database Syst Rev* 2015;2:CD008106.
 170. Blazina ME, Kerlan RK, Jobe FW et al. Jumper's knee. *Orth Clin North Am* 1973;4(3):665-78.
 171. Khan KM, Bonar F, Desmond PM et al. Patellar tendinosis (jumper's knee): findings at histopathologic examination, US and MR imaging. *Radiology* 1996;200(3): 821-7.
 172. Maffulli N, Khan KM, Puddu G. Overuse tendon conditions: time to change a confusing terminology. *Arthroscopy* 1998;14(8):840-3.
 173. Maffulli N, Binfield PM, Leach WJ et al. Surgical management of tendinopathy of the main body of the patellar tendon in athletes. *Clin J Sport Med* 1999;9(2):58-62.
 174. Visentini PJ, Khan KM, Cook JL et al. The VISA score: an index of severity of symptoms in patients with jumper's knee (patellar tendinosis). *J Sci Med Sport* 1998;1(1):22-8.
 175. Khan KM, Maffulli N, Coleman BD et al. Patellar tendinopathy: some aspects of basic science and clinical management. *Br J Sports Med* 1998;32:346-55.
 176. Richards DP, Ajemian SV, Wiley JP et al. Knee joint dynamics predict patellar tendinitis in elite volleyball players. *Am J Sports Med* 1996;24(5):676-83.
 177. Cook JL, Khan KM, Kiss ZS et al. Patellar tendinopathy in junior basketball players: a controlled clinical and ultrasonographic

- study of 268 patellar tendons in players aged 14–18 years. *Scand J Med Sci Sports* 2000;10(4):216–20.
178. Edwards S, Steele JR, McGhee DE et al. Landing strategies of athletes with an asymptomatic patellar tendon abnormality. *Med Sci Sports Exerc* 2010;42(11):2072–80.
 179. Malliaras P, Cook JL, Kent P. Reduced ankle dorsiflexion range may increase the risk of patellar tendon injury among volleyball players. *J Sci Med Sport* 2006;9(4):304–9.
 180. van Ark M, Docking SI, van den Akker-Scheek I et al. Does the adolescent patellar tendon respond to 5 days of cumulative load during a volleyball tournament? *Scand J Med Sci Sports* 2016;26(2):189–96.
 181. Rio E, Kidgell D, Moseley GL et al. Tendon neuroplastic training: changing the way we think about tendon rehabilitation: a narrative review. *Br J Sports Med* 2016;50(4):209–15.
 182. Fredberg U, Bolvig L, Andersen NT. Prophylactic training in asymptomatic soccer players with ultrasonographic abnormalities in Achilles and patellar tendons. *Am J Sports Med* 2008;36(3):451–60.
 183. Visnes H, Hoksrud A, Cook J et al. No effect of eccentric training on jumper's knee in volleyball players during the competitive season. *Clin J Sport Med* 2005;15(4):227–34.
 184. Fredberg U, Bolvig L. Significance of ultrasonographically detected asymptomatic tendinosis in the patellar and Achilles tendons of elite soccer players: a longitudinal study. *Am J Sports Med* 2002;30(4):488–91.
 185. Visnes H, Bahr R. The evolution of eccentric training as treatment for patellar tendinopathy (jumper's knee): a critical review of exercise programmes. *Br J Sports Med* 2007;41(4):217–23.
 186. Stasinopoulos D, Stasinopoulos I. Comparison of effects of exercise programme, pulsed ultrasound and transverse friction in the treatment of chronic patellar tendinopathy. *Clin Rehabil* 2004;18(4):347–52.
 187. Wilson J, Sevier T, Helfst R et al. Comparison of rehabilitation methods in the treatment of patellar tendinitis. *J Sport Rehabil* 2000;9:304–14.
 188. Pellicchia GL, Hamel H, Behnke P. Treatment of infrapatellar tendinitis: a combination of modalities and transverse friction massage versus iontophoresis. *J Sport Rehabil* 1994;3:135–45.
 189. Cook JL, Purdam CR. Is tendon pathology a continuum? A pathology model to explain the clinical presentation of load-induced tendinopathy. *Br J Sports Med* 2009;43(6):409–16.
 190. Fallon K, Purdam C, Cook J et al. A 'polypill' for acute tendon pain in athletes with tendinopathy? *J Sci Med Sport* 2008;11(3):235–8.
 191. de Vries A, Zwerver J, Diercks R et al. Effect of patellar strap and sports tape on pain in patellar tendinopathy: a randomized controlled trial. *Scand J Med Sci Sports* 2015 Sep 17. doi: 10.1111/sms.12556. [Epub ahead of print].
 192. Mani-Babu S, Morrissey D, Waugh C et al. The effectiveness of extracorporeal shock wave therapy in lower limb tendinopathy: a systematic review. *Am J Sports Med* 2015;43(3):752–61.
 193. Bosch G, de Mos M, van Binsbergen R et al. The effect of focused extracorporeal shock wave therapy on collagen matrix and gene expression in normal tendons and ligaments. *Equine Vet J* 2009;41(4):335–41.
 194. de Vos RJ, Heijboer MP, Weinans H et al. Tendon structure's lack of relation to clinical outcome after eccentric exercises in chronic midportion Achilles tendinopathy. *J Sport Rehabil* 2012;21(1):34–43.
 195. de Jonge S, Tol JL, Weir A et al. The tendon structure returns to asymptomatic values in nonoperatively treated Achilles tendinopathy but is not associated with symptoms: a prospective study. *Am J Sports Med* 2015;43(12):2950–8.
 196. Drew BT, Smith TO, Littlewood C et al. Do structural changes (eg, collagen/matrix) explain the response to therapeutic exercises in tendinopathy: a systematic review. *Br J Sports Med* 2014;48(12):966–72.
 197. Docking SI, Cook J. Pathological tendons maintain sufficient aligned fibrillar structure on ultrasound tissue characterization (UTC). *Scand J Med Sci Sports* 2016;26(6):675–83.
 198. de Vos RJ, Weir A, Tol JL et al. No effects of PRP on ultrasonographic tendon structure and neovascularisation in chronic midportion Achilles tendinopathy. *Br J Sports Med* 2011;45(5):387–92.
 199. de Vos RJ, van Veldhoven PL, Moen MH et al. Autologous growth factor injections in chronic tendinopathy: a systematic review. *Br Med Bull* 2010;95:63–77.
 200. Sanderson LM, Bryant A. Effectiveness and safety of prolotherapy injections for management of lower limb tendinopathy and fasciopathy: a systematic review. *J Foot Ankle Res* 2015;8:57.
 201. Jensen K, Di Fabio RP. Evaluation of eccentric exercise in treatment of patellar tendinitis. *Phys Ther* 1989;69(3):211–6.
 202. Cannell LJ, Taunton JE, Clement DB et al. A randomised clinical trial of the efficacy of drop squats or leg extension/leg curl exercises to treat clinically diagnosed jumper's knee in athletes: pilot study. *Br J Sports Med* 2001;35:60–4.
 203. Purdam CR, Cook JL, Hopper DM et al. Discriminative ability of functional loading tests for adolescent jumper's knee. *Phys Ther Sport* 2003;4:3–9.
 204. Young MA, Cook JL, Purdam CR et al. Eccentric decline squat protocol offers superior results at 12 months compared with traditional eccentric protocol for patellar tendinopathy in volleyball players. *Br J Sports Med* 2005;39(2):102–5.
 205. Bahr R, Fossan B, Loken S et al. Surgical treatment compared with eccentric training for patellar tendinopathy (jumper's knee). A randomized, controlled trial. *J Bone Joint Surg Am* 2006;88(8):1689–98.
 206. Kongsgaard M, Kovanen V, Aagaard P et al. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scand J Med Sci Sport* 2009;19(6):790–802.

207. Alfredson H, Öhberg L. Neovascularisation in chronic painful patellar tendinosis—promising results after sclerosing neovessels outside the tendon challenge the need for surgery. *Med Sci Sports Exerc* 2005;13(2):74–80.
208. Hoksrud A, Öhberg L, Alfredson H et al. Ultrasound-guided sclerosis of neovessels in painful chronic patellar tendinopathy. *Am J Sports Med* 2006;34(11):1738–46.
209. Alfredson H, Öhberg L, Forsgren S. Is vasculo-neural ingrowth the cause of pain in chronic Achilles tendinosis? An investigation using ultrasonography and colour Doppler, immunohistochemistry and diagnostic injections. *Knee Surg Sports Traumatol Arthrosc* 2003;11(5):334–8.
210. Fredberg U, Bolvig L, Pfeiffer-Jensen M et al. Ultrasonography as a tool for diagnosis, guidance of local steroid injection and, together with pressure algometry, monitoring of the treatment of athletes with chronic jumper's knee and Achilles tendinitis: a randomized, double-blind, placebo-controlled study. *Scand J Rheumatol* 2004;33(2):94–101.
211. Liddle AD, Rodriguez-Merchan EC. Platelet-rich plasma in the treatment of patellar tendinopathy: a systematic review. *Am J Sports Med* 2015;43(10):2583–90.
212. Hoksrud A, Öhberg L, Alfredson H et al. Ultrasound-guided sclerosis of neovessels in painful chronic patellar tendinopathy: a randomized controlled trial. *Am J Sports Med* 2006;34(11):1738–46.
213. Coleman BD, Khan KM, Kiss ZS et al. Open and arthroscopic patellar tenotomy for chronic patellar tendinopathy: a retrospective outcome study. *Am J Sports Med* 2000;28(2):183–90.
214. Willberg L, Sunding K, Forssblad M et al. Sclerosing polidocanol injections or arthroscopic shaving to treat patellar tendinopathy/jumper's knee? A randomised controlled study. *Br J Sports Med* 2011;45(5):411–5.
215. Brockmeyer M, Diehl N, Schmitt C et al. Results of surgical treatment of chronic patellar tendinosis (jumper's knee): a systematic review of the literature. *Arthroscopy* 2015;31(12):2424–9.e3.
216. Coleman BD, Khan KM, Maffulli N et al. Studies of surgical outcome after patella tendinopathy: clinical significance of methodological deficiencies and guidelines for future studies. *Scand J Med Sci Sports* 2000;10:2–11.
217. Paavola M, Paakkala T, Kannus P et al. Ultrasonography in the differential diagnosis of Achilles tendon injuries and related disorders. A comparison between pre-operative ultrasonography and surgical findings. *Acta Radiol* 1998;39(6):612–19.
218. Reinus WR, Shady KL, Mirowitz SA et al. MR diagnosis of rotator cuff tears of the shoulder: value of using T2-weighted fat-saturated images. *Am J Roentgenol* 1995;164(6):1451–5.
219. Saddik D, McNally EG, Richardson M. MRI of Hoffa's fat pad. *Skeletal Radiol* 2004;33(8):433–44.
220. Ducher G, Cook J, Spurrier D et al. Ultrasound imaging of the patellar tendon attachment to the tibia during puberty: a 12-month follow-up in tennis players. *Scand J Med Sci Sports* 2010;20(1):e35–e40.

CHAPTER 37

1. Baker RL, Fredericson M. Iliotibial Band Syndrome in Runners: Biomechanical Implications and Exercise Interventions. *Phys Med Rehabil Clin N Am* 2016;27(1):53–77.
2. Fairclough J, Hayashi K, Toumi H et al. Is iliotibial band syndrome really a friction syndrome? *J Sci Med Sport* 2007;10(2):74–6.
3. Ellis R, Hing W, Reid D. Iliotibial band friction syndrome—a systematic review. *Man Ther* 2007;12(3):200–8.
4. Falvey EC, Clark RA, Franklyn-Miller A et al. Iliotibial band syndrome: an examination of the evidence behind a number of treatment options. *Scand J Med Sci Sports* 2010;20:580–7.
5. Aderem J, Louw QA. Biomechanical risk factors associated with iliotibial band syndrome in runners: a systematic review. *BMC Musculoskelet Disord* 2015;16:356.
6. Huang BK, Campos JC, Michael Peschka PG et al. Injury of the gluteal aponeurotic fascia and proximal iliotibial band: anatomy, pathologic conditions, and MR imaging. *Radiographics* 2013;33(5):1437–52.
7. Noehren B, Davis I, Hamill J. ASB clinical biomechanics award winner 2006 prospective study of the biomechanical factors associated with iliotibial band syndrome. *Clin Biomech* 2007;22(9):951–6.
8. Fredericson M, Cookingham CL, Chaudhari AM et al. Hip abductor weakness in distance runners with iliotibial band syndrome. *Clin J Sport Med* 2000;10:169–75.
9. Fredericson M, Wolf C. Iliotibial band syndrome in runners: innovations in treatment. *Sports Med* 2005;35(5):451–9.
10. Messier SP, Edwards DG, Martin DF et al. Etiology of iliotibial band friction syndrome in distance runners. *Med Sci Sports Exerc* 1995;27(7):951–60.
11. Weckström K, Söderström J. Radial extracorporeal shockwave therapy compared with manual therapy in runners with iliotibial band syndrome. *J Back Musculoskelet Rehabil* 2016;29(1):161–70.
12. Schwellnus MP, Theunissen L, Noakes TD et al. Anti-inflammatory and combined anti-inflammatory/analgesic medication in the early management of iliotibial band friction syndrome. A clinical trial. *S Afr Med J* 1991;79(10):602–6.
13. Gunter P, Schwellnus MP. Local corticosteroid injection in iliotibial band friction syndrome in runners: a randomised controlled trial. *Br J Sports Med* 2004;38(3):269–72; discussion 72.
14. Willis RB. Meniscal injuries in children and adolescents. *Op Tech Sports Med* 2006;14(3):197–202.
15. Kocher MS, Klingele K, Rassman SO. Meniscal disorders: normal, discoid, and cysts. *Orthop Clin North Am* 2003;34(3):329–40.
16. Sharma L, Song J, Felson DT et al. The role of knee alignment in disease progression and functional decline in knee osteoarthritis. *JAMA* 2001;286(2):188–95.
17. Brouwer GM, van Tol AW, Bergink AP et al. Association between valgus and varus alignment and the development and progression of radiographic osteoarthritis of the knee. *Arthritis Rheum* 2007;56(4):1204–11.

18. Altman RD, Moskowitz R. Intraarticular sodium hyaluronate (Hyalgan) in the treatment of patients with osteoarthritis of the knee: a randomized clinical trial. *J Rheumatol* 1998;25:2203-12.
19. Petrella RJ, Bartha C. Home based exercise therapy for older patients with knee osteoarthritis: a randomized clinical trial. *J Rheumatol* 2000;27(9):2215-21.
20. Squire MW, Callaghan JJ, Goetz DD et al. Unicompartmental knee replacement. A minimum 15 year followup study. *Clin Orthop* 1999;367:61-72.
21. Merican AM, Amis AA. Anatomy of the lateral retinaculum of the knee. *J Bone Joint Surg Br* 2008;90-B(4):527-34.
22. Shellock FG, Stone KR, Crues JV. Development and clinical application of kinematic MRI of the patellofemoral joint using an extremity MR system. *Med Sci Sports Exerc* 1999;31(5):788-91.
23. Panni AS, Tartarone M, Patricola A et al. Long-term results of lateral retinacular release. *Arthroscopy* 2005;21(5):526-31.
24. Change-Larsen K, Littlewood C, Garth A. Prone hip extension with lower abdominal hollowing improves the relative timing of gluteus maximus activation in relation to biceps femoris. *Man Ther* 2010;15(1):61-5.
25. Longo UG, Garau G, Denaro V et al. Surgical management of tendinopathy of biceps femoris tendon in athletes. *Disabil Rehabil* 2008;30(20-22):1602-7.
26. Forster BB, Lee JS, Kelly S et al. Proximal tibiofibular joint: an often-forgotten cause of lateral knee pain. *Am J Roentgenol* 2007;188(4):W359-66.
27. Scott J, Lee H, Barsoum W et al. The effect of tibiofemoral loading on proximal tibiofibular joint motion. *J Anat* 2007; 211(5):647-53.
28. Sekiya JK, Kuhn JE. Instability of the proximal tibiofibular joint. *J Am Acad Orthop Surg* 2003;11(2):120-8.
29. Luhmann SJ, Schoenecker PL, Dobbs MB et al. Adolescent patellofemoral pain: implicating the medial patellofemoral ligament as the main pain generator. *J Child Orthop* 2008;2: 269-77.
30. Fithian DC, Powers CM, Khan N. Rehabilitation of the knee after medial patellofemoral ligament reconstruction. *Clin Sports Med* 2010;29:283-90.
31. Bedi A, Kelly NH, Baad M et al. Dynamic contact mechanics of the medial meniscus as a function of radial tear, repair, and partial meniscectomy. *J Bone Joint Surg Am* 2010;92(6):1398-408.
32. Nicholson S, Dickman K, Maradiegue A. Reducing premature osteoarthritis in the adolescent through appropriate screening. *J Ped Nurs* 2009;24(1):69-74.
33. Bout-Tabaku S, Best TM. The adolescent knee and risk for osteoarthritis—an opportunity or responsibility for sport medicine physicians? *Curr Sports Med Rep* 2010;9(6):329-31.
34. Bashir A, Grey ML, Hartke J et al. Nondestructive imaging of human cartilage glycosaminoglycan concentration by MRI. *Magn Reson Med* 1999;41:857-65.
35. Olsen O-E, Myklebust G, Engebretsen L et al. Exercises to prevent lower limb injuries in youth sports: cluster randomised controlled trial. *BMJ* 2005;330(7489):449.
36. Twomey D, Finch C, Roediger E et al. Preventing lower limb injuries: is the latest evidence being translated into the football field? *J Sci Med Sport* 2009;12(4):452-6.
37. Pasanen K, Parkkari J, Pasanen M et al. Neuromuscular training and the risk of leg injuries in female floorball players: cluster randomised controlled study. *Br J Sports Med* 2008;42(10):802-5.
38. Kirkley A, Webster-Bogaert S, Litchfield R et al. The effect of bracing on varus gonarthrosis. *J Bone Joint Surg Am* 1999;81A:539-48.
39. Komistek RD, Dennis DA, Northcutt EJ et al. An in vivo analysis of the effectiveness of the osteoarthritic knee brace during heel-strike of gait. *J Arthroplasty* 1999;14:738-42.
40. Bellamy N, Campbell J, Robinson V et al. Viscosupplementation for the treatment of osteoarthritis of the knee. *Cochrane Database Syst Rev* 2005;18(2):CD005321.
41. Kessler MA, Behrend H, Henz S et al. Function, osteoarthritis and activity after ACL-rupture: 11 years follow-up results of conservative versus reconstructive treatment. *Knee Surg Sports Traumatol Arthrosc* 2008;16:442-8.
42. Øiestad BE, Engebretsen L, Storheim K et al. Knee osteoarthritis after anterior cruciate ligament injury: a systematic review. *Am J Sports Med* 2009;37(7):1434-43.
43. Rennie WJ, Saifuddin A. Pes anserine bursitis: incidence in symptomatic knees and clinical presentation. *Skeletal Radiol* 2005;34(7):395-8.
44. Alvarez-Nemegyei J. Risk factors for pes anserinus tendinitis/bursitis syndrome: a case control study. *J Clin Rheumatol* 2007;13(2):63-5.
45. Finnoff JT, Nutz DJ, Henning PT et al. Accuracy of ultrasound-guided versus unguided pes anserinus bursa injections. *PMR* 2010;2(8):732-9.
46. Niitsu M, Ikeda K, Iijima T et al. MR imaging of Pellegrini-Stieda disease. *Radiat Med* 1999;17: 405-9.
47. Theivendran K, Lever CJ, Hart WJ. Good result after surgical treatment of Pellegrini-Stieda syndrome. *Knee Surg Sports Traumatol Arthrosc* 2009;17(10):1231-3.
48. Fritschy D, Fasel J, Imbert J-C et al. The popliteal cyst. *Knee Surg Sports Traumatol Arthrosc* 2006;14:623-8.
49. Muché JA, Lento PH. Posterior knee pain and its causes: a clinician's guide to expediting diagnosis. *Phys Sportsmed* 2004;32(3):23-30.
50. Di Sante L, Paoloni M, Ioppolo F et al. Ultrasound-guided aspiration and corticosteroid injection of Baker's cysts in knee osteoarthritis: a prospective observational study. *Am J Phys Med Rehabil* 2010;89(12):970-5.
51. LaPrade RF, Wozniczka JK, Stellmaker MP et al. Analysis of the static function of the popliteus tendon and evaluation of an anatomic reconstruction: the 'fifth ligament' of the knee. *Am J Sports Med* 2010;38(3):543-9.
52. Guha AR, Gorgees KA, Walker DI. Popliteus tendon rupture: a case report and review of the literature. *Br J Sports Med* 2003;37(4):358-60.
53. Garrick JG, Webb DR. *Sports injuries: diagnosis and management*. Philadelphia: WB Saunders, 1990.

54. Soda N, Fujihashi Y, Aoki T. In vivo ultrasound imaging of the popliteus muscle: investigation of functional characteristics. *J Phys Ther Sci* 2016;28(3):979-82.
55. Smith J, Finnoff JT, Santaella-Sante B et al. Sonographically guided popliteus tendon sheath injection: techniques and accuracy. *J Ultrasound Med* 2010;29(5):775-82.

CHAPTER 38

1. Pedowitz R, Hargens A. Acute and chronic compartment syndromes. In: Garrett W, Speer K, Kirkendall D (eds). *Principles and Practice of Orthopaedic Sports medicine*. Philadelphia: Lippincott Williams & Wilkins, 2000:87-97.
2. Schneiders AG, Sullivan SJ, Hendrick PA et al. The ability of clinical tests to diagnose stress fractures: a systematic review and meta-analysis. *J Orthop Sports Phys Ther* 2012;42(9):760-71.
3. Winters M, Moen MH, Zimmermann WO et al. The medial tibial stress syndrome score: a new patient-reported outcome measure. *Br J Sports Med*. 2016;50(19):1192-9.
4. Blackman PG. A review of chronic exertional compartment syndrome in the lower leg. *Med Sci Sports Exerc* 2000;32(3 Suppl):S4-10.
5. Fredericson M, Bergman AG, Hoffman KL et al. Tibial stress reaction in runners. Correlation of clinical symptoms and scintigraphy with a new magnetic resonance imaging grading system. *Am J Sports Med* 1995;23(4):472-81.
6. Yao L, Johnson C, Gentili A et al. Stress injuries of bone: analysis of MR imaging staging criteria. *Acad Radiol* 1998;5(1):34-40.
7. Arendt E, Agel J, Heikes C et al. Stress injuries to bone in college athletes: a retrospective review of experience at a single institution. *Am J Sports Med* 2003;31(6):959-68.
8. Batt ME, Ugalde V, Anderson MW et al. A prospective controlled study of diagnostic imaging for acute shin splints. *Med Sci Sports Exerc* 1998;30(11):1564-71.
9. van den Brand JGH, Nelson T, Verleisdonk EJMM et al. The diagnostic value of intracompartmental pressure measurement, magnetic resonance imaging, and near-infrared spectroscopy in chronic exertional compartment syndrome: a prospective study in 50 patients. *Am J Sports Med* 2005;33(5):699-704.
10. Rajasekaran S, Beavis C, Aly AR et al. The utility of ultrasound in detecting anterior compartment thickness changes in chronic exertional compartment syndrome: a pilot study. *Clin J Sport Med* 2013;23(4):305-11.
11. Gershuni DH, Gosink BB, Hargens AR et al. Ultrasound evaluation of the anterior musculofascial compartment of the leg following exercise. *Clin Orthop Relat Res* 1982;167(1):185-90.
12. Martinson H, Stokes MJ. Measurement of anterior tibial muscle size using real-time ultrasound imaging. *Eur J App Physiol Occ Physiol* 1991;63(3-4):250-4.
13. Birtles DB, Minden D, Wickes SJ et al. Chronic exertional compartment syndrome: muscle changes with isometric exercise. *Med Sci Sports Exerc* 2002;34(12):1900-6.
14. Birtles DB, Rayson MP, Jones DA et al. Effect of eccentric exercise on patients with chronic exertional compartment syndrome. *Eur J Appl Physiol* 2003;88(6):565-71.
15. Dahl M, Hansen P, Stål P et al. Stiffness and thickness of fascia do not explain chronic exertional compartment syndrome. *Clin Orth Rel Res* 2011;469(12):3495-500.
16. Sigmund EE, Novikov DS, Sui D et al. Time-dependent diffusion in skeletal muscle with the random permeable barrier model (RPBM): application to normal controls and chronic exertional compartment syndrome patients. *NMR Biomed* 2014;27(5):519-28.
17. Sigmund EE, Sui D, Ukpebor O et al. Stimulated echo diffusion tensor imaging and SPAIR T2-weighted imaging in chronic exertional compartment syndrome of the lower leg muscles. *J Magn Reson Imaging* 2013;38(5):1073-82.
18. Hislop M, Tierney P, Murray P et al. Chronic exertional compartment syndrome: the controversial 'fifth' compartment of the leg. *Am J Sports Med* 2003;31(5):770-6.
19. Pedowitz RA, Hargens AR, Mubarak SJ et al. Modified criteria for the objective diagnosis of chronic compartment syndrome of the leg. *Am J Sports Med* 1990;18(1):35-40.
20. Aweid O, Del Buono A, Malliaras P et al. Systematic review and recommendations for intracompartmental pressure monitoring in diagnosing chronic exertional compartment syndrome of the leg. *Clin J Sport Med* 2012;22(4):356-70.
21. Dharm-Datta S, Minden DF, Rosell PA et al. Dynamic pressure testing for chronic exertional compartment syndrome in the UK military population. *J R Army Med Corps* 2013;159(2):114-18.
22. Popp KL, Hughes JM, Smock AJ et al. Bone geometry, strength, and muscle size in runners with a history of stress fracture. *Med Sci Sports Exerc* 2009;41(12):2145-50.
23. Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016;50(5):273-80.
24. Fields KB, Sykes JC, Walker KM et al. Prevention of running injuries. *Curr Sports Med Rep* 2010;9(3):176-82.
25. Mountjoy M, Sundgot-Borgen J, Burke L et al. The IOC consensus statement: beyond the Female Athlete Triad—Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med* 2014;48(7):491-7.
26. Crowell HP, Milner CE, Hamill J et al. Reducing impact loading during running with the use of real-time visual feedback. *J Orthop Sports Phys Ther* 2010;40(4):206-13.
27. Edwards WB, Taylor D, Rudolph TJ et al. Effects of running speed on a probabilistic stress fracture mode. *Clin Biomech (Bristol, Avon)* 2010;25(4):372-7.
28. Davis IS, Bowser BJ, Mullineaux DR. Greater vertical impact loading in female runners with medically diagnosed injuries: a prospective investigation. 2016;50(14):887-92.
29. Napier C, Cochrane CK, Taunton JE et al. Gait modifications to change lower extremity gait biomechanics in runners: a systematic review. *Br J Sports Med* 2015;49(21):1382-8.

30. Dickson TB, Jr., Kichline PD. Functional management of stress fractures in female athletes using a pneumatic leg brace. *Am J Sports Med* 1987;15(1):86-9.
31. Whitelaw GP, Wetzler MJ, Levy AS et al. A pneumatic leg brace for the treatment of tibial stress fractures. *Clin Orthop Relat Res* 1991;(270):301-5.
32. Swenson EJ, Jr., DeHaven KE, Sebastianelli WJ et al. The effect of a pneumatic leg brace on return to play in athletes with tibial stress fractures. *Am J Sports Med* 1997;25(3):322-8.
33. Mollon B, da Silva V, Busse JW et al. Electrical stimulation for long-bone fracture-healing: a meta-analysis of randomized controlled trials. *J Bone Joint Surg Am* 2008;90(11):2322-30.
34. Gan TY, Kuah DE, Graham KS et al. Low-intensity pulsed ultrasound in lower limb bone stress injuries: a randomized controlled trial. *Clin J Sport Med* 2014;24(6):457-60.
35. Shima Y, Engebretsen L, Iwasa J et al. Use of bisphosphonates for the treatment of stress fractures in athletes. *Knee Surg Sports Traumatol Arthrosc* 2009;17(5):542-50.
36. Mallee WH, Weel H, van Dijk CN et al. Surgical versus conservative treatment for high-risk stress fractures of the lower leg (anterior tibial cortex, navicular and fifth metatarsal base): a systematic review. *Br J Sports Med* 2015;49(6):370-6.
37. Young AJ, McAllister DR. Evaluation and treatment of tibial stress fractures. *Clin Sports Med* 2006;25(1):117-28.
38. Detmer DE. Chronic shin splints. Classification and management of medial tibial stress syndrome. *Sports Med* 1986;3(6):436-46.
39. Craig DI. Current developments concerning medial tibial stress syndrome. *Phys Sportsmed* 2009;37(4):39-44.
40. Moen MH, Tol JL, Weir A et al. Medial tibial stress syndrome: a critical review. *Sports Med* 2009;39(7):523-46.
41. Bennett JE, Reinking MF, Pluemer B et al. Factors contributing to the development of medial tibial stress syndrome in high school runners. *J Orthop Sports Phys Ther* 2001;31(9):504-10.
42. Hubbard TJ, Carpenter EM, Cordova ML. Contributing factors to medial tibial stress syndrome: a prospective investigation. *Med Sci Sports Exerc* 2009;41(3):490-6.
43. Pell RF, Khanuja HS, Cooley GR. Leg pain in the running athlete. *J Am Acad Orthop Surg* 2004;12(6):396-404.
44. Magnusson HI, Westlin NE, Nyqvist F et al. Abnormally decreased regional bone density in athletes with medial tibial stress syndrome. *Am J Sports Med* 2001;29(6):712-15.
45. Magnusson HI, Ahlborg HG, Karlsson C et al. Low regional tibial bone density in athletes with medial tibial stress syndrome normalizes after recovery from symptoms. *Am J Sports Med* 2003;31(4):596-600.
46. Burne SG, Khan KM, Boudville PB et al. Risk factors associated with exertional medial tibial pain: a 12 month prospective clinical study. *Br J Sports Med* 2004;38(4):441-5.
47. Craig DI. Medial tibial stress syndrome: evidence-based prevention. *J Athl Train* 2008;43(3):316-18.
48. Holen KJ, Engebretsen L, Grontvedt T et al. Surgical treatment of medial tibial stress syndrome (shin splint) by fasciotomy of the superficial posterior compartment of the leg. *Scand J Med Sci Sports* 1995;5(1):40-3.
49. Fraimont M, Adamson G. Chronic exertional compartment syndrome. *J Am Acad Orthop Surg* 2003;11(4):268-76.
50. Barbour TD, Briggs CA, Bell SN et al. Histology of the fascial-periosteal interface in lower limb chronic deep posterior compartment syndrome. *Br J Sports Med* 2004;38(6):709-17.
51. Birtles DB, Rayson MP, Casey A et al. Venous obstruction in healthy limbs: a model for chronic compartment syndrome? *Med Sci Sports Exerc* 2003;35(10):1638-44.
52. Edmundsson D, Toolanen G, Thornell L et al. Evidence for low muscle capillary supply as a pathogenic factor in chronic compartment syndrome. *Scand J Med Sci Sports* 2010;20(6):805-13.
53. Balduini FC, Shenton DW, O'Connor KH et al. Chronic exertional compartment syndrome: correlation of compartment pressure and muscle ischemia utilizing ³¹P-NMR spectroscopy. *Clin Sports Med* 1993;12(1):151-65.
54. Owens S, Edwards P, Miles K et al. Chronic compartment syndrome affecting the lower limb: MIBI perfusion imaging as an alternative to pressure monitoring: two case reports. *Br J Sports Med* 1999;33:49-53.
55. Edmundsson D, Svensson O, Toolanen G. Intermittent claudication in diabetes mellitus due to chronic exertional compartment syndrome of the leg: an observational study of 17 patients. *Acta Orthop* 2008;79(4):534-9.
56. Williams EH, Detmer DE, Guyton GP et al. Non-invasive neurosensory testing used to diagnose and confirm successful surgical management of lower extremity deep distal posterior compartment syndrome. *J Brachial Plex Peripher Nerve Inj* 2009;4:4.
57. Litwiller DV, Amrami KK, Dahm DL et al. Chronic exertional compartment syndrome of the lower extremities: improved screening using a novel dual birdcage coil and in-scanner exercise protocol. *Skeletal Radiol* 2007;36(11):1067-75.
58. Oturai PS, Lorenzen T, Norregaard J et al. Evaluation of Tc-99m-tetrofosmin single-photon emission computed tomography for detection of chronic exertional compartment syndrome of the leg. *Scand J Med Sci Sports* 2006;16(4):282-6.
59. Hutchinson MR, Bederka B, Kopplin M. Anatomic structures at risk during minimal-incision endoscopically assisted fascial compartment releases in the leg. *Am J Sports Med* 2003;31(5):764-9.
60. Detmer DE, Sharpe K, Suft RL et al. Chronic compartment syndrome: diagnosis, management, and outcomes. *Am J Sports Med* 1985;13(3):162-70.
61. Farr D, Selesnick H. Chronic exertional compartment syndrome in a collegiate soccer player: a case report and literature review. *Am J Orthop* 2008;37(7):374-7.
62. Franklyn-Miller A, Roberts A, Hulse D et al. Biomechanical overload syndrome: defining a new diagnosis. *Br J Sports Med* 2014;48(6):415-6.

63. Barton CJ, Bonanno DR, Carr J et al. Running retraining to treat lower limb injuries: a mixed-methods study of current evidence synthesised with expert opinion. *Br J Sports Med* 2016;50(9):513–26.
64. Diebal AR, Gregory R, Alitz C et al. Forefoot running improves pain and disability associated with chronic exertional compartment syndrome. *Am J Sports Med* 2012;40(5):1060–7.
65. Soares A, Andriolo RB, Atallah AN et al. Botulinum toxin for myofascial pain syndromes in adults. *Cochrane Database Syst Rev* 2014;7:CD007533.
66. Isner-Horobeti ME, Dufour SP, Blaes C et al. Intramuscular pressure before and after botulinum toxin in chronic exertional compartment syndrome of the leg: a preliminary study. *Am J Sports Med* 2013;41(11):2558–66.
67. Wittstein J, Moorman CT, 3rd, Levin LS. Endoscopic compartment release for chronic exertional compartment syndrome: surgical technique and results. *Am J Sports Med* 2010;38(8):1661–6.
68. Esmail AN, Flynn JM, Ganley TJ et al. Acute exercise-induced compartment syndrome in the anterior leg. A case report. *Am J Sports Med* 2001;29(4):509–12.
69. Livingston KS, Meehan WP, 3rd, Hresko MT et al. Acute exertional compartment syndrome in young athletes: a descriptive case series and review of the literature. *Pediatr Emerg Care* 2016 doi: 10.1097/pec.0000000000000647. [Published online Feb 10]
70. Howard JL, Mohtadi NG, Wiley JP. Evaluation of outcomes in patients following surgical treatment of chronic exertional compartment syndrome in the leg. *Clin J Sport Med* 2000;10(3):176–84.
71. Micheli LJ, Solomon R, Solomon J et al. Surgical treatment for chronic lower-leg compartment syndrome in young female athletes. *Am J Sports Med* 1999;27(2):197–201.
72. Winkes MB, Hoogveen AR, Houterman S et al. Compartment pressure curves predict surgical outcome in chronic deep posterior compartment syndrome. *Am J Sports Med* 2012;40(8):1899–905.
73. Schepesis AA, Gill SS, Foster TA. Fasciotomy for exertional anterior compartment syndrome: is lateral compartment release necessary? *Am J Sports Med* 1999;27(4):430–5.
74. van Zoest WJ, Hoogveen AR, Scheltinga MR et al. Chronic deep posterior compartment syndrome of the leg in athletes: postoperative results of fasciotomy. *Int J Sports Med* 2008;29(5):419–23.
75. Schepesis AA, Fitzgerald M, Nicoletta R. Revision surgery for exertional anterior compartment syndrome of the lower leg: technique, findings, and results. *Am J Sports Med* 2005;33(7):1040–7.
76. Campano D, Robaina JA, Kusnezov N et al. Surgical Management for Chronic Exertional Compartment Syndrome of the Leg: A Systematic Review of the Literature. *Arthroscopy* 2016;32(7):1478–86.

77. Joy SM, Raudales R. Popliteal Artery Entrapment Syndrome. *Curr Sports Med Rep* 2015;14(5):364–7.

CHAPTER 39

1. Balias R, Alomar X, Rodas G et al. The soleus muscle: MRI, anatomic and histologic findings in cadavers with clinical correlation of strain injury distribution. *Skeletal Radiol* 2013;42:521–30.
2. Ekstrand J, Häggglund M, Walden M. Epidemiology of muscle injuries in professional football (soccer). *Am J Sports Med* 2011;39(6):1226–32.
3. Koulouris G, Ting AY, Jhamb A et al. Magnetic resonance imaging findings of injuries to the calf muscle complex. *Skeletal Radiol* 2007;36(10):921–7.
4. Orchard JW, Farhart P, Leopold C. Lumbar spine region pathology and hamstring and calf injuries in athletes: is there a connection? *Br J Sports Med* 2004;38(4):502–4.
5. Van Middelkoop M, Kolkman J, Van Ochten J et al. Risk factors for lower extremity injuries among male marathon runners. *Scand J Med Sci Sports* 2008;18(6):691–7.
6. Bizzini M, Junge A, Bahr R et al. Injuries and musculoskeletal complaints in referees and assistant referees selected for the 2006 FIFA World Cup: retrospective and prospective survey. *Br J Sports Med* 2009;43(7):490–7.
7. Urbano FL. Homan's sign in the diagnosis of deep venous thrombosis. *Hosp Physician* 2001;37(3):22–4.
8. Orchard JW, Alcott E, James T et al. Exact moment of a gastrocnemius muscle strain captured on video. *Br J Sports Med* 2002;36(3):222–3.
9. Bleakley CM, Glasgow P, MacAuley DC. PRICE needs updating, should we call the POLICE? *Br J Sports Med* 2012;46:220–1.
10. Chen WM, Park J, Park SB. Role of gastrocnemius-soleus muscle in forefoot force transmission at heel rise—a 3D finite element analysis. *J Biomech* 2012;45(10):1783–9.
11. Reynolds JF, Noakes TD, Schwellnus MP et al. Non-steroidal anti-inflammatory drugs fail to enhance healing of acute hamstring injuries treated with physiotherapy. *S Afr Med J* 1995;85(6):517–22.
12. Jarvinen TA, Jarvinen TL, Kaariainen M et al. Muscle injuries: biology and treatment. *Am J Sports Med* 2005;33(5):745–64.
13. Dixon J. Gastrocnemius vs soleus strain: how to differentiate and deal with calf muscle injuries. *Curr Rev Musculoskelet Med* 2009;2:74–7.
14. Radford JA, Burns J, Buchbinder R et al. Does stretching increase ankle dorsiflexion range of motion? A systematic review. *Br J Sports Med* 2006;40(10):870–5.
15. Gabbe BJ, Finch CF, Wajswelner H et al. Predictors of lower extremity injuries at the community level of Australian football. *Clin J Sport Med* 2004;14:56–63.
16. Connell DA, Schneider-Kolsky ME, Hoving JL et al. Longitudinal study comparing sonographic and MRI assessments of acute and

- healing hamstring injuries. *AJR Am J Roentgenol* 2004;183(4):975-84.
17. Delgado GJ, Chung CB, Lektrakul N et al. Tennis leg: clinical US study of 141 patients and anatomic investigation of four cadavers with MR imaging and US. *Radiology* 2002;224(1):112-9.
 18. Jennings A, Peterson R. Delayed reconstruction of medial head of gastrocnemius rupture: a surgical option. *Foot Ankle Int* 2013;34(6):904-7.
 19. Christodoulou A, Terzidis I, Natsis K et al. Soleus accessorius, an anomalous muscle in a young athlete: case report and analysis of the literature. *Br J Sports Med* 2004;38(6):e38.
 20. Rossi R, Bonasia DE, Tron A et al. Accessory soleus in the athletes: literature review and case report of a massive muscle in a soccer player. *Knee Surg Sports Traumatol Arthrosc* 2009;17(8):990-5.
 21. Doda N, Peh WC, Chawla A. Symptomatic accessory soleus muscle: diagnosis and follow-up on magnetic resonance imaging. *Br J Radiol* 2006;79(946):e129-32.
 22. Buschmann WR, Cheung Y, Jahss MH. Magnetic resonance imaging of anomalous leg muscles: accessory soleus, peroneus quartus and the flexor digitorum longus accessorius. *Foot Ankle* 1991;12(2):109-16.
 23. Levien LJ, Veller MG. Popliteal artery entrapment syndrome: more common than previously recognized. *J Vasc Surg* 1999;30:587-98.
 24. Hislop M, Kennedy D, Cramp B et al. Functional popliteal artery entrapment syndrome: poorly understood and frequently missed? A review of clinical features, appropriate investigations, and treatment options. *J Sports Med* 2014;2014:105953. doi: 10.1155/2014/105953. Epub 2014 Sep 7. Review.
 25. Stager A, Clement D. Popliteal artery entrapment syndrome. *Sports Med* 1999;28(1):61-70.
 26. Turnipseed WD. Functional popliteal artery entrapment syndrome: a poorly understood and often missed diagnosis that is frequently mistreated. *J Vasc Surg* 2009;49(5):1189-95.
 27. O'Leary DP, O'Brien G, Fulton G. Popliteal artery entrapment syndrome. *Int J Surg Case Rep* 2010;1(2):13-5.
 28. McAree BJ, O'Donnell ME, Davison GW et al. Bilateral popliteal artery occlusion in a competitive bike rider: case report and clinical review. *Vasc Endovascular Surg* 2008;42(4):380-5.
 29. di Marzo L, Cavallaro A. Popliteal vascular entrapment. *World J Surg* 2005;29 Suppl 1:S43-5.
 30. Pillai J. A current interpretation of popliteal vascular entrapment. *J Vasc Surg* 2008;48(6 Suppl):61S-65S.
 31. Rignault DP, Pailler JL, Lunel F. The 'functional' popliteal entrapment syndrome. *Int Angiol* 1985;4(3):341-3.
 32. Barbaras AP. Popliteal artery entrapment syndrome. *Br J Hosp Med* 1985;34(5):304.
 33. Altinatis U, Helgstrand U, Hansen M. Popliteal artery entrapment syndrome: ultrasound imaging, intra-operative findings and clinical outcome. *Vasc Endovasc Surg* 2013;47(7):513-8.
 34. Baltopoulos P, Filippou DK, Sigala F. Popliteal artery entrapment syndrome: anatomic or functional syndrome? *Clin J Sport Med* 2004;14(1):8-12.
 35. Willson TD, Revesz E, Podbielski FJ et al. External iliac artery dissection secondary to endofibrosis in a cyclist. *J Vasc Surg* 2010;52(1):219-21.
 36. Smith WB, Olmsted KA, Zierler ER. Iliac Artery Endofibrosis in Athletes: a Case Series. *J Vasc Ultrasound* 2008;32:193-9.
 37. Giannoukas AD, Berczi V, Anoop U et al. Endofibrosis of iliac arteries in high-performance athletes: diagnostic approach and minimally invasive endovascular treatment. *Cardiovasc Intervent Radiol* 2006;29(5):866-9.
 38. Abraham P, Bouye P, Quere I et al. Past, present and future of arterial endofibrosis in athletes: a point of view. *Sports Med* 2004;34(7):419-25.
 39. Fabre T, Montero C, Gaujard E et al. Chronic calf pain in athletes due to sural nerve entrapment. A report of 18 cases. *Am J Sports Med* 2000;28(5):679-82.
 40. Jaureguito JW, Greenwald AE, Wilcox JF et al. The incidence of deep venous thrombosis after arthroscopic knee surgery. *Am J Sports Med* 1999;27(6):707-10.
 41. Wells PS, Hirsh J, Anderson DR et al. Accuracy of clinical assessment of deep-vein thrombosis. *Lancet*, 1995;345(8961):1326-30.

CHAPTER 40

1. Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016;50(5):273-80.
2. van der Worp MP, ten Haaf DS, van Cingel R et al. Injuries in runners; a systematic review on risk factors and sex differences. *PLoS One* 2015;10(2):e0114937.
3. Kujala UM, Sarna S, Kaprio J. Cumulative incidence of Achilles tendon rupture and tendinopathy in male former elite athletes. *Clin J Sport Med* 2005;15(3):133-5.
4. O'Brien M. The anatomy of the Achilles tendon. *Foot Ankle Clin* 2005;10(2):225-38.
5. Wang D, Sandlin MI, Cohen JR et al. Operative versus nonoperative treatment of acute Achilles tendon rupture: an analysis of 12 570 patients in a large healthcare database. *Foot Ankle Surg* 2015;21(4):250-3.
6. Huttunen TT, Kannus P, Rolf C et al. Acute achilles tendon ruptures: incidence of injury and surgery in Sweden between 2001 and 2012. *Am J Sports Med* 2014;42(10):2419-23.
7. Thompson TC, Doherty JH. Spontaneous rupture of tendon of Achilles: a new clinical diagnostic test. *J Trauma* 1962;2:126-9.
8. Simmonds FA. The diagnosis of the ruptured Achilles tendon. *Practitioner* 1957;179(1069):56-8.
9. Hutchison AM, Evans R, Bodger O et al. What is the best clinical test for Achilles tendinopathy? *Foot Ankle Surg* 2013;19(2):112-7.
10. Maffulli N. The clinical diagnosis of subcutaneous tear of the Achilles tendon. A prospective study in 174 patients. *Am J Sports Med* 1998;26(2):266-70.

11. Reiman M, Burgi C, Strube E et al. The utility of clinical measures for the diagnosis of Achilles tendon injuries: a systematic review with meta-analysis. *J Athl Train* 2014;49(6):820-9.
12. Hartgerink P, Fessell DP, Jacobson JA et al. Full- versus partial-thickness Achilles tendon tears: sonographic accuracy and characterization in 26 cases with surgical correlation. *Radiology* 2001;220(2):406-12.
13. Garras DN, Raikin SM, Bhat SB et al. MRI is unnecessary for diagnosing acute Achilles tendon ruptures: clinical diagnostic criteria. *Clin Orthop Relat Res* 2012;470(8):2268-73.
14. Kayser R, Mahlfeld K, Heyde CE. Partial rupture of the proximal Achilles tendon: a differential diagnostic problem in ultrasound imaging. *Br J Sports Med* 2005;39(11):838-42; discussion 842.
15. Khan KM, Forster BB, Robinson J et al. Are ultrasound and magnetic resonance imaging of value in assessment of Achilles tendon disorders? A two year prospective study. *Br J Sports Med* 2003;37(2):149-53.
16. Maffulli N, Kenward MG, Testa V et al. Clinical diagnosis of Achilles tendinopathy with tendinosis. *Clin J Sport Med* 2003;13(1):11-5.
17. Robinson JM, Cook JL, Purdam C et al. The VISA-A questionnaire: a valid and reliable index of the clinical severity of Achilles tendinopathy. *Br J Sports Med* 2001;35(5):335-41.
18. Iversen JV, Bartels EM, Langberg H. The Victorian Institute of Sports Assessment-Achilles questionnaire (VISA-A)-a reliable tool for measuring Achilles tendinopathy. *Int J Sports Phys Ther* 2012;7(1):76-84.
19. de Vos RJ, Weir A, van Schie HT et al. Platelet-rich plasma injection for chronic Achilles tendinopathy: a randomized controlled trial. *JAMA* 2010;303(2):144-9.
20. Sayana MK, Maffulli N. Eccentric calf muscle training in non-athletic patients with Achilles tendinopathy. *J Sci Med Sport* 2007;10(1):52-8.
21. Tumilty S, Munn J, Abbott JH et al. Laser therapy in the treatment of Achilles tendinopathy: a pilot study. *Photomed Laser Surg* 2008;26(1):25-30.
22. Kang S, Thordarson DB, Charlton TP. Insertional Achilles tendinitis and Haglund's deformity. *Foot Ankle Int* 2012;33(6):487-91.
23. Docking SI, Rosengarten SD, Daffy J et al. Structural integrity is decreased in both Achilles tendons in people with unilateral Achilles tendinopathy. *J Sci Med Sport* 2015;18(4):383-7.
24. Giombini A, Dragoni S, Di Cesare A et al. Asymptomatic Achilles, patellar, and quadriceps tendinopathy: a longitudinal clinical and ultrasonographic study in elite fencers. *Scand J Med Sci Sports* 2013;23(3):311-6.
25. Comin J, Cook JL, Malliaras P et al. The prevalence and clinical significance of sonographic tendon abnormalities in asymptomatic ballet dancers: a 24-month longitudinal study. *Br J Sports Med* 2013;47(2):89-92.
26. Åström M, Rausing A. Chronic Achilles tendinopathy. A survey of surgical and histopathologic findings. *Clin Orthop Relat Res* 1995;316:151-64.
27. Asplund CA, Best TM. Achilles tendon disorders. *BMJ* 2013;346:f1262.
28. Kannus P, Józsa L. Histopathological changes preceding spontaneous rupture of a tendon. A controlled study of 891 patients. *J Bone Joint Surg Am* 1991;73(10):1507-25.
29. Åström M, Gentz CF, Nilsson P et al. Imaging in chronic Achilles tendinopathy: a comparison of ultrasonography, magnetic resonance imaging and surgical findings in 27 histologically verified cases. *Skeletal Radiol* 1996;25(7):615-20.
30. Vasta S, Di Martino A, Zampogna B et al. Role of VEGF, nitric oxide, and sympathetic neurotransmitters in the pathogenesis of tendinopathy: a review of the current evidences. *Front Aging Neurosci* 2016;8:186.
31. Alfredson H, Forsgren S, Thorsen K et al. Glutamate NMDAR1 receptors localised to nerves in human Achilles tendons. Implications for treatment? *Knee Surg Sports Traumatol Arthrosc* 2001;9(2):123-6.
32. Alfredson H, Thorsen K, Lorentzon R. In situ microdialysis in tendon tissue: high levels of glutamate, but not prostaglandin E2 in chronic Achilles tendon pain. *Knee Surg Sports Traumatol Arthrosc* 1999;7(6):378-81.
33. Alfredson H, Lorentzon M, Bäckman S et al. cDNA-arrays and real-time quantitative PCR techniques in the investigation of chronic Achilles tendinosis. *J Orthop Res* 2003;21(6):970-5.
34. Magnan B, Bondi M, Pierantoni S et al. The pathogenesis of Achilles tendinopathy: a systematic review. *Foot Ankle Surg* 2014;20(3):154-9.
35. Rees JD, Stride M, Scott A. Tendons-time to revisit inflammation. *Br J Sports Med* 2014;48(21):1553-7.
36. Scott A, Khan KM, Cook JL et al. What is 'inflammation'? Are we ready to move beyond Celsus? *Br J Sports Med* 2004;38(3):248-9.
37. Danielson P. Reviving the 'biochemical' hypothesis for tendinopathy: new findings suggest the involvement of locally produced signal substances. *Br J Sports Med* 2009;43(4):265-8.
38. Rio E, Moseley L, Purdam C et al. The pain of tendinopathy: physiological or pathophysiological? *Sports Med* 2014;44(1):9-23.
39. Gaida JE, Alfredson H, Kiss ZS et al. Asymptomatic Achilles tendon pathology is associated with a central fat distribution in men and a peripheral fat distribution in women: a cross sectional study of 298 individuals. *BMC Musculoskelet Disord* 2010;11:41.
40. Klein EE, 'Weil L, Jr., Weil LS, Sr.' et al. Body mass index and Achilles tendonitis: a 10-year retrospective analysis. *Foot Ankle Spec* 2013;6(4):276-82.
41. Scott RT, Hyer CF, Granata A. The correlation of Achilles tendinopathy and body mass index. *Foot Ankle Spec* 2013;6(4):283-5.
42. Batista F, Nery C, Pinzur M et al. Achilles tendinopathy in diabetes mellitus. *Foot Ankle Int* 2008;29(5):498-501.
43. Rees J, Gaida JE, Silbernagel KG et al. Rehabilitation of tendon problems in patients with diabetes mellitus. *Adv Exp Med Biol* 2016;920:199-208.

44. Saragiotto BT, Yamato TP, Hespanhol Junior LC et al. What are the main risk factors for running-related injuries? *Sports Med* 2014;44(8):1153-63.
45. van Gent RN, Siem D, van Middelkoop M et al. Incidence and determinants of lower extremity running injuries in long distance runners: a systematic review. *Br J Sports Med* 2007;41(8):469-80; discussion 480.
46. Posthumus M, September AV, Schwellnus MP et al. Investigation of the Sp1-binding site polymorphism within the COL1A1 gene in participants with Achilles tendon injuries and controls. *J Sci Med Sport* 2009;12(1):184-9.
47. Raleigh SM, van der Merwe L, Ribbans WJ et al. Variants within the MMP3 gene are associated with Achilles tendinopathy: possible interaction with the COL5A1 gene. *Br J Sports Med* 2009;43(7):514-20.
48. Lewis T, Cook J. Fluoroquinolones and tendinopathy: a guide for athletes and sports clinicians and a systematic review of the literature. *J Athl Train* 2014;49(3):422-7.
49. Kirchgessner T, Larbi A, Omoumi P et al. Drug-induced tendinopathy: from physiology to clinical applications. *Joint Bone Spine* 2014;81(6):485-92.
50. Gerster JC. Plantar fasciitis and Achilles tendinitis among 150 cases of seronegative spondylarthritis. *Rheumatol Rehabil* 1980;19(4):218-22.
51. Longo UG, Ronga M, Maffulli N. Achilles tendinopathy. *Sports Med Arthrosc* 2009;17(2):112-26.
52. Munteanu SE, Barton CJ. Lower limb biomechanics during running in individuals with Achilles tendinopathy: a systematic review. *J Foot Ankle Res* 2011;4:15.
53. Mahieu NN, Witvrouw E, Stevens V et al. Intrinsic risk factors for the development of Achilles tendon overuse injury: a prospective study. *Am J Sports Med* 2006;34(2):226-35.
54. Kaufman KR, Brodine SK, Shaffer RA et al. The effect of foot structure and range of motion on musculoskeletal overuse injuries. *Am J Sports Med* 1999;27(5):585-93.
55. Van Ginckel A, Thijs Y, Hesar NG et al. Intrinsic gait-related risk factors for Achilles tendinopathy in novice runners: a prospective study. *Gait Posture* 2009;29(3):387-91.
56. Rabin A, Kozol Z, Finestone AS. Limited ankle dorsiflexion increases the risk for mid-portion Achilles tendinopathy in infantry recruits: a prospective cohort study. *J Foot Ankle Res* 2014;7(1):48.
57. Hein T, Janssen P, Wagner-Fritz U et al. Prospective analysis of intrinsic and extrinsic risk factors on the development of Achilles tendon pain in runners. *Scand J Med Sci Sports* 2014;e24(3):201-12.
58. Freckleton G, Cook J, Pizzari T. The predictive validity of a single leg bridge test for hamstring injuries in Australian Rules football players. *Br J Sports Med* 2014;48(8):713-7.
59. Fredberg U, Bolvig L, Andersen NT. Prophylactic training in asymptomatic soccer players with ultrasonographic abnormalities in Achilles and patellar tendons: the Danish Super League study. *Am J Sports Med* 2008;36(3):451-60.
60. Counsel P, Comin J, Davenport M et al. Pattern of fascicular involvement in midportion Achilles tendinopathy at ultrasound. *Sports Health* 2015;7(5):424-8.
61. Marti B, Vader JP, Minder CE et al. On the epidemiology of running injuries. The 1984 Bern Grand-Prix study. *Am J Sports Med* 1988;16(3):285-94.
62. Nielsen RØ, Parner ET, Nohr EA et al. Excessive progression in weekly running distance and risk of running-related injuries: an association which varies according to type of injury. *J Orthop Sports Phys Ther* 2014;44(10):739-47.
63. Orchard JW, Blanch P, Paoloni J et al. Cricket fast bowling workload patterns as risk factors for tendon, muscle, bone and joint injuries. 2015;49(16):1064-8.
64. Windt J, Gabbett TJ. How do training and competition workloads relate to injury? The workload-injury aetiology model. *Br J Sports Med* 2016 Jul 14. doi: 10.1136/bjsports-2016-096040.
65. Khan KM, Scott A. Mechanotherapy: how physical therapists' prescription of exercise promotes tissue repair. *Br J Sports Med* 2009;43(4):247-52.
66. Alfredson H, Pietilä T, Jonsson P et al. Heavy-load eccentric calf muscle training for the treatment of chronic Achilles tendinosis. *Am J Sports Med* 1998;26(3):360-6.
67. Shalabi A, Kristoffersen-Wilberg M, Svensson L et al. Eccentric training of the gastrocnemius-soleus complex in chronic Achilles tendinopathy results in decreased tendon volume and intratendinous signal as evaluated by MRI. *Am J Sports Med* 2004;32(5):1286-96.
68. Silbernagel KG, Thomeé R, Thomeé P et al. Eccentric overload training for patients with chronic Achilles tendon pain—a randomised controlled study with reliability testing of the evaluation methods. *Scand J Med Sci Sports* 2001;11(4):197-206.
69. Malliaras P, Barton CJ, Reeves ND et al. Achilles and patellar tendinopathy loading programmes: a systematic review comparing clinical outcomes and identifying potential mechanisms for effectiveness. *Sports Med* 2013;43(4):267-86.
70. Kongsgaard M, Kovanen V, Aagaard P et al. Corticosteroid injections, eccentric decline squat training and heavy slow resistance training in patellar tendinopathy. *Scand J Med Sci Sports* 2009; 19(6):790-802.
71. Frohm A, Saartok T, Halvorsen K et al. Eccentric treatment for patellar tendinopathy: a prospective randomised short-term pilot study of two rehabilitation protocols. *Br J Sports Med* 2007;41(7):e7.
72. Allison GT, Purdam C. Eccentric loading for Achilles tendinopathy—strengthening or stretching? *Br J Sports Med* 2009;43(4):276-9.
73. Silbernagel KG, Thomeé R, Eriksson BI et al. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomized controlled study. *Am J Sports Med* 2007;35(6):897-906.
74. Cook JL, Docking SI. 'Rehabilitation will increase the "capacity" of your . . . insert musculoskeletal tissue here. . . ' Defining

- 'tissue capacity': a core concept for clinicians. *Br J Sports Med* 2015;49(23):1484-5.
75. Rio E, Kidgell D, Moseley GL et al. Tendon neuroplastic training: changing the way we think about tendon rehabilitation: a narrative review. *Br J Sports Med* 2016;50(4):209-15.
 76. Rio E, Kidgell D, Purdam C et al. Isometric exercise induces analgesia and reduces inhibition in patellar tendinopathy. *Br J Sports Med* 2015;49(19):1277-83.
 77. Masci L, Spang C, van Schie HT et al. How to diagnose plantaris tendon involvement in midportion Achilles tendinopathy—clinical and imaging findings. *BMC Musculoskelet Disord* 2016;17:97.
 78. Spang C, Alfredson H, Ferguson M et al. The plantaris tendon in association with mid-portion Achilles tendinosis: tendinosis-like morphological features and presence of a non-neuronal cholinergic system. *Histol Histopathol* 2013;28(5):623-32.
 79. Spang C, Harandi VM, Alfredson H et al. Marked innervation but also signs of nerve degeneration in between the Achilles and plantaris tendons and presence of innervation within the plantaris tendon in midportion Achilles tendinopathy. *J Musculoskelet Neuronal Interact* 2015;15(2):197-206.
 80. Alfredson H. Midportion Achilles tendinosis and the plantaris tendon. *Br J Sports Med* 2011;45(13):1023-5.
 81. Bedi HS, Jowett C, Ristanis S et al. Plantaris Excision and Ventral Paratendinous Scraping for Achilles Tendinopathy in an Athletic Population. *Foot Ankle Int* 2016;37(4):386-93.
 82. Maffulli N, Walley G, Sayana MK et al. Eccentric calf muscle training in athletic patients with Achilles tendinopathy. *Disabil Rehabil* 2008;30(20-22):1677-84.
 83. Alfredson H, Ohberg L. Sclerosing injections to areas of neo-vascularisation reduce pain in chronic Achilles tendinopathy: a double-blind randomised controlled trial. *Knee Surg Sports Traumatol Arthrosc* 2005;13(4):338-44.
 84. Lind B, Ohberg L, Alfredson H. Sclerosing polidocanol injections in mid-portion Achilles tendinosis: remaining good clinical results and decreased tendon thickness at 2-year follow-up. *Knee Surg Sports Traumatol Arthrosc* 2006;14(12):1327-32.
 85. Kearney RS, Parsons N, Metcalfe D et al. Injection therapies for Achilles tendinopathy. *Cochrane Database Syst Rev* 2015(5):Cd010960.
 86. Coombes BK, Bisset L, Vicenzino B. Efficacy and safety of corticosteroid injections and other injections for management of tendinopathy: a systematic review of randomised controlled trials. *Lancet* 2010;376(9754):1751-67.
 87. Maffulli N, Spiezia F, Longo UG et al. High volume image guided injections for the management of chronic tendinopathy of the main body of the Achilles tendon. *Phys Ther Sport* 2013;14(3):163-7.
 88. Chen J, Yu Q, Wu B et al. Autologous tenocyte therapy for experimental Achilles tendinopathy in a rabbit model. *Tissue Eng Part A* 2011;17(15-16):2037-48.
 89. Mani-Babu S, Morrissey D, Waugh C et al. The effectiveness of extracorporeal shock wave therapy in lower limb tendinopathy: a systematic review. *Am J Sports Med* 2015;43(3):752-61.
 90. Rasmussen S, Christensen M, Mathiesen I et al. Shockwave therapy for chronic Achilles tendinopathy: a double-blind, randomized clinical trial of efficacy. *Acta Orthop* 2008; 79(2):249-56.
 91. Stergioulas A, Stergioula M, Aarskog R et al. Effects of low-level laser therapy and eccentric exercises in the treatment of recreational athletes with chronic Achilles tendinopathy. *Am J Sports Med* 2008;36(5):881-7.
 92. Bjordal JM, Lopes-Martins RA, Iversen VV. A randomised, placebo controlled trial of low level laser therapy for activated Achilles tendinitis with microdialysis measurement of peritendinous prostaglandin E2 concentrations. *Br J Sports Med* 2006;40(1):80.
 93. Alfredson H. Ultrasound and Doppler-guided mini-surgery to treat midportion Achilles tendinosis: results of a large material and a randomised study comparing two scraping techniques. *Br J Sports Med* 2011;45(5):407-10.
 94. Alfredson H, Spang C, Forsgren S. Unilateral surgical treatment for patients with midportion Achilles tendinopathy may result in bilateral recovery. *Br J Sports Med* 2014;48(19):1421-4.
 95. Alfredson H, Ohberg L, Zeisig E et al. Treatment of midportion Achilles tendinosis: similar clinical results with US and CD-guided surgery outside the tendon and sclerosing polidocanol injections. *Knee Surg Sports Traumatol Arthrosc* 2007;15(12):1504-9.
 96. Scott A, Khan KM. Corticosteroids: short-term gain for long-term pain? *Lancet* 2010;376(9754):1714-5.
 97. de Mos M, van der Windt AE, Jahr H et al. Can platelet-rich plasma enhance tendon repair? A cell culture study. *Am J Sports Med* 2008;36(6):1171-8.
 98. Jo CH, Kim JE, Yoon KS et al. Platelet-rich plasma stimulates cell proliferation and enhances matrix gene expression and synthesis in tenocytes from human rotator cuff tendons with degenerative tears. *Am J Sports Med* 2012;40(5):1035-45.
 99. Kearney RS, Parsons N, Costa ML. Achilles tendinopathy management: a pilot randomised controlled trial comparing platelet-rich plasma injection with an eccentric loading programme. *Bone Joint Res* 2013;2(10):227-32.
 100. de Vos RJ, Weir A, Tol JL et al. No effects of PRP on ultrasonographic tendon structure and neovascularisation in chronic midportion Achilles tendinopathy. *Br J Sports Med* 2011;45(5):387-92.
 101. Andersson G, Backman LJ, Christensen J et al. Nerve distributions in insertional Achilles tendinopathy—a comparison of bone, bursae and tendon. *Histol Histopathol* 2016 Jun 20. doi:10.14670/HH-11-790. .
 102. McLauchlan GJ, Handoll HH. Interventions for treating acute and chronic Achilles tendinitis. *Cochrane Database Syst Rev* 2001(2):CD000232.
 103. Wheeler PC, Mahadevan D, Bhatt R et al. A Comparison of Two Different High-Volume Image-Guided Injection Procedures for Patients With Chronic Noninsertional Achilles Tendinopathy: A Pragmatic Retrospective Cohort Study. *J Foot Ankle Surg.* 2016 Sep-Oct;55(5):976-9.

104. Muller SA, Todorov A, Heisterbach PE et al. Tendon healing: an overview of physiology, biology, and pathology of tendon healing and systematic review of state of the art in tendon bioengineering. *Knee Surg Sports Traumatol Arthrosc* 2015;23(7):2097-105.
105. Boudreault J, Desmeules F, Roy JS et al. The efficacy of oral non-steroidal anti-inflammatory drugs for rotator cuff tendinopathy: a systematic review and meta-analysis. *J Rehabil Med* 2014;46(4):294-306.
106. Matava M, Brater DC, Gritter N et al. Recommendations of the National Football League Physician Society task force on the use of Toradol® ketorolac in the National Football League. *Sports Health* 2012;4(5):377-83.
107. Rompe JD, Furia J, Maffulli N. Eccentric loading versus eccentric loading plus shock-wave treatment for midportion Achilles tendinopathy: a randomized controlled trial. *Am J Sports Med* 2009;37(3):463-70.
108. Huisstede BM, Gebremariam L, van der Sande R et al. Evidence for effectiveness of extracorporeal shock-wave therapy (ESWT) to treat calcific and non-calcific rotator cuff tendinosis—a systematic review. *Man Ther* 2011;16(5):419-33.
109. van der Worp H, van den Akker-Scheek I, van Schie H et al. ESWT for tendinopathy: technology and clinical implications. *Knee Surg Sports Traumatol Arthrosc* 2013;21(6):1451-8.
110. Bosch G, de Mos M, van Binsbergen R et al. The effect of focused extracorporeal shock wave therapy on collagen matrix and gene expression in normal tendons and ligaments. *Equine Vet J* 2009;41(4):335-41.
111. Baltes TP, Zwiers R, Wiegerinck JI et al. Surgical treatment for midportion Achilles tendinopathy: a systematic review. *Knee Surg Sports Traumatol Arthrosc* 2016 Mar 12. doi: 10.1007/s00167-016-4062-9. [Epub ahead of print]
112. Cook JL, Purdam C. Is compressive load a factor in the development of tendinopathy? *Br J Sports Med* 2012; 46(3):163-8.
113. Rufai A, Ralphs JR, Benjamin M. Structure and histopathology of the insertional region of the human Achilles tendon. *J Orthop Res* 1995;13(4):585-93.
114. Lyman J, Weinhold PS, Almekinders LC. Strain behavior of the distal Achilles tendon: implications for insertional Achilles tendinopathy. *Am J Sports Med* 2004;32(2):457-61.
115. Haglund P. Beitrag zur Klinik der Achillessehne. *Zeitschr Orthop Chir* 1928;49:49-58.
116. van Dijk CN, van Sterkenburg MN, Wiegerinck JI et al. Terminology for Achilles tendon related disorders. *Knee Surg Sports Traumatol Arthrosc* 2011;19(5):835-41.
117. Benjamin M, Ralphs JR. Tendons and ligaments—an overview. *Histol Histopathol* 1997;12(4):1135-44.
118. Benjamin M, Ralphs JR. Fibrocartilage in tendons and ligaments—an adaptation to compressive load. *J Anat* 1998;193 (Pt 4):481-94.
119. Jonsson P, Alfredson H, Sunding K et al. New regimen for eccentric calf-muscle training in patients with chronic insertional Achilles tendinopathy: results of a pilot study. *Br J Sports Med* 2008;42(9):746-9.
120. Hoksrud A, Torgalsen T, Harstad H et al. Ultrasound-guided sclerosis of neovessels in patellar tendinopathy: a prospective study of 101 patients. *Am J Sports Med* 2012;40(3):542-7.
121. Ohberg L, Alfredson H. Sclerosing therapy in chronic Achilles tendon insertional pain—results of a pilot study. *Knee Surg Sports Traumatol Arthrosc* 2003;11(5):339-43.
122. Rompe JD, Furia J, Maffulli N. Eccentric loading compared with shock wave treatment for chronic insertional Achilles tendinopathy. A randomized, controlled trial. *J Bone Joint Surg Am* 2008;90(1):52-61.
123. Nesse E, Finsen V. Poor results after resection for Haglund's heel: analysis of 35 heels in 23 patients after 3 years. *Acta Orthop Scand* 1994;65(1):107-9.
124. Fridrich F. [Tendon-splitting approach for the surgical treatment of Haglund's deformity and associated condition. Evaluation and results]. *Acta Chir Orthop Traumatol Cech* 2009;76(3):212-7.
125. DeVries JG, Summerhays B, Guehlstorf DW. Surgical correction of Haglund's triad using complete detachment and reattachment of the Achilles tendon. *Journal of Foot & Ankle Surgery* 2009;48(4):447-51.
126. Anderson JA, Suero E, O'Loughlin PF et al. Surgery for retrocalcaneal bursitis: a tendon-splitting versus a lateral approach. *Clin Orthop Relat Res* 2008;466(7):1678-82.
127. Stephens MM. Haglund's deformity and retrocalcaneal bursitis. *Orthop Clin North Am* 1994;25(1):41-6.
128. Goldberg-Stein S, Berko N, Thornhill B et al. Fluoroscopically guided retrocalcaneal bursa steroid injection: description of the technique and pilot study of short-term patient outcomes. *Skeletal Radiol* 2016;45(8):1107-12.
129. Vallone G, Vittorio T. Complete Achilles tendon rupture after local infiltration of corticosteroids in the treatment of deep retrocalcaneal bursitis. *J Ultrasound* 2014;17(2):165-7.
130. Wiegerinck JI, Kok AC, van Dijk CN. Surgical treatment of chronic retrocalcaneal bursitis. *Arthroscopy* 2012;28(2):283-93.
131. Mukish P, Reybet-Degat PY, Demangel A et al. Accessory soleus muscle: a difficult diagnosis: a case report and a review of the literature. *Eur J Orthop Surg Traumatol* 2012;22 Suppl 1:205-9.
132. Del Nero FB, Ruiz CR, Aliaga Júnior R. The presence of accessory soleus muscle in humans. *Einstein (Sao Paulo)* 2012;10(1):79-81.
133. Isner-Horobeti M-E, Muff G, Lonsdorfer-Wolf E et al. Use of botulinum toxin type A in symptomatic accessory soleus muscle: first five cases. *Scand J Med Sci Sports* 2015 Dec 2. doi: 10.1111/sms.12616.
134. Leswick DA, Chow V, Stoneham GW. Resident's corner. Answer to case of the month #94. Accessory soleus muscle. *Can Assoc Radiol J* 2003;54(5):313-5.
135. Zellers JA, Carmont MR, Grävare Silbernagel K. Return to play post-Achilles tendon rupture: a systematic review and meta-analysis of rate and measures of return to play. *Br J Sports Med* 2016 Jun 3. pii: bjsports-2016-096106.

136. Claessen FM, de Vos RJ, Reijman M et al. Predictors of primary Achilles tendon ruptures. *Sports Med* 2014;44(9):1241-59.
137. Carmont MR, Silbernagel KG, Nilsson-Helander K et al. Cross cultural adaptation of the Achilles tendon total rupture score with reliability, validity and responsiveness evaluation. *Knee Surg Sports Traumatol Arthrosc* 2013;21(6):1356-60.
138. Silbernagel KG, Nilsson-Helander K, Thomeé R et al. A new measurement of heel-rise endurance with the ability to detect functional deficits in patients with Achilles tendon rupture. *Knee Surg Sports Traumatol Arthrosc* 2010;18(2):258-64.
139. Brorsson A, Olsson N, Nilsson-Helander K et al. Recovery of calf muscle endurance 3 months after an Achilles tendon rupture. *Scand J Med Sci Sports* 2016;26(7):844-53.
140. Nilsson-Helander K, Thomeé R, Silbernagel KG et al. The Achilles tendon total rupture score (ATRS): development and validation. *Am J Sports Med* 2007;35(3):421-6.
141. Olsson N, Karlsson J, Eriksson BI et al. Ability to perform a single heel-rise is significantly related to patient-reported outcome after Achilles tendon rupture. *Scand J Med Sci Sports* 2014;24(1):152-8.
142. Olsson N, Petzold M, Brorsson A et al. Predictors of clinical outcome after acute Achilles tendon ruptures. *Am J Sports Med* 2014;42(6):1448-55.
143. Silbernagel KG, Brorsson A, Olsson N et al. Sex differences in outcome after an acute Achilles tendon rupture. *Orthop J Sports Med* 2015;3(6):2325967115586768.
144. Khan RJ, Fick D, Keogh A et al. Treatment of acute Achilles tendon ruptures. A meta-analysis of randomized, controlled trials. *J Bone Joint Surg Am* 2005;87A(10):2202-10.
145. Paavola M, Orava S, Leppilahti J et al. Chronic Achilles tendon overuse injury: complications after surgical treatment. An analysis of 432 consecutive patients. *Am J Sports Med* 2000;28(1):77-82.
146. Huang J, Wang C, Ma X et al. Rehabilitation regimen after surgical treatment of acute Achilles tendon ruptures: a systematic review with meta-analysis. *Am J Sports Med* 2015;43(4):1008-16.
147. Olsson N, Silbernagel KG, Eriksson BI et al. Stable surgical repair with accelerated rehabilitation versus nonsurgical treatment for acute Achilles tendon ruptures: a randomized controlled study. *Am J Sports Med* 2013;41(12):2867-76.
148. Nilsson-Helander K, Silbernagel KG, Thomeé R et al. Acute Achilles tendon rupture: a randomized, controlled study comparing surgical and nonsurgical treatments using validated outcome measures. *Am J Sports Med* 2010;38(11):2186-93.
149. Lunsford BR, Perry J. The standing heel-rise test for ankle plantar flexion: criterion for normal. *Phys Ther* 1995;75(8):694-8.
2. Bahr R, Bahr IA. Incidence of acute volleyball injuries: a prospective cohort study of injury mechanisms and risk factors. *Scand J Med Sci Sports* 1997;7(3):166-71.
3. Morrison KE, Kaminski TW. Foot characteristics in association with inversion ankle injury. *J Athl Train* 2007;42(1):135-42.
4. Jonckheer P, Willems T, De Ridder R et al. Evaluating fracture risk in acute ankle sprains: any news since the Ottawa ankle rules? A systematic review. *Eur J Gen Pract* 2016;22(1):31-41.
5. Lin CW, Uegaki K, Coupe VM et al. Economic evaluations of diagnostic tests, treatment and prevention for lateral ankle sprains: a systematic review. *Br J Sports Med* 2013;47(18):114-9 doi: 10.1136/bjsports-2012.
6. Dimmick S, Kennedy D, Daunt N. Evaluation of thickness and appearance of anterior talofibular and calcaneofibular ligaments in normal versus abnormal ankles with MRI. *J Med Imaging Radiat Oncol* 2008;52(6):559-63.
7. Cook CE, Hegedus EJ. *Orthopedic Physical Examination Tests*. 2nd ed. Upper Saddle River, New Jersey: Pearson Education Inc, 2013.
8. Jain N, Symes T, Doorgakant A et al. Clinical audit of the management of stable ankle fractures. *Ann R Coll Surg Engl* 2008;90(6):483-7.
9. Holme E, Magnusson SP, Becher K et al. The effect of supervised rehabilitation on strength, postural sway, position sense and re-injury risk after acute ankle ligament sprain. *Scand J Med Sci Sports* 1999;9(2):104-9.
10. Verhagen E, van der Beek A, Twisk J et al. The effect of a proprioceptive balance board training program for the prevention of ankle sprains: a prospective controlled trial. *Am J Sports Med* 2004;32(6):1385-93. doi: 10.1177/0363546503262177.
11. Verhagen EA, van Tulder M, van der Beek AJ et al. An economic evaluation of a proprioceptive balance board training programme for the prevention of ankle sprains in volleyball. *Br J Sports Med* 2005;39(2):111-5.
12. Beynnon BD, Renstrom PA, Alosa DM et al. Ankle ligament injury risk factors: a prospective study of college athletes. *J Orthop Res* 2001;19(2):213-20.
13. Verhagen EA, Van der Beek AJ, Bouter LM et al. A one season prospective cohort study of volleyball injuries. *Br J Sports Med* 2004;38(4):477-81.
14. Verhagen EA, Bay K. Optimising ankle sprain prevention: a critical review and practical appraisal of the literature. *Br J Sports Med* 2010;44(15):1082-8.
15. McKeon PO, Hertel J. Systematic review of postural control and lateral ankle instability, part II: is balance training clinically effective? *J Athl Train* 2008;43(3):305-15.
16. Abian-Vicen J, Alegre LM, Fernandez-Rodriguez JM et al. Prophylactic ankle taping: elastic versus inelastic taping. *Foot Ankle Int* 2009;30(3):218-25.
17. Hupperets MD, Verhagen EA, van Mechelen W. Effect of sensorimotor training on morphological, neurophysiological and functional characteristics of the ankle: a critical review. *Sports Med* 2009;39(7):591-605.

CHAPTER 41

1. Doherty C, Delahunt E, Caulfield B et al. The incidence and prevalence of ankle sprain injury: a systematic review and meta-analysis of prospective epidemiological studies. *Sports Med* 2014;44(1):123-40.

18. Verhagen EA, van Mechelen W, de Vente W. The effect of preventive measures on the incidence of ankle sprains. *Clin J Sport Med* 2000;10(4):291-6.
19. Verhagen EA, van der Beek AJ, van Mechelen W. The effect of tape, braces and shoes on ankle range of motion. *Sports Med* 2001;31(9):667-77.
20. Mason-Mackay AR, Whatman C, Reid D. The effect of ankle bracing on lower extremity biomechanics during landing: a systematic review. *J Sci Med Sport* 2016;19(7):531-40.
21. Postle K, Pak D, Smith TO. Effectiveness of proprioceptive exercises for ankle ligament injury in adults: a systematic literature and meta-analysis. *Man Ther* 2012;17(4):285-91.
22. Karlsson J, Lansinger O. Chronic lateral instability of the ankle in athletes. *Sports Med* 1993;16(5):355-65.
23. Lephart S, Conley K. The role of proprioception in chronic ankle instability. In: Schmidt R, Benesch S, Lipke K., eds. *Chronic Ankle Instability: Manuscripts of the International Ankle Symposium*. Ulm: Libri Verlag, 2000:254-67.
24. Hupperets MD, Verhagen EA, van Mechelen W. Effect of unsupervised home based proprioceptive training on recurrences of ankle sprain: randomised controlled trial. *BMJ* 2009;339:b2684. doi: 10.1136/bmj.b2684.
25. Kerkhoffs GM, Handoll HH, de Bie R et al. Surgical versus conservative treatment for acute injuries of the lateral ligament complex of the ankle in adults. *Cochrane Database Syst Rev* 2002(3):CD000380.
26. Petersen W, Rembitzki IV, Koppenburg AG et al. Treatment of acute ankle ligament injuries: a systematic review. *Arch Orthop Trauma Surg* 2013;133(8):1129-41.
27. Kaikkonen A, Kannus P, Jarvinen M. Surgery versus functional treatment in ankle ligament tears. A prospective study. *Clin Orthop Relat Res* 1996;326:194-202.
28. Pijnenburg ACM, Bogaard K, Krips R et al. Operative and functional treatment of rupture of the lateral ligament of the ankle: a randomised, prospective trial. *J Bone Joint Surg Br*, 2003;85B(4):525-30.
29. Babis GC, Papagelopoulos PJ, Tsarouchas J et al. Operative treatment for Maisonneuve fracture of the proximal fibula. *Orthopedics* 2000;23(7):687-90.
30. Wikstrom EA, Naik S, Lodha N et al. Bilateral balance impairments after lateral ankle trauma: a systematic review and meta-analysis. *Gait & Posture* 2010;31(4):407-14.
31. Lee M, Kwon JW, Choi WJ et al. Comparison of outcomes for osteochondral lesions of the talus with and without chronic lateral ankle instability. *Foot Ankle Int* 2015;36(9):1050-7.
32. Rolf C, Guntner P, Ekenman I et al. Dislocation of the tibialis posterior tendon: diagnosis and treatment. *J Foot Ankle Surg* 1997;36(1):63-5.
33. van Dijk CN, Longo UG, Loppini M et al. Classification and diagnosis of acute isolated syndesmotic injuries: ESSKA-AFAS consensus and guidelines. *Knee Surg Sports Traumatol Arthrosc*. 2016;24(4):1200-16.
34. Ogilvie-Harris DJ, Reed SC, Hedman TP. Disruption of the ankle syndesmosis: biomechanical study of the ligamentous restraints. *Arthroscopy* 1994;19(5):558-60.
35. Gerber JP, Williams GN, Scoville CR et al. Persistent disability associated with ankle sprains: a prospective examination of an athletic population. *Foot Ankle Int* 1998;19(10):653-60.
36. Beumer A. Chronic instability of the anterior syndesmosis of the ankle. *Acta Orthop Suppl* 2007;78(327):4-36.
37. Oae K, Takao M, Naito K et al. Injury of the tibiofibular syndesmosis: value of MR imaging for diagnosis. *Radiology* 2003;227(1):155-61.
38. Shehab D, Elgazzar A, Collier BD et al. Impact of three-phase bone scintigraphy on the diagnosis and treatment of complex regional pain syndrome type I or reflex sympathetic dystrophy. *Med Princ Pract* 2006;15(1):46-51.
39. Forouzanfar T, Koke AJ, van Kleef M et al. Treatment of complex regional pain syndrome type I. *Eur J Pain* 2002;6(2):105-22.
40. Oerlemans HM, Oostendorp RA, de Boo T et al. Adjuvant physical therapy versus occupational therapy in patients with reflex sympathetic dystrophy/complex regional pain syndrome type I. *Arch Phys Med Rehabil* 2000;81(1):49-56.
41. van de Vusse AC, Stomp-van den Berg SG, Kessels AH et al. Randomised controlled trial of gabapentin in complex regional pain syndrome type 1 [ISRCTN84121379]. *BMC Neurol* 2004;4:13.
42. Manicourt DH, Brasseur JP, Boutsen Y et al. Role of alendronate in therapy for posttraumatic complex regional pain syndrome type I of the lower extremity. *Arthritis Rheum* 2004;50(11):3690-7.
43. Cepeda MS, Carr DB, Lau J. Local anesthetic sympathetic blockade for complex regional pain syndrome. *Cochrane Database Syst Rev* 2005(4):CD004598.

CHAPTER 42

1. Martin RL, Irrgang JJ, Burdett RG et al. Evidence of validity for the Foot and Ankle Ability Measure (FAAM). *Foot Ankle Int* 2005;26(11):968-83.
2. Carcia CR, Martin RL, Drouin JM. Validity of the Foot and Ankle Ability Measure in athletes with chronic ankle instability. *J Athl Train* 2008;43(2):179-83.
3. Moreira TS, Magalhaes LC, Silva RD et al. Translation, cross-cultural adaptation and validity of the Brazilian version of the Foot and Ankle Ability Measure questionnaire. *Disabil Rehabil* 2016;38(25):2479-90.
4. Weel H, Zwiers R, Azim D et al. Validity and reliability of a Dutch version of the Foot and Ankle Ability Measure. *Knee Surg Sports Traumatol Arthrosc* 2016;24(4):1348-54.
5. Nauck T, Lohrer H. Translation, cross-cultural adaption and validation of the German version of the Foot and Ankle Ability Measure for patients with chronic ankle instability. *Br J Sports Med* 2011;45(10):785-90.
6. Sartorio F, Vercelli S, Bravini E et al. Foot and Ankle Ability Measure: cross-cultural translation and validation of the

- Italian version of the ADL module (FAAM-I/ADL). *Med Lav* 2014;105(5):357-65.
7. Uematsu D, Suzuki H, Sasaki S et al. Evidence of validity for the Japanese version of the Foot and Ankle Ability Measure. *J Athl Train* 2015;50(1):65-70.
 8. Kim JA, Ha S, Kang T et al. Reliability and validity of Korean version Ankle Instability Instrument and Foot and Ankle Ability Measure: 2782 Board #305 June 3, 9:30 AM-11:00 AM. *Med Sci Sports Exerc* 2016;48(5 Suppl 1):783.
 9. Mazaheri M, Salavati M, Negahban H et al. Reliability and validity of the Persian version of Foot and Ankle Ability Measure (FAAM) to measure functional limitations in patients with foot and ankle disorders. *Osteoarth Cart* 2010;18(6):755-9.
 10. Arunakul M, Arunakul P, Suesiritumrong C et al. Validity and reliability of Thai version of the Foot and Ankle Ability Measure (FAAM) subjective form. *J Med Assoc Thai* 2015;98(6):561-7.
 11. Celik D, Malkoc M, Martin R. Evidence for reliability, validity and responsiveness of Turkish Foot and Ankle Ability Measure (FAAM). *Rheumatol Int* 2016 May 2. [Epub ahead of print]
 12. Morris C, Doll HA, Wainwright A et al. The Oxford Ankle Foot Questionnaire for children: scaling, reliability and validity. *J Bone Joint Surg Br* 2008;90(11):1451-6.
 13. Morris C, Doll H, Davies N et al. The Oxford Ankle Foot Questionnaire for children: responsiveness and longitudinal validity. *Qual Life Res* 2009;18(10):1367-76.
 14. Morris C, Liabo K, Wright P et al. Development of the Oxford Ankle Foot Questionnaire: finding out how children are affected by foot and ankle problems. *Child Care Health Dev* 2007;33(5):559-68.
 15. Martinkevich P, Moller-Madsen B, Gottlieb M et al. Validation of the translated Oxford Ankle Foot Questionnaire in 82 Danish children aged between five and 16 years. *Bone Joint J* 2015;97-b(3):420-6.
 16. Martinelli N, Romeo G, Bonifacini C et al. Validation of the Italian version of the Oxford Ankle Foot Questionnaire for children. *Qual Life Res* 2016;25(1):117-23.
 17. Hale SA, Hertel J. Reliability and sensitivity of the Foot and Ankle Disability Index in subjects with chronic ankle instability. *J Athl Train* 2005;40(1):35-40.
 18. Martin R, Burdett R, Irrgang J. Development of the Foot and Ankle Disability Index (FADI) [abstract]. *J Orthop Sports Phys Ther* 1999;29:A32-3.
 19. Johanson NA, Liang MH, Daltroy L et al. American Academy of Orthopaedic Surgeons lower limb outcomes assessment instruments. Reliability, validity, and sensitivity to change. *J Bone Joint Surg Am* 2004;86-a(5):902-9.
 20. Goldstein CL, Schemitsch E, Bhandari M et al. Comparison of different outcome instruments following foot and ankle trauma. *Foot Ankle Int* 2010;31(12):1075-80.
 21. Gonzalez-Sanchez M, Velasco-Ramos E, Ruiz-Munoz M et al. Cross-cultural adaptation and validation of the Spanish version of the American Academy of Orthopaedic Surgeons-Foot and Ankle Module (AAOS-FAMsp). *J Orthop Surg Res* 2016;11(1):74.
 22. Kim JB, Kim JK, Seo SG et al. Validity, reliability, and responsiveness of the Korean version of American Academy of Orthopedic Surgeons Foot and Ankle questionnaire. *J Foot Ankle Surg* 2015;54(1):46-50.
 23. Kitaoka HB, Alexander IJ, Adelaar RS et al. Clinical rating systems for the ankle-hindfoot, midfoot, hallux, and lesser toes. *Foot Ankle Int* 1994;15(7):349-53.
 24. Coster MC, Rosengren BE, Bremander A et al. Comparison of the Self-reported Foot and Ankle Score (SEFAS) and the American Orthopedic Foot and Ankle Society Score (AOFAS). *Foot Ankle Int* 2014;35(10):1031-6.
 25. Baumhauer JF, Nawoczenski DA, DiGiovanni BF et al. Reliability and validity of the American Orthopaedic Foot and Ankle Society Clinical Rating Scale: a pilot study for the hallux and lesser toes. *Foot Ankle Int* 2006;27(12):1014-9.
 26. SooHoo NF, Shuler M, Fleming LL. Evaluation of the validity of the AOFAS Clinical Rating Systems by correlation to the SF-36. *Foot Ankle Int* 2003;24(1):50-5.
 27. Dawson J, Coffey J, Doll H et al. A patient-based questionnaire to assess outcomes of foot surgery: validation in the context of surgery for hallux valgus. *Qual Life Res* 2006;15(7):1211-22.
 28. Malviya A, Makwana N, Laing P. Correlation of the AOFAS scores with a generic health QUALY score in foot and ankle surgery. *Foot Ankle Int* 2007;28(4):494-8.
 29. Dawson J, Doll H, Coffey J et al. Responsiveness and minimally important change for the Manchester-Oxford Foot Questionnaire (MOXFQ) compared with AOFAS and SF-36 assessments following surgery for hallux valgus. *Osteoarth Cart* 2007;15(8):918-31.
 30. Conceicao CS, Neto MG, Neto AC et al. Analysis of the psychometric properties of the American Orthopaedic Foot and Ankle Society Score (AOFAS) in rheumatoid arthritis patients: application of the Rasch model. *Rev Bras Reumatol Engl Ed* 2016;56(1):8-13.
 31. Rodrigues R, Masiero D, Mizusaki J et al. Translation, cultural adaptation, and validity of the American Orthopaedic Foot and Ankle Society (AOFAS) Ankle-Hindfoot Scale. *Acta Ortop Bras* 2008;16(2):107-11.
 32. Kostuj T, Krummenauer F, Schaper K et al. Analysis of agreement between the German translation of the American Foot and Ankle Society's Ankle and Hindfoot Scale (AOFAS-AHS) and the Foot Function Index in its validated German translation by Naal et al. (FFI-D). *Arch Orthop Trauma Surg* 2014;134(9):1205-10.
 33. Leigheb M, Janicka P, Andorno S et al. Italian translation, cultural adaptation and validation of the 'American Orthopaedic Foot and Ankle Society's (AOFAS) Ankle-Hindfoot Scale'. *Acta Biomed* 2016;87(1):38-45.
 34. Roos EM, Brandsson S, Karlsson J. Validation of the Foot and Ankle Outcome Score for ankle ligament reconstruction. *Foot Ankle Int* 2001;22(10):788-94.
 35. Sierevelt IN, Beimers L, van Bergen CJ et al. Validation of the Dutch language version of the Foot and Ankle Outcome Score. *Knee Surg Sports Traumatol Arthrosc* 2015;23(8):2413-9.

36. Sierevelt IN, van Eekeren IC, Haverkamp D et al. Evaluation of the Dutch version of the Foot and Ankle Outcome Score (FAOS): responsiveness and minimally important change. *Knee Surg Sports Traumatol Arthrosc* 2016;24(4):1339-47.
37. van den Akker-Scheek I, Seldentuis A, Reininga IH et al. Reliability and validity of the Dutch version of the Foot and Ankle Outcome Score (FAOS). *BMC Musculoskelet Disord* 2013;14:183.
38. van Bergen CJ, Sierevelt IN, Hoogervorst P et al. Translation and validation of the German version of the Foot and Ankle Outcome Score. *Arch Orthop Trauma Surg* 2014;134(7):897-901.
39. Lee KM, Chung CY, Kwon SS et al. Transcultural adaptation and testing psychometric properties of the Korean version of the Foot and Ankle Outcome Score (FAOS). *Clin Rheumatol* 2013;32(10):1443-50.
40. Negahban H, Mazaheri M, Salavati M et al. Reliability and validity of the Foot and Ankle Outcome Score: a validation study from Iran. *Clin Rheumatol* 2010;29(5):479-86.
41. Karatepe AG, Gunaydin R, Kaya T et al. Validation of the Turkish version of the Foot and Ankle Outcome Score. *Rheumatol Int* 2009;30(2):169-73.
42. Chen L, Lyman S, Do H et al. Validation of Foot and Ankle Outcome Score for hallux valgus. *Foot Ankle Int* 2012;33(12):1145-55.
43. Hogan MV, Mani SB, Chan JY et al. Validation of the Foot and Ankle Outcome Score for hallux rigidus. *HSS J* 2016;12(1):44-50.
44. Mani SB, Brown HC, Nair P et al. Validation of the Foot and Ankle Outcome Score in adult-acquired flatfoot deformity. *Foot Ankle Int* 2013;34(8):1140-6.
45. Mani SB, Do H, Vulcano E et al. Evaluation of the Foot and Ankle Outcome Score in patients with osteoarthritis of the ankle. *Bone Joint J* 2015;97-b(5):662-7.
46. Binkley JM, Stratford PW, Lott SA et al. The Lower Extremity Functional Scale (LEFS): scale development, measurement properties, and clinical application. North American Orthopaedic Rehabilitation Research Network. *Phys Ther* 1999;79(4):371-83.
47. Alcock G, Stratford P. Validation of the Lower Extremity Functional Scale on athletic subjects with ankle sprains. *Physiother Can* 2002;54:233-40.
48. Lin CW, Moseley AM, Refshauge KM et al. The Lower Extremity Functional Scale has good clinimetric properties in people with ankle fracture. *Phys Ther* 2009;89(6):580-8.
49. Gabel CP, Melloh M, Burkett B et al. Lower Limb Functional Index: development and clinimetric properties. *Phys Ther* 2012;92(1):98-110.
50. Pan SL, Liang HW, Hou WH et al. Responsiveness of SF-36 and Lower Extremity Functional Scale for assessing outcomes in traumatic injuries of lower extremities. *Injury* 2014;45(11):1759-63.
51. Alnahdi AH, Alrashid GI, Alkhaldi HA et al. Cross-cultural adaptation, validity and reliability of the Arabic version of the Lower Extremity Functional Scale. *Disabil Rehabil* 2016;38(9):897-904.
52. Pereira LM, Dias JM, Mazuquin BF et al. Translation, cross-cultural adaptation and analysis of the psychometric properties of the Lower Extremity Functional Scale (LEFS): LEFS-BRAZIL. *Braz J Phys Ther* 2013;17(3):272-80.
53. Hoogboom TJ, de Bie RA, den Broeder AA et al. The Dutch Lower Extremity Functional Scale was highly reliable, valid and responsive in individuals with hip/knee osteoarthritis: a validation study. *BMC Musculoskelet Disord* 2012;13:117.
54. Repo JP, Tukiainen EJ, Roine RP et al. Reliability and validity of the Finnish version of the Lower Extremity Functional Scale (LEFS). *Disabil Rehabil* 2016 Jul 6:1-7. [Epub ahead of print]
55. René F, Casimiro L, Tremblay M et al. Fiabilité test retest et validité de construit de la version française de L'Échelle fonctionnelle des membres inférieurs (ÉFMI), partie II. *Physiother Can* 2011;63(2):249-55.
56. Cacchio A, De Blasis E, Necozone S et al. The Italian version of the Lower Extremity Functional Scale was reliable, valid, and responsive. *J Clin Epidemiol* 2010;63(5):550-7.
57. Negahban H, Hessam M, Tabatabaei S et al. Reliability and validity of the Persian Lower Extremity Functional Scale (LEFS) in a heterogeneous sample of outpatients with lower limb musculoskeletal disorders. *Disabil Rehabil* 2014;36(1):10-5.
58. Cruz-Diaz D, Lomas-Vega R, Osuna-Perez MC et al. The Spanish Lower Extremity Functional Scale: a reliable, valid and responsive questionnaire to assess musculoskeletal disorders in the lower extremity. *Disabil Rehabil* 2014;36(23):2005-11.
59. Hou WH, Yeh TS, Liang HW. Reliability and validity of the Taiwan Chinese version of the Lower Extremity Functional Scale. *J Formos Med Assoc* 2014;113(5):313-20.
60. Citaker S, Kafa N, Hazar Kanik Z et al. Translation, cross-cultural adaptation and validation of the Turkish version of the Lower Extremity Functional Scale on patients with knee injuries. *Arch Orthop Trauma Surg* 2016;136(3):389-95.
61. Cuesta-Vargas AI, Gabel CP, Bennett P. Cross-cultural adaptation and validation of a Spanish version of the Lower Limb Functional Index. *Health Qual Life Outcomes* 2014;12:75.
62. Duruturk N, Tonga E, Gabel CP et al. Cross-cultural adaptation, reliability and validity of the Turkish version of the Lower Limb Functional Index. *Disabil Rehabil* 2015 Jul 26:1-6. [Epub ahead of print]
63. McNair PJ, Prapavessis H, Collier J et al. The lower-limb tasks questionnaire: an assessment of validity, reliability, responsiveness, and minimal important differences. *Arch Phys Med Rehabil* 2007;88(8):993-1001.
64. Chhabra A, Soldatos T, Chalian M, 3-tesla magnetic resonance imaging evaluation of posterior tibial tendon dysfunction with relevance to clinical staging. *J Foot Ankle Surg* 2011;50:320-328.
65. Simpson MR, Howard TM. Tendinopathies of the foot and ankle. *Am Fam Physician* 2009;80(10):1107-14.
66. Rodriguez D, Devos Bevernage B, Maldague P et al. Tarsal tunnel syndrome and flexor hallucis longus tendon hypertrophy. *Orthop Traumatol Surg Res* 2010;96(7):829-31.
67. Patel AT, Gaines K, Malamut R et al. Usefulness of electrodiagnostic techniques in the evaluation of suspected

- tarsal tunnel syndrome: an evidence-based review. *Muscle Nerve* 2005;32(2):236-40.
68. Oh SJ, Meyer RD. Entrapment neuropathies of the tibial (posterior tibial) nerve. *Neurol Clin* 1999;17(3):593-615, vii.
 69. Skalley TC, Schon LC, Hinton RY et al. Clinical results following revision tibial nerve release. *Foot Ankle Int* 1994;15(7):360-7.
 70. Pfeiffer WH, Cracchiolo A, 3rd. Clinical results after tarsal tunnel decompression. *J Bone Joint Surg Am* 1994;76(8):1222-30.
 71. Lo LD, Schweitzer ME, Fan JK et al. MR imaging findings of entrapment of the flexor hallucis longus tendon. *AJR Am J Roentgenol* 2001;176(5):1145-8.
 72. Khan K, Brown J, Way S et al. Overuse injuries in classical ballet. *Sports Med* 1995;19(5):341-57.
 73. Brukner P, Bennell K, Matheson G. *Stress Fractures*. Melbourne: Blackwells Scientific Asia, 1999.
 74. Jowett AJ, Birks CL, Blackney MC. Medial malleolar stress fracture secondary to chronic ankle impingement. *Foot Ankle Int* 2008;29(7):716-21.
 75. Irion V, Miller TL, Kaeding CC. The treatment and outcomes of medial malleolar stress fractures: a systematic review of the literature. *Sports Health* 2014;6(6):527-30.
 76. Counsel PD, Davenport M, Brown A et al. Ultrasound-guided radiofrequency denervation of the medial calcaneal nerve. *Clin J Sport Med* 2016;26(6):465-470.
 77. Clarke HD, Kitaoka HB, Ehman RL. Peroneal tendon injuries. *Foot Ankle Int* 1998;19(5):280-8.
 78. Tjin ATER, Schweitzer ME, Karasick D. MR imaging of peroneal tendon disorders. *AJR Am J Roentgenol* 1997;168(1):135-40.
 79. Maffulli N, Ferran NA, Oliva F et al. Recurrent subluxation of the peroneal tendons. *Am J Sports Med* 2006;34(6):986-92.
 80. Hight RM. Diagnosis of anterolateral ankle impingement: comparison between magnetic resonance imaging and clinical examination. *Am J Sports Med* 1998;26(1):152-3.
 81. Liu SH, Nuccion SL, Finerman G. Diagnosis of anterolateral ankle impingement. Comparison between magnetic resonance imaging and clinical examination. *Am J Sports Med* 1997;25(3):389-93.
 82. Bradshaw C, Khan K, Brukner P. Stress fracture of the body of the talus in athletes demonstrated with computer tomography. *Clin J Sport Med* 1996;6(1):48-51.
 83. Rossi F, Dragoni S. Talar body fatigue stress fractures: three cases observed in elite female gymnasts. *Skeletal Radiol* 2005;34(7):389-94.
 84. Kim YS, Lee HM, Kim JP et al. Fatigue stress fracture of the talar body: an uncommon cause of ankle pain. *J Foot Ankle Surg* 2016;55(5):1113-6.
 85. van den Bekerom MP, Raven EE. The distal fascicle of the anterior inferior tibiofibular ligament as a cause of tibiotalar impingement syndrome: a current concepts review. *Knee Surg Sports Traumatol Arthrosc* 2007;15(4):465-71.
 86. Talusan PG, Toy J, Perez JL et al. Anterior ankle impingement: diagnosis and treatment. *J Am Acad Orthop Surg* 2014;22(5):333-9.
 87. Parma A, Buda R, Vannini F et al. Arthroscopic treatment of ankle anterior bony impingement: the long-term clinical outcome. *Foot Ankle Int* 2014;35(2):148-55.
 88. Walsh SJ, Twaddle BC, Rosenfeldt MP et al. Arthroscopic treatment of anterior ankle impingement: a prospective study of 46 patients with 5-year follow-up. *Am J Sports Med* 2014;42(11):2722-6.
 89. Zwiers R, Wiegelerinck JI, Murawski CD et al. Arthroscopic treatment for anterior ankle impingement: a systematic review of the current literature. *Arthroscopy* 2015;31(8):1585-96.

CHAPTER 43

1. McPoil TG, Martin RRL, Cornwall MW et al. Heel pain-plantar fasciitis: clinical practice guidelines linked to the international classification of functioning, disability, and health from the orthopaedic section of the American Physical Therapy Association. *J Orth Phys Ther* 2008;38(4):a1-18.
2. De Garceau D, Dean D, Requejo SM et al. The association between diagnosis of plantar fasciitis and windlass test results. *Foot Ankle Int* 2003;24(3):251-5.
3. Budiman-Mak E, Conrad KJ, Roach KE. The Foot Function Index: a measure of foot pain and disability. *J Clin Epidemiol* 1991;44(6):561-70.
4. SooHoo NF, Samimi DB, Vyas RM et al. Evaluation of the validity of the Foot Function Index in measuring outcomes in patients with foot and ankle disorders. *Foot Ankle Int* 2006;27(1):38-42.
5. Madeley NJ, Wing KJ, Topliss C et al. Responsiveness and validity of the SF-36, Ankle Osteoarthritis Scale, AOFAS Ankle Hindfoot Score, and Foot Function Index in end stage ankle arthritis. *Foot Ankle Int* 2012;33(1):57-63.
6. Budiman-Mak E, Conrad K, Stuck R et al. Theoretical model and Rasch analysis to develop a revised Foot Function Index. *Foot Ankle Int* 2006;27(7):519-27.
7. Jorgensen JE, Andreassen J, Rathleff MS. Translation and validation of the Danish Foot Function Index (FFI-DK). *Scand J Med Sci Sports* 2015;25(4):e408-13.
8. Pourtier-Piotte C, Pereira B, Soubrier M et al. French validation of the Foot Function Index (FFI). *Ann Phys Rehabil Med* 2015;58(5):276-82.
9. Naal FD, Impellizzeri FM, Huber M et al. Cross-cultural adaptation and validation of the Foot Function Index for use in German-speaking patients with foot complaints. *Foot Ankle Int* 2008;29(12):1222-8.
10. Martinelli N, Scotto GM, Sartorelli E et al. Reliability, validity and responsiveness of the Italian version of the Foot Function Index in patients with foot and ankle diseases. *Qual Life Res* 2014;23(1):277-84.
11. Vetrano M, Vulpiani MC, Erroi D et al. Cross-cultural adaptation and reliability of the Italian version of the Foot Function Index (FFI-I) for patients with plantar fasciitis. *J Sports Med Phys Fitness* 2014;54(5):636-43.

12. Huh JW, Eun IS, Ko YC et al. Reliability and validity of the Korean Version of the Foot Function Index. *J Foot Ankle Surg* 2016;55(4):759-61.
13. Paez-Moguer J, Budiman-Mak E, Cuesta-Vargas AI. Cross-cultural adaptation and validation of the Foot Function Index to Spanish. *Foot Ankle Surg* 2014;20(1):34-9.
14. Wu SH, Liang HW, Hou WH. Reliability and validity of the Taiwan Chinese version of the Foot Function Index. *J Formos Med Assoc* 2008;107(2):111-8.
15. Rowan K. The development and validation of a multi-dimensional measure of chronic foot pain: the Rowan Foot Pain Assessment Questionnaire (ROFPAQ). *Foot Ankle Int* 2001;22(10):795-809.
16. Landorf KB, Radford JA, Hudson S. Minimal important difference (MID) of two commonly used outcome measures for foot problems. *J Foot Ankle Res* 2010;3:7.
17. Bennett PJ, Patterson C, Wearing S et al. Development and validation of a questionnaire designed to measure foot-health status. *J Am Podiatr Med Assoc* 1998;88(9):419-28.
18. Ferreira AF, Laurindo IM, Rodrigues PT et al. Brazilian version of the Foot Health Status Questionnaire (FHSQ-BR): cross-cultural adaptation and evaluation of measurement properties. *Clinics (Sao Paulo)* 2008;63(5):595-600.
19. Cuesta-Vargas A, Bennett P, Jimenez-Cebrian AM et al. The psychometric properties of the Spanish version of the Foot Health Status Questionnaire. *Qual Life Res* 2013;22(7):1739-43.
20. Macran S, Kind P, Collingwood J et al. Evaluating podiatry services: testing a treatment specific measure of health status. *Qual Life Res* 2003;12(2):177-88.
21. Lemont H, Ammirati KM, Usen N. Plantar fasciitis: a degenerative process (fasciosis) without inflammation. *J Am Podiatr Med Assoc* 2003;93(3):234-7.
22. Snider MP, Clancy WG, McBeath AA. Plantar fascia release for chronic plantar fasciitis in runners. *Am J Sports Med* 1983;11(4):215-9.
23. Tountas AA, Fornasier VL. Operative treatment of subcalcaneal pain. *Clin Orthop Relat Res* 1996;332:170-8.
24. Dunn JE, Link CL, Felson DT et al. Prevalence of foot and ankle conditions in a multiethnic community sample of older adults. *Am J Epidemiol* 2004;159(5):491-8.
25. Hill CL, Gill TK, Menz HB et al. Prevalence and correlates of foot pain in a population-based study: the North West Adelaide health study. *J Foot Ankle Res* 2008;1(1):2.
26. Di Caprio F, Buda R, Mosca M et al. Foot and lower limb diseases in runners: assessment of risk factors. *J Sports Sci Med* 2010;9(4):587-96.
27. van Leeuwen KD, Rogers J, Winzenberg T et al. Higher body mass index is associated with plantar fasciopathy/'plantar fasciitis': systematic review and meta-analysis of various clinical and imaging risk factors. *Br J Sports Med* 2016;50(16):972-81.
28. Tenforde AS, Sayres LC, McCurdy ML et al. Overuse injuries in high school runners: lifetime prevalence and prevention strategies. *PM&R* 2011;3(2):125-31; quiz 131.
29. Bolivar YA, Munuera PV, Padillo JP. Relationship between tightness of the posterior muscles of the lower limb and plantar fasciitis. *Foot Ankle Int* 2013;34(1):42-8.
30. McMillan AM, Landorf KB, Barrett JT et al. Diagnostic imaging for chronic plantar heel pain: a systematic review and meta-analysis. *J Foot Ankle Res* 2009;2:32.
31. Smith S, Tinley P, Gilheany M et al. The inferior calcaneal spur-anatomical and histological considerations. *Foot* 2007;17:25-31.
32. Chimutengwende-Gordon M, O'Donnell P, Singh D. Magnetic resonance imaging in plantar heel pain. *Foot Ankle Int* 2010;31(10):865-70.
33. Guermazi A, Roemer FW, Crema M. *Imaging in Sports-specific Musculoskeletal Injuries*. Switzerland: Springer International Publishing Switzerland, 2016.
34. Thomas JL, Christensen JC, Kravitz SR et al. The diagnosis and treatment of heel pain: a clinical practice guideline-revision 2010. *J Foot Ankle Surg* 2010;49(3 Suppl):S1-19.
35. Landorf KB. Plantar heel pain and plantar fasciitis. *BMJ Clin Evid* 2015 Nov 25; pii: 1111.
36. Martin RL, Irrgang JJ, Cionti SF. Outcome study of subjects with insertional plantar fasciitis. *Foot Ankle Int* 1998;19:803-11.
37. Landorf KB, Radford JA, Keenan A-M et al. Effectiveness of low-Dye taping for the short-term management of plantar fasciitis. *J Am Pod Med Assoc* 2005;95(6):525-30.
38. Radford JA, Landorf KB, Buchbinder R et al. Effectiveness of low-Dye taping for the short-term treatment of plantar heel pain: a randomized trial. *BMC Musculoskelet Disord* 2006;7:64-70.
39. Pfeffer G, Bacchetti P, Deland J et al. Comparison of custom and prefabricated orthoses in the initial treatment of proximal plantar fasciitis. *Foot Ankle Int* 1999;20(4):214-21.
40. Gross MT, Byers JM, Krafft JL et al. The impact of custom semirigid foot orthotics on pain and disability for individuals with plantar fasciitis. *J Orthop Sports Phys Ther* 2002;32:149-57.
41. Yamashita MH. Evaluation and selection of shoe wear and orthoses for the runner. *Phys Med Rehabil Clin N Am* 2005;16(3):801-29.
42. DiGiovanni BF, Nawoczenski DA, Lintal ME et al. Tissue-specific plantar fascia-stretching exercise enhances outcomes in patients with chronic heel pain. A prospective randomized study. *J Bone Joint Surg Br* 2003;85-A(7):1270-7.
43. Porter D, Barrill E, Oneacre K et al. The effects of duration and frequency of Achilles tendon stretching on dorsiflexion and outcome in painful heel syndrome: a randomized, blinded, control study. *Foot Ankle Int* 2002;23(7):619-24.
44. Celik D, Kus G, Sirma SO. Joint mobilization and stretching exercise vs steroid injection in the treatment of plantar fasciitis: a randomized controlled study. *Foot Ankle Int* 2016;37(2):150-6.

45. Ajimsha MS, Binsu D, Chithra S. Effectiveness of myofascial release in the management of plantar heel pain: a randomized controlled trial. *Foot (Edinb)* 2014;24(2):66-71.
46. Donley BG, Moore T, Sfera J et al. The efficacy of oral nonsteroidal anti-inflammatory medication (NSAID) in the treatment of plantar fasciitis: a randomized, prospective, placebo-controlled study. *Foot Ankle Int* 2007;28(1):20-3.
47. Cotchett MP, Munteanu SE, Landorf KB. Effectiveness of trigger point dry needling for plantar heel pain: a randomized controlled trial. *Phys Ther* 2014;94(8):1083-94.
48. Tatli YZ, Kapasi S. The real risks of steroid injection for plantar fasciitis, with a review of conservative therapies. *Curr Rev Musculoskelet Med* 2009;2:3-9.
49. Karls SL, Snyder KR, Neibert PJ. Effectiveness of corticosteroid injections in the treatment of plantar fasciitis. *J Sport Rehabil* 2016;25(2):202-7.
50. Li Z, Xia C, Yu A et al. Ultrasound-versus palpation-guided injection of corticosteroid for plantar fasciitis: a meta-analysis. *PLoS One* 2014;9(3):e92671.
51. Sellman JR. Plantar fascia rupture associated with corticosteroid injection. *Foot and Ankle International* 1994;15(7):376-81.
52. Acevedo JL, Beskin JL. Complications of plantar fascia rupture associated with corticosteroid injection. *Foot Ankle Int* 1998;19:91-7.
53. Snow DM, Reading J, Dalal R. Lateral plantar nerve injury following steroid injection for plantar fasciitis. *Br J Sports Med* 2005;39(12):e41; discussion e41.
54. Rathleff MS, Molgaard CM, Fredberg U et al. High-load strength training improves outcome in patients with plantar fasciitis: a randomized controlled trial with 12-month follow-up. *Scand J Med Sci Sports* 2015;25(3):e292-300.
55. Batt ME, Tanji JL, Skattum N. Plantar fasciitis: a prospective randomized clinical trial of the tension night splint. *Clin J Sport Med* 1996;6:158-62.
56. Powell M, Post WR, Keener J et al. Effective treatment of chronic plantar fasciitis with dorsiflexion night splints: a crossover prospective randomised outcome study. *Foot Ankle Int* 1998;19(1):10-8.
57. Probe RA, Baca M, Adams R et al. Night splint treatment for plantar fasciitis. A prospective randomized study. *Clin Orthop Relat Res* 1999;368:190-5.
58. Speed C. A systematic review of shockwave therapies in soft tissue conditions: focusing on the evidence. *Br J Sports Med* 2014;48(21):1538-42.
59. Sammarco GJ, Helfrey RB. Surgical treatment of recalcitrant plantar fasciitis. *Foot Ankle Int* 1996;17:520-6.
60. Jerosch J, Schunck J, Liebsch D et al. Indication, surgical technique and results of endoscopic fascial release in plantar fasciitis (EFRPF). *Knee Surg Sports Traumatol Arthrosc* 2004;12:471-7.
61. O'Malley MJ, Page A, Cook R. Endoscopic plantar fasciotomy for chronic heel pain. *Foot Ankle Int* 2000;21:505-10.
62. Spears IR, Miller-Young JE, Sharma J et al. The potential influence of the heel counter on internal stress during static standing: a combined finite element and positional MRI investigation. *J Biomech* 2007;40(12):2774-80.
63. Sormaa MJ, Niva MH, Kiuru MJ et al. Stress injuries of the calcaneus detected with magnetic resonance imaging in military recruits. *J Bone Joint Surg Am* 2006;88(10):2237-42.
64. Baxter D, Pfeffer G. Treatment of chronic heel pain by surgical release of the first branch of the lateral plantar nerve. *Clin Orthop Relat Res* 1992;279:229-36.
65. Chundru U, Liebeskind A, Seidelmann F et al. Plantar fasciitis and calcaneal spur formation are associated with abductor digiti minimi atrophy on MRI of the foot. *Skeletal Radiol* 2008;37(6):505-10.
66. Meadows JR, Finnoff JT. Lower extremity nerve entrapments in athletes. *Curr Sports Med Rep* 2014;13(5):299-306.
67. Lareau CR, Sawyer GA, Wang JH et al. Plantar and medial heel pain: diagnosis and management. *J Am Acad Orthop Surg* 2014;22(6):372-80.
68. Hoffman D, Bianchi S. Sonographic evaluation of plantar hindfoot and midfoot pain. *J Ultrasound Med* 2013;32(7):1271-84.
69. Presley JC, Maida E, Pawlina W et al. Sonographic visualization of the first branch of the lateral plantar nerve (Baxter nerve): technique and validation using perineural injections in a cadaveric model. *J Ultrasound Med* 2013;32(9):1643-52.
70. Khan KM, Brukner PD, Kearney C et al. Tarsal navicular stress fracture in athletes. *Sports Med* 1994;17:65-76.
71. Karas MA, Hoy DJ. Compensatory midfoot dorsiflexion in the individual with heelcord tightness: implications for orthotic device designs. *J Prosthet Orthot* 2002;14:82-93.
72. Khan KM, Fuller PJ, Brukner PD et al. Outcome of conservative and surgical management of navicular stress fracture in athletes. Eighty-six cases proven with computerized tomography. *Am J Sports Med* 1992;20(6):657-66.
73. Kiss ZA, Khan KM, Fuller PJ. Stress fractures of the tarsal navicular bone: CT findings in 55 cases. *AJR Am J Roentgenol* 1993;160:111-5.
74. Torg JS, Moyer J, Gaughan JP et al. Management of tarsal navicular stress fractures. *Am J Sports Med* 2010;38(5):1048-53.
75. Mayer SW, Joyner PW, Almekinders LC et al. Stress fractures of the foot and ankle in athletes. *Sports Health* 2014;6(6):481-91.
76. Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016;50(5):273-80.
77. Mallee WH, Weel H, van Dijk CN et al. Surgical versus conservative treatment for high-risk stress fractures of the lower leg (anterior tibial cortex, navicular and fifth metatarsal base): a systematic review. *Br J Sports Med* 2015;49(6):370-6.
78. Lattermann C, Goldstein J, Wukich D et al. Practical management of Lisfranc injuries in athletes. *Clin J Sport Med* 2007;17(4):311-5.
79. Rhim B, Hunt JC. Lisfranc injury and Jones fracture in sports. *Clin Podiatr Med Surg* 2011;28(1):69-86.

80. Crim J. MR imaging evaluation of subtle Lisfranc injuries: the midfoot sprain. *Magn Reson Imaging Clin N Am* 2008;16(1):19-27, v.
81. Libby B, Ersoy H, Pomeranz SJ. Imaging of the Lisfranc injury. *J Surg Orthop Adv* 2015;24(1):79-82.
82. Woodward S, Jacobson JA, Femino JE et al. Sonographic evaluation of Lisfranc ligament injuries. *J Ultrasound Med* 2009;28(3):351-7.
83. Nunley JA, Vertullo CJ. Classification, investigation, and management of midfoot sprains. Lisfranc injuries in the athlete. *Am J Sports Med* 2002;30(6):871-8.
84. Welck MJ, Zinchenko R, Rudge B. Lisfranc injuries. *Injury* 2015;46(4):536-41.
85. Marx RC, Mizel MS. What's new in foot and ankle surgery. *J Bone Joint Surg Am* 2012;94(10):952-7.
86. Marshall P, Hamilton WG. Cuboid subluxation in ballet dancers. *Am J Sports Med* 1992;20(2):169-75.
87. Patterson SM. Cuboid syndrome: a review of the literature. *J Sports Sci Med* 2006;5:597-606.
88. Hermel MB, Gershon-Cohen J. The nutcracker fracture of the cuboid by indirect violence. *Radiology* 1953;60:850-4.
89. Jennings J, Davies GJ. Treatment of cuboid syndrome secondary to lateral ankle sprains: a case series. *J Orthop Sports Phys Ther* 2005;35:409-15.
90. Joo SY, Jeong C. Stress fracture of the tarsal cuboid bone in early childhood. *Eur J Orthop Surg Traumatol* 2015;25:595-9.
91. Yu SM, Dardani M, Yu JS. MRI of isolated cuboid stress fractures in adults. *AJR Am J Roentgenol* 2013;201(6):1325-30.
92. Khan KM, Brukner PD, Bradshaw C. Stress fracture of the medial cuneiform bone in a runner. *Clin J Sport Med* 1993;3(4):262-4.
93. Bui-Mansfield LT, Thomas WR. Magnetic resonance imaging of stress injury of the cuneiform bones in patients with plantar fasciitis. *J Comput Assist Tomogr* 2009;33(4):593-6.
94. Creighton R, Sonogar A, Gordon G. Stress fracture of the tarsal middle cuneiform bone. A case report. *J Am Pod Med Assoc* 1990;80:489-95.
95. Solomon LB, Ruhli FJ, Taylor J et al. A dissection and computer tomograph study of tarsal coalitions in 100 cadaver feet. *J Orthop Res* 2003;21(2):352-8.
96. Thomason K, Stephens MM. Tarsal coalition. *Curr Orth Prac* 2012;23:229-34.
97. Moraleda L, Gantsoudas G, Mubarak S. C sign: talocalcaneal coalition or flatfoot deformity? *J Pediatr Orth* 2014;34:814-9.
98. Sharp RJ, Wade CM, Hennessy MS et al. The role of MRI and ultrasound imaging in Morton's neuroma and the effect of size of lesion on symptoms. *J Bone Joint Surg Br* 2003;85(7):999-1005.
99. Brukner PD, Bennell KL, Matheson GO. *Stress Fractures*. Melbourne: Blackwells Scientific Asia, 1999.
100. Bennell K, Matheson G, Meeuwisse W et al. Risk factors for stress fractures. *Sports Med* 1999;28(2):91-122.
101. Nagel A, Fernholz F, Kibele C et al. Long-distance running increases plantar pressures beneath the metatarsal heads: a barefoot walking investigation of 200 marathon runners. *Gait Posture* 2008;27(1):152-5.
102. Weist R, Eils E, Rosenbaum D. The influence of muscle fatigue on electromyogram and plantar pressure patterns as an explanation for the incidence of metatarsal stress fractures. *Am J Sports Med* 2004;32(8):1893-8.
103. Snyder R, DeAngelis J, Koester M et al. Does shoe insole modification prevent stress fractures? A systematic review. *HSS Journal* 2009;5(2):92-8.
104. Harrington T, Crichton KJ, Anderson IF. Overuse ballet injury of the base of the second metatarsal. *Am J Sports Med* 1993;21:591-8.
105. Micheli LJ, Sohn RS, Solomon R. Stress fractures of the second metatarsal involving Lisfranc's joint in ballet dancers: a new overuse of the foot. *J Bone Joint Surg Am* 1985;67A:1372-5.
106. O'Malley MJ, Hamilton WG, Munyak J et al. Stress fractures at the base of the second metatarsal in ballet dancers. *Foot Ankle Int* 1996;17(2):89-94.
107. Chuckpaiwong B, Cook C, Pietrobon R et al. Second metatarsal stress fracture in sport: comparative risk factors between proximal and non-proximal locations. *Br J Sports Med* 2007;41:510-4.
108. Jones R. Fracture of the base of the fifth metatarsal bone by indirect violence. *Ann Surg* 1902;35:697-700.
109. Lawrence SJ, Botte MJ. Jones' fractures and related fractures of the proximal fifth metatarsal. *Foot Ankle* 1993;14(6):358-65.
110. Zwitser EW, Breederveld RS. Fractures of the fifth metatarsal; diagnosis and treatment. *Injury* 2010;41(6):555-62.
111. Ekstrand J, van Dijk CN. Fifth metatarsal fractures among male professional footballers: a potential career-ending disease. *Br J Sports Med* 2013;47(12):754-8.
112. Torg JS, Balduini FC, Zelko RR et al. Fractures of the base of the fifth metatarsal distal to the tuberosity. Classification and guidelines for non-surgical and surgical management. *J Bone Joint Surg Am* 1984;66(2):209-14.
113. Leumann A, Pagenstert G, Fuhr P et al. Intramedullary screw fixation in proximal fifth-metatarsal fractures in sports: clinical and biomechanical analysis. *Arch Orthop Trauma Surg* 2008;128:1425-30.
114. Mologne TS, Lundeen JM, Clapper MF et al. Early screw fixation versus casting in the treatment of acute Jones fractures. *Am J Sports Med* 2005;33(7):970-5.
115. Den Hartog BD. Fracture of the proximal fifth metatarsal. *J Am Acad Orthop Surg* 2009;17(7):458-64.
116. Porter D, Duncan M, Meyer S. Fifth metatarsal Jones fracture fixation with a 4.5-mm cannulated stainless steel screw in the competitive and recreational athlete. *Am J Sports Med* 2005;33:726-33.
117. Hunt KJ, Anderson RB. Treatment of Jones fracture nonunions and refractures in the elite athlete: outcomes of intramedullary screw fixation with bone grafting. *Am J Sports Med* 2011;39(9):1948-54.
118. Wright R, Fischer D, Shively R et al. Refracture of proximal fifth metatarsal (Jones) fracture after intramedullary screw fixation in athletes. *Am J Sports Med* 2000;28(5):732-6.

119. Larson CM, Almekinders LC, Taft TN et al. Intramedullary screw fixation of Jones fractures—analysis of failure. *Am J Sports Med* 2002;30(1):55–60.
120. Bigsby E, Halliday R, Middleton RG et al. Functional outcome of fifth metatarsal fractures. *Injury* 2014;45(12):2009–12.
121. Perez HR, Roberts J. Flexor tendon sheath as a source of pain in lesser metatarsal overload. *J Am Podiatr Med Assoc* 2009;99(2):129–34.
122. Panchbhavi VK, Trevino S. Clinical tip: a new clinical sign associated with metatarsophalangeal joint synovitis of the lesser toes. *Foot Ankle Int* 2007;28(5):640–1.
123. McCormick JJ, Anderson RB. Rehabilitation following turf toe injury and plantar plate repair. *Clin Sports Med* 2010;29:313–23.
124. Anderson RB. Turf toe injuries of the hallux metatarsophalangeal joint. *Tech Foot Ankle Surg* 2002;1(2):102–11.
125. Grady JF, Axe TM, Zager EJ et al. A retrospective analysis of 772 patients with hallux limitus. *J Am Podiatr Med Assoc* 2002;92(2):102–8.
126. Kalish SR, Willis FB. Hallux limitus and dynamic splinting: a retrospective series. *FAOJ* 2009;2(4):1.
127. Nix S, Smith M, Vicenzino B. Prevalence of hallux valgus in the general population: a systematic review and meta-analysis. *J Foot Ankle Res* 2010;3:21.
128. Shine J, Weil L, Weil LS et al. Scarf osteotomy for the correction of adolescent hallux valgus. *Foot Ankle Spec* 2010;3:10–4.
129. Munuera PV, Polo J, Rebollo J. Length of the first metatarsal and hallux in hallux valgus in the initial stage. *Internat Orth (SICOT)* 2008;32:489–95.
130. Maffulli N, Longo UG, Marinozzi A et al. Hallux valgus: effectiveness and safety of minimally invasive surgery. A systematic review. *Brit Med Bull* 2011;97:149–67.
131. Schuh R, Hofstaetter SG, Adams SB et al. Rehabilitation after hallux valgus surgery: importance of physical therapy to restore weight bearing of the first ray during the stance phase. *Phys Ther* 2009;89(9):934–45.
132. Dedmond BT, Cory JW, McBryde A, Jr. The hallucal sesamoid complex. *J Am Acad Orthop Surg* 2006;14(13):745–53.
133. Bichara DA, Henn RF, 3rd, Theodore GH. Sesamoidectomy for hallux sesamoid fractures. *Foot Ankle Int* 2012;33(9):704–6.
134. Yu GV, Judge MS, Hudson JR et al. Predislocation syndrome. Progressive subluxation/dislocation of the lesser metatarsophalangeal joint. *J Am Podiatr Med Assoc* 2002;92(4):182–99.
135. Kier R, Abrahamian H, Caminear D et al. MR arthrography of the second and third metatarsophalangeal joints for the detection of tears of the plantar plate and joint capsule. *AJR Am J Roentgenol* 2010;194(4):1079–81.
136. Owens R, Gougoulias N, Guthrie H et al. Morton's neuroma: clinical testing and imaging in 76 feet, compared to a control group. *Foot Ankle Surg* 2011;17(3):197–200.
137. Bignotti B, Signori A, Sormani MP et al. Ultrasound versus magnetic resonance imaging for Morton neuroma: systematic review and meta-analysis. *Eur Radiol* 2015;25(8):2254–62.
138. Xu Z, Duan X, Yu X et al. The accuracy of ultrasonography and magnetic resonance imaging for the diagnosis of Morton's neuroma: a systematic review. *Clin Radiol* 2015;70(4):351–8.
139. Morgan P, Monaghan W, Richards S. A systematic review of ultrasound-guided and non-ultrasound-guided therapeutic injections to treat Morton's neuroma. *J Am Podiatr Med Assoc* 2014;104(4):337–48.
140. Markovic M, Crichton K, Read JW et al. Effectiveness of ultrasound-guided corticosteroid injection in the treatment of Morton's neuroma. *Foot Ankle Int* 2008;29(5):483–7.
141. Fanucci E, Masala S, Fabiano S et al. Treatment of intermetatarsal Morton's neuroma with alcohol injection under US guide: 10-month follow-up. *Eur Radiol* 2004;14(3):514–8.
142. Hughes RJ, Ali K, Jones H et al. Treatment of Morton's neuroma with alcohol injection under sonographic guidance: follow-up of 101 cases. *AJR Am J Roentgenol* 2007;188(6):1535–9.
143. Chuter GS, Chua YP, Connell DA et al. Ultrasound-guided radiofrequency ablation in the management of interdigital (Morton's) neuroma. *Skeletal Radiol* 2013;42(1):107–11.
144. Friedman T, Richman D, Adler R. Sonographically guided cryoneurolysis: preliminary experience and clinical outcomes. *J Ultrasound Med* 2012;31(12):2025–34.
145. Villas C, Florez B, Alfonso M. Neurectomy versus neurolysis for Morton's neuroma. *Foot Ankle Int* 2008;29(6):578–80.
146. Pitsis GC, Best JP, Sullivan MR. Unusual stress fractures of the proximal phalanx of the great toe: a report of two cases. *Br J Sports Med* 2004;38:e31.

CHAPTER 44

1. Caine DJ, Maffulli N. Epidemiology of pediatric sports injuries. An important area of medicine and sports science research. In: Caine DJ, Maffulli N, eds. *Epidemiology of Pediatric Sports Injuries Individual Sports*. Basel: Karger, 2005:1–7.
2. Malina R, Bouchard C, Bar-Or O. *Growth, Maturation and Physical Activity*. 2nd ed. Champaign: Human Kinetics, 2004.
3. Beunen G, Malina R. Growth and biologic maturation: relevance to athletic performance. In: Hebestreit H, Bar-Or O, eds. *The Young Athlete Encyclopaedia of Sports Medicine (Vol XIII)*. Blackwell Publishing, Malden, MA, USA 2008:3–17.
4. Alexander CJ. Effect of growth rate on the strength of the growth plate-shaft junction. *Skel Radiol* 1976;1(2):67–76.
5. Peterson CA, Peterson HA. Analysis of the incidence of injuries to the epiphyseal growth plate. *J Trauma* 1972;12(4):275–81.
6. Bailey DA, Wedge JH, McCulloch RG et al. Epidemiology of fractures of the distal end of the radius in children as associated with growth. *J Bone Joint Surg Am* 1989;71(8):1225–31.
7. Micheli LJ, Fehlandt AF. Overuse injuries to tendons and apophyses in children and adolescents. *Clin Sports Med* 1992;11(4):713–26.
8. Feldman D, Shrier I, Rossignol M et al. Adolescent growth is not associated with changes in flexibility. *Clin J Sport Med* 1999;9(1):24–9.

9. Ogden JA. *Skeletal Injuries in the Child*. New York: Springer-Verlag, 2000.
10. Salter RB, Harris WR. Injuries involving the epiphyseal plate. *J Bone Joint Surg Am* 1963;45(3):587-622.
11. DiFiori JP, Caine DJ, Malina RM. Wrist pain, distal radial physeal injury, and ulnar variance in the young gymnast. *Am J Sports Med* 2006;34(5):840-9.
12. Frush TJ, Lindenfeld TN. Peri-epiphyseal and overuse injuries in adolescent athletes. *Sports Health* 2009;1(3):201-11.
13. Erickson BJ, Chalmers PN, Bush-Joseph CA et al. Ulnar Collateral Ligament Reconstruction of the Elbow: A Systematic Review of the Literature. *Orthop J Sports Med* 2015;3(12):2325967115618914. [Epub ahead of print]
14. Smith MV, Bedi A, Chen NC. Surgical treatment for osteochondritis dissecans of the capitellum. *Sports Health* 2012;4(5):425-32.
15. Caine D, Howe W, Ross W et al. Does repetitive physical loading inhibit radial growth in female gymnasts? *Clin J Sport Med* 1997;7(4):302-8.
16. Sward L. The thoracolumbar spine in young elite athletes. Current concepts on the effects of physical training. *Sports Med* 1992;13(5):357-64.
17. Herring JA. *Tachdjian's Pediatric Orthopaedics: From the Texas Scottish Rite Hospital for Children*. 5th ed. Philadelphia: W.B. Saunders, 2013.
18. Peck DM. Apophyseal injuries in the young athlete. *Am Fam Phys* 1995;51(8):1891-5.
19. Chaudhry S, Philips D, Feldman D. Legg-Calvé-Perthes disease. An overview with recent literature. *Bull Hosp Joint Dis* 2014;72(1):18-27.
20. Loder RT, Skopelja EN. The epidemiology and demographics of slipped capital femoral epiphysis. *ISRN Orthopedics* 2011;2011:Article ID 486512.
21. Kasper JC, Gerhardt MB, Mandelbaum BR. Stress injury leading to slipped capital femoral epiphysis in a competitive adolescent tennis player: a case report. *Clin J Sport Med* 2007;17(1):72-4.
22. Maffulli N, Bruns W. Injuries in young athletes. *Europ J Pediatr* 2000;159(1-2):59-63.
23. Gaida JE, Cook J. Treatment options for patellar tendinopathy: critical review. *Curr Sports Med Rep* 2011;10(5):255-70.
24. van Ark M, Docking SI, van den Akker-Scheek I et al. Does the adolescent patellar tendon respond to 5 days of cumulative load during a volleyball tournament? *Scand J Med Sci Sports* 2016;26(2):189-96.
25. Williamson L, Bowness P, Mowat A et al. Difficulties in diagnosing acute rheumatic fever-arthritis may be short lived and carditis silent. *BMJ* 2000;320(7231):362-5.
26. Kim SJ, Shim DW, Park KW. Functional outcome of transphyseal reconstruction of the anterior cruciate ligament in skeletally immature patients. *Knee Surg Relat Res* 2012;24(3):173-9.
27. Fabricant PD, Jones KJ, Delos D et al. Reconstruction of the anterior cruciate ligament in the skeletally immature athlete: a review of current concepts: AAOS exhibit selection. *J Bone Joint Surg Am* 2013;95(5):e28.
28. Goddard M, Bowman N, Salmon LJ et al. Endoscopic anterior cruciate ligament reconstruction in children using living donor hamstring tendon allografts. *Am J Sports Med* 2013;41(3):567-74.
29. Kumar S, Ahearne D, Hunt DM. Transphyseal anterior cruciate ligament reconstruction in the skeletally immature: follow-up to a minimum of sixteen years of age. *J Bone Joint Surg Am* 2013;95(1):e1.
30. Moksnes H, Engebretsen L, Eitzen I et al. Functional outcomes following a non-operative treatment algorithm for anterior cruciate ligament injuries in skeletally immature children 12 years and younger. A prospective cohort with 2 years follow-up. *Br J Sports Med* 2013;47(8):488-94.
31. Demange MK, Camanho GL. Nonanatomic anterior cruciate ligament reconstruction with double-stranded semitendinosus grafts in children with open physes: minimum 15-year follow-up. *Am J Sports Med* 2014;42(12):2926-32.
32. Calvo R, Figueroa D, Gili F et al. Transphyseal anterior cruciate ligament reconstruction in patients with open physes: 10-year follow-up study. *Am J Sports Med* 2015;43(2):289-94.
33. Lawrence JT, Argawal N, Ganley TJ. Degeneration of the knee joint in skeletally immature patients with a diagnosis of an anterior cruciate ligament tear: is there harm in delay of treatment? *Am J Sports Med* 2011;39(12):2582-7.
34. Kocher MS, Saxon HS, Hovis WD et al. Management and complications of anterior cruciate ligament injuries in skeletally immature patients: survey of the Herodicus Society and The ACL Study Group. *J Pediatr Orthop* 2002;22(4):452-7.
35. Caine D, DiFiori J, Maffulli N. Physeal injuries in children's and youth sports: reasons for concern? *Br J Sports Med* 2006;40(9):749-60.
36. Renstrom P, Ljungqvist A, Arendt E et al. Non-contact ACL injuries in female athletes: an International Olympic Committee current concepts statement. *Br J Sports Med* 2008;42(6):394-412.
37. Funahashi KM, Moksnes H, Maletis GB et al. Anterior cruciate ligament injuries in adolescents with open physis: effect of recurrent injury and surgical delay on meniscal and cartilage injuries. *Am J Sports Med* 2014;42(5):1068-73.
38. Frosch KH, Stengel D, Brodhun T et al. Outcomes and risks of operative treatment of rupture of the anterior cruciate ligament in children and adolescents. *Arthroscopy* 2010;26(11):1539-50.
39. Kaeding CC, Flanigan D, Donaldson C. Surgical techniques and outcomes after anterior cruciate ligament reconstruction in preadolescent patients. *Arthroscopy* 2010;26(11):1530-8.
40. Reider B. A matter of timing. *Am J Sports Med* 2015;43(2):273-4.
41. Mohtadi N, Grant J. Managing anterior cruciate ligament deficiency in the skeletally immature individual: a systematic review of the literature. *Clin J Sport Med* 2006;16(6):457-64.
42. Moksnes H, Engebretsen L, Risberg MA. The current evidence for treatment of ACL injuries in children is low: a systematic review. *J Bone Joint Surg Am* 2012;94(12):1112-9.

43. Chotel F, Henry J, Seil R et al. Growth disturbances without growth arrest after ACL reconstruction in children. *Knee Surg Sports Traumatol Arthrosc* 2010;18(11):1496-500.
44. Lee K, Siegel MJ, Lau DM et al. Anterior cruciate ligament tears: MR imaging-based diagnosis in a pediatric population. *Radiology* 1999;213(3):697-704.
45. Bollen S, Pease F, Ehrenraich A et al. Changes in the four-strand hamstring graft in anterior cruciate ligament reconstruction in the skeletally-immature knee. *J Bone Joint Surg Br* 2008;90(4):455-9.
46. Lafrance RM, Giordano B, Goldblatt J et al. Pediatric tibial eminence fractures: evaluation and management. *J Am Acad Orthop Surg* 2010;18(7):395-405.
47. Roth PB. Fracture of the spine of the tibia. *J Bone Joint Surg Am* 1928;10(3):509-18.
48. Lubowitz JH, Elson WS, Guttman D. Part II: arthroscopic treatment of tibial plateau fractures: intercondylar eminence avulsion fractures. *Arthroscopy* 2005;21(1):86-92.
49. Di Caprio F, Buda R, Ghermandi R et al. Combined arthroscopic treatment of tibial plateau and intercondylar eminence avulsion fractures. *J Bone Joint Surg Am* 2010;92 Suppl 2:161-9.
50. Moksnes H, Engebretsen L, Risberg MA. Management of anterior cruciate ligament injuries in skeletally immature individuals. *J Orthop Sports Phys Ther* 2012;42(3):172-83.
51. Faigenbaum AD, Kraemer WJ, Blimkie CJ et al. Youth resistance training: updated position statement paper from the National Strength and Conditioning Association. *J Strength Cond Res* 2009;23(5 Suppl):S60-79.
52. Hewett TE, Ford KR, Myer GD. Anterior cruciate ligament injuries in female athletes: part 2, a meta-analysis of neuromuscular interventions aimed at injury prevention. *Am J Sports Med* 2006;34(3):490-8.
53. Steffen K, Emery CA, Romiti M et al. High adherence to a neuromuscular injury prevention programme (FIFA 11+) improves functional balance and reduces injury risk in Canadian youth female football players: a cluster randomised trial. *Br J Sports Med* 2013;47(12):794-802.
54. McKay CD, Steffen K, Romiti M et al. The effect of coach and player injury knowledge, attitudes and beliefs on adherence to the FIFA 11+ programme in female youth soccer. *Br J Sports Med* 2014;48(17):1281-6.
55. Kvist J, Kartus J, Karlsson J et al. Results from the Swedish national anterior cruciate ligament register. *Arthroscopy* 2014;30(7):803-10.
56. Wiggins AJ, Grandhi RK, Schneider DK et al. Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Am J Sports Med* 2016;44(7):1861-76.
57. Funahashi KM, Moksnes H, Maletis GB et al. Anterior cruciate ligament injuries in adolescents with open physis: effect of recurrent injury and surgical delay on meniscal and cartilage injuries. *Am J Sports Med* 2014;42(5):1068-73.
58. Moksnes H, Engebretsen L. It takes more than timing: letter to the editor. *Am J Sports Med* 2015;43(6):NP 14-5.
59. Moksnes H, Grindem H. Prevention and rehabilitation of paediatric anterior cruciate ligament injuries. *Knee Surg Sports Traumatol Arthrosc* 2016;24(3):730-6.
60. Sever JW. Apophysitis of the os calcis. *NY Med J* 1912;95:1025.
61. Ogden JA, Ganey TM, Hill JD et al. Sever's injury: a stress fracture of the immature calcaneal metaphysis. *J Ped Orth* 2004;24(5):488-92.
62. Chiodo WA, Cook KD. Pediatric heel pain. *Clin Podiatr Med Surg* 2010;27(3):355-67 doi: 10.1016/j.cpm.2010.03.001. [Epub ahead of print]

CHAPTER 45

1. United States Bone and Joint Initiative. Summary Data. The Burden of Musculoskeletal Diseases in the United States (BMUS), 3rd ed. Rosemont, IL, 2014. <http://www.boneandjointburden.org/2014-report/vid1/summary-data>.
2. Geuze E, Westenberg HG, Jochims A et al. Altered pain processing in veterans with posttraumatic stress disorder. *Arch Gen Psychiatry* 2007;64(1):76-85.
3. Dar R, Arieli D, Frenk H. The effect of past-injury on pain threshold and tolerance. *Pain* 1995;60(2):189-93.
4. Harper P. No pain, no gain: pain behaviour in the armed forces. *Br J Nurs* 2006;15(10):548-51.
5. Davidson PL, Chalmers DJ, Wilson BD et al. Lower limb injuries in New Zealand Defence Force personnel: descriptive epidemiology. *Aust N Z J Public Health* 2008;32(2):167-73.
6. Jones BH, Knapik JJ. Physical training and exercise-related injuries. Surveillance, research and injury prevention in military populations. *Sports Med* 1999;27(2):111-25.
7. Kaufman KR, Brodine S, Shaffer R. Military training-related injuries: surveillance, research, and prevention. *Am J Prev Med* 2000;18(3 Suppl):54-63.
8. Lutz C. US and Coalition Casualties in Iraq and Afghanistan. Secondary US and Coalition Casualties in Iraq and Afghanistan 2013. 21 Feb 2013. Watson Institute for International Studies, Brown University. <http://watson.brown.edu/costsofwar/files/cow/imce/papers/2013/USandCoalition.pdf>.
9. Belmont PJ, Owens BD, Schoenfeld AJ. Musculoskeletal injuries in Iraq and Afghanistan: epidemiology and outcomes following a decade of war. *J Am Acad Orthop Surg*. 2016;24(6):341-8. doi: 10.5435/JAAOS-D-15-00123.
10. Belmont PJ, Jr., McCrisky BJ, Hsiao MS et al. The nature and incidence of musculoskeletal combat wounds in Iraq and Afghanistan (2005-2009). *J Orthop Trauma* 2013;27(5):e107-13. doi: 10.1097/BOT.0b013e3182703188.
11. Hauret KG, Taylor BJ, Clemmons NS et al. Frequency and causes of nonbattle injuries air evacuated from operations Iraqi Freedom and Enduring Freedom, U.S. Army, 2001-2006. *Am J Prev Med* 2010;38(1 Suppl):S94-107.
12. Reynolds KL, Heckel HA, Witt CE et al. Cigarette smoking, physical fitness, and injuries in infantry soldiers. *Am J Prev Med* 1994;10(3):145-50.

13. Hauret KG, Jones BH, Bullock SH et al. Musculoskeletal injuries; description of an under-recognized injury problem among military personnel. *Am J Prev Med* 2010;38(1 Suppl):S61-70.
14. Rudzki SJ. Injuries in Australian Army recruits. Part I: decreased incidence and severity of injury seen with reduced running distance. *Mil Med* 1997;162:472-6.
15. Jordaan G, Schwellnus MP. The incidence of overuse injuries in military recruits during basic military training. *Mil Med* 1994;159(6):421-6.
16. Heir T, Eide G. Age, body composition, aerobic fitness and health condition as risk factors for musculoskeletal injuries in conscripts. *Scand J Med Sci Sports* 1996;6(4):222-7.
17. Penn-Barwell J, Fries CA, Bennett PM et al. Mortality, survival and residual injury burden of Royal Navy and Royal Marine combat casualties sustained in 11-years of operations in Iraq and Afghanistan. *J R Nav Med Serv* 2014;100(2):161-5.
18. Workcover Authority New South Wales. Workers Compensation statistical bulletin. Sydney: Workcover, 1998.
19. Roy TC, Lopez HP. A comparison of deployed occupational tasks performed by different types of military battalions and resulting low back pain. *Mil Med* 2013;178(8):e937-43. doi: 10.7205/MILMED-D-12-00539.
20. Konitzer LN, Fargo MV, Brininger TL et al. Association between back, neck, and upper extremity musculoskeletal pain and the individual body armor. *J Hand Ther* 2008;21(2):143-8; quiz 149. doi: 10.1197/j.jht.2007.10.017.
21. Smith TA, Cashman TM. The incidence of injury in light infantry soldiers. *Mil Med* 2002;167(2):104-8.
22. Vann RD, Vorosmarti J. Military diving operations and medical support. Military preventive medicine: mobilization and deployment: Department of Defense, Office of The Surgeon General, US Army, Borden Institute, 2003:955-94.
23. Broadhurst RS, Morrison LJ, Howsare CR et al. Military diving medicine. Military preventive medicine: mobilization and deployment: Department of Defense, Office of The Surgeon General, US Army, Borden Institute, 2003:575-610.
24. Diebal AR, Gregory R, Alitz C et al. Forefoot running improves pain and disability associated with chronic exertional compartment syndrome. *Am J Sports Med* 2012;40(5):1060-7. doi: 10.1177/0363546512439182.
25. Cornelius A. Keeping runners running: the secrets of running assessment—advice and exercise progressions. BJSM podcasts: *BMJ*, 2015. <http://ow.ly/eHYD304BgdQ> (accessed September 26th, 2016)
26. Franklyn-Miller A, Roberts A, Hulse D et al. Biomechanical overload syndrome: defining a new diagnosis. *Br J Sports Med* 2014;48(6):415-6.
27. Protzman RR, Griffis CC. Comparative stress fracture incidence in males and females in an equal training environment. *Athletic Training* 1977;12:126-30.
28. Reinker KA, Ozbourne S. A comparison of male and female basic orthopaedic pathology in basic training. *Mil Med* 1979;144:532-6.
29. Brudvig TJ, Gudjer TD, Obermeyer L. Stress fractures in 259 trainees: a one-year study of incidence as related to age, sex and race. *Mil Med* 1983;148:666-9.
30. Jones BH, Harris JM, Tuyethoa NV et al. Exercise-induced stress fractures and stress reactions of bone: epidemiology, etiology and classification. In: Holloszy JO, ed. *Exercise and Sport Science Review*. Baltimore: Williams and Wilkins, 1989:379-422.
31. Larson NS, Amin R, Olsen C et al. Effect of diet cola on urine calcium excretion. *Endocrine Rev* 2010;31(3)(1 Suppl):P2-198.
32. Zhao L, Chang Q, Huang T et al. Prospective cohort study of the risk factors for stress fractures in Chinese male infantry recruits. *J Int Med Res* 2016;44(4):787-95.
33. Shaffer RA, Rauh MJ, Brodine SK et al. Predictors of stress fracture susceptibility in young female recruits. *Am J Sports Med* 2006;34(1):108-15.
34. Research into foot lesions among Canadian field forces. 13th Commonwealth Defence Conference on Operational Clothing and Combat Equipment. Malaysia, 1981.
35. Dalen A, Nilsson J, Thorstensson A. *Factors influencing a prolonged foot march*. Stockholm, Sweden: Karolinska Institute, 1978.
36. Benseck CK, Kish R. *Lower extremity disorders among men and women in Army Basic training and the effects of two types of boots*. US Army Natick Research & Development Laboratories, Natick, MA: p 1-99, 1983.
37. Kelly RA. The incidence and prevention of foot problems among male phase one British Army recruits at an Army Training Regiment. *J R Army Med Corps* 2015;161(1 Suppl):i56-i59.
38. Herring KM, Richie DH. Friction blisters and sock fibre composition. *J Am Podiatr Med Assoc* 1990;80:63-71.
39. Kinoshita H. Effects of different loads and carrying systems on selected biomechanical parameters describing gait. *Ergonomics* 1985;28:1347-62.
40. Allan JR. *A study of foot blisters*. United Kingdom: Army Operational Research Establishment (AORE), 1964.
41. Bullock SH, Jones BH, Gilchrist J et al. Prevention of physical training-related injuries recommendations for the military and other active populations based on expedited systematic reviews. *Am J Prev Med* 2010;38(1):S156-81.
42. Thompson KJ, Hamlet MP, Jones BH et al. *Impact of sock systems on frequency and severity of blister injury in a Marine recruit population*. Natick, MA: USARIEM, 1993.
43. Quinn J. *The effects of two new foot powders on the incidence of foot infection and blisters in recruits during basic training*. Farnborough, UK: Army Personnel Research Establishment (APRE), 1967.
44. Military footwear and the occurrence of blisters. 8th Commonwealth Defence Conference On Clothing and General Stores. Melbourne, 1965.
45. Whittingham DGV. *A further investigation of marching in flying boots: Exercise Orthoped II*. Farnborough UK: RAF Institute of Aviation Medicine, 1951.
46. Kragh JF, Jr., Jones BH, Amaroso PJ et al. Parachuting injuries among Army Rangers: a prospective survey of an elite airborne battalion. *Mil Med* 1996;161(7):416-9.

47. Hughes CD, Weinrauch PC. Military static line parachute injuries in an Australian commando battalion. *ANZ J Surg* 2008;78(10):848-52.
48. Amoroso PJ, Ryan JB, Bickley B et al. Braced for impact: reducing military paratroopers' ankle sprains using outside-the-boot braces. *J Trauma* 1998;45(3):575-80.
49. Schumacher JT, Jr., Creedon JF, Pope RW. The effectiveness of the parachutist ankle brace in reducing ankle injuries in an airborne ranger battalion. *Mil Med* 2000;165(12):944-8.
50. Knapik JJ, Spiess A, Swedler DI et al. Systematic review of the parachute ankle brace: injury risk reduction and cost effectiveness. *Am J Prev Med* 2010;38(1 Suppl):S182-8.
51. Pope RP, Herbert R, Kirwan JD et al. Predicting attrition in basic military training. *Mil Med* 1999;164(10):710-4.
52. Knapik JJ, Canham-Chervak M, Hauret KG et al. Discharges during US Army basic training: injury rates and risk factors. *Mil Med* 2001;166(7):641-7.
53. Pointer S, Harrison J, Bradley C. *National Injury Prevention Plan priorities for 2004 and beyond*. Canberra: Australian Institute of Health and Welfare, 2003.
54. National Center for Injury Prevention and Control. *Violence prevention: A Public Health study*. Geneva: WHO Collaborating Centre on Injury Control, 1998.
55. Pope RP. Injury surveillance and systematic investigation identify a rubber matting hazard for anterior cruciate ligament rupture on an obstacle course. *Mil Med* 2002;167(4):359-62.
56. Pope RP. Rubber matting on an obstacle course causes anterior cruciate ligament ruptures and its removal eliminates them. *Mil Med* 2002;167(4):355-8.
57. Langton N, Khan KM, Lusina SJ. FIFA's Football for Health: applying Kotter's eight-step programme for transformational change to a mass participation activity. *Br J Sports Med* 2010;44(8):537-9.
58. Kotter J. Leading change: why transformation efforts fail. *Harvard Bus Rev* 2007;96-103.
59. Darakjy S, Marin RE, Knapik JJ et al. Injuries and illnesses among armor brigade soldiers during operational training. *Mil Med* 2006;171(11):1051-6.
60. Rudzki SJ, Cunningham MJ. The effect of a modified physical training program in reducing injury and medical discharge rates in Australian Army recruits. *Mil Med* 1999;164(9):648-52.
61. Blacker SD, Wilkinson DM, Bilzon JL et al. Risk factors for training injuries among British Army recruits. *Mil Med* 2008;173(3):278-86.
62. Blacker S, Wilkinson D, Rayson M. Gender differences in the physical demands of British Army recruit training. *Mil Med* 2009;174(8):811-6.
63. Jones BH, Bovee M, Knapik J. *Associations among body composition, physical fitness and injury in men and women army trainees. Body composition and physical performance: applications for the military services*. Washington DC: National Academy Press, 1992:141-73.
64. Tomasi LF, Peterson JH, Pettit GP et al. Women's response to Army training. *Phys Sportsmed* 1977;5(6):32-7.
65. Daniels WL, Kowal DM, Vogel JA et al. Physiological effects of a military training program on male and female cadets. *Aviat Space Environ Med* 1979;50(6):562-6.
66. Knapik JJ, Sharp MA, Canham-Chervak M et al. Risk factors for training-related injuries among men and women in basic combat training. *Med Sci Sports Exerc* 2001;33(6):946-54.
67. Leger LA, Gadoury C. Validity of the 20 m shuttle run test with 1 Min stages to predict V02 max in adults. *Can J Sport Sci* 1989;14:21-6.
68. Paliczka VJ, Nichols AK, Boreham CAG. A multi-stage shuttle run as a predictor of running performance and maximal oxygen uptake in adults. *Br J Sports Med* 1987;21:163-5.
69. Allsopp AJ, Scarpello EG, Andrews S et al. Survival of the fittest? The scientific basis for the Royal Navy pre-joining fitness test. *J R Nav Med Serv* 2003;89(1):11-8.
70. Pope RP. Prediction and prevention of lower limb injuries and attrition in Army recruits. Thesis. Wagga Wagga, NSW: Charles Sturt University, 2002.
71. Burke BG, Sauser WT, Kemery ER et al. Intelligence and physical fitness as predictors of success in early infantry training. *Percept Mot Skills* 1989;69:263-71.
72. Lunt H. A pre-joining fitness test improves pass rates of Royal Navy recruits. *Occup Med* 2007;57(5):377-9.
73. Knapik JJ, Sharp MA, Darakjy S et al. Temporal changes in the physical fitness of US Army recruits. *Sports Med* 2006;37(7):613-34.
74. Niebuhr DW, Scott CT, Li Y et al. Preaccession fitness and body composition as predictors of attrition in US Army recruits. *Mil Med* 2009;174(7):695-701.
75. Marti B, Vader JP, Minder CE et al. On the epidemiology of running injuries. *Am J Sports Med* 1988;16:285-94.
76. Macera CA, Pate RR, Powell KE et al. Predicting lower extremity injuries among habitual runners. *Arch Intern Med* 1989;149:2565-8.
77. Walter SD, Hart LE, McIntosh JM et al. The Ontario Cohort study of running related injuries. *Arch Int Med* 1989;149:2561-4.
78. Taunton JE, Ryan MB, Clement DB et al. A prospective study of running injuries: the Vancouver Sun Run 'In Training' clinics. *Br J Sports Med* 2003;37(3):329-44.
79. Powell KE, Kohl HW, Casperson CJ et al. An epidemiological perspective on the causes of running injuries. *Phys Sportsmed* 1986;14(6):100-14.
80. Ekstrand J. Soccer injuries and their prevention. Medical Dissertations no. 130. Linköping University, 1982.
81. Goss DL, Christopher GE, Faulk RT et al. Functional training program bridges rehabilitation and return to duty. *J Spec Oper Med* 2009;9(2):29-48.
82. Gijon-Nogueron G, Fernandez-Villarejo M. Risk Factors and Protective Factors for Lower-Extremity Running Injuries A Systematic Review. *J Am Podiatr Med Assoc* 2015;105(6):532-40.
83. Wright AA, Taylor JB, Ford KR et al. Risk factors associated with lower extremity stress fractures in runners: a

- systematic review with meta-analysis. *Br J Sports Med* 2015;49(23):1517-23.
84. Koplan JP, Powell KE, Sikes RK et al. An epidemiological study of the benefits and risks of running. *JAMA* 1982;248:3118-21.
 85. Jacobs SJ, Berson BL. Injuries to runners: a study of entrants to a 10 000m race. *Am J Sports Med* 1986;14:151-5.
 86. Bovens AM, Janssen GM, Vermeer HG et al. Occurrence of running injuries in adults following a supervised training programme. *Int J Sports Med* 1989;10:S186-90.
 87. Lysholm J, Wiklander J. Injuries in runners. *Am J Sports Med* 1987;15:168-71.
 88. Kowal DM. Nature and causes of injuries in women resulting from an endurance training programme. *Am J Sports Med* 1980;8:265-9.
 89. Andrews JR. Overuse syndromes of the lower extremity. *Clin in Sports Med* 1983;2:137-48.
 90. Pollock ML, Gettmann LR, Milesis LA et al. Effects of frequency and duration of training on attrition and incidence of injury. *Med Sci Sports* 1977;9:31-6.
 91. Gabbett TJ. The training-injury prevention paradox: should athletes be training smarter and harder? *Br J Sports Med* 2016;50(5):273-80.
 92. Musculoskeletal injury project. Cincinnati, OH: American College of Sports Medicine, 1996:29 May-1 June.
 93. Trank TV, Ryman DH, Minagawa RY et al. Running mileage, movement mileage, and fitness in male US Navy recruits. *Med Sci Sports Exerc* 2001;33(6):1033-8.
 94. Knapik JJ, Hauret KG, Arnold S et al. Injury and fitness outcomes during implementation of physical readiness training. *Int J Sports Med* 2003;24(5):372-81.
 95. Jones BH, Cowan DN, Tomlinson JP et al. Epidemiology of injuries associated with physical training among young men in the Army. *Med Sci Sports Exerc* 1993;25(2):197-203.
 96. UASCHPPM. Recommendations for the prevention of physical training (PT)-related injuries: Results of a systematic evidence based review by the Joint Services Physical Training Injury Prevention Working Group (JSPTIPWG). Aberdeen Proving Ground, MD: United States Army Centre for Health Promotion and Preventative Medicine, 2008.
 97. Rudzki SJ. Weight-load marching as a method of conditioning Australian Army recruits. *Mil Med* 1989;154(4):201-5.
 98. Pope RP. Prevention of pelvic stress fractures in female army recruits. *Mil Med* 1999;164(5):370-3.
 99. Knapik JJ, Darakjy S, Scott SJ et al. Evaluation of a standardized physical training program for basic combat training. *J Strength Cond Res* 2005;19(2):246-53.
 100. Burns AS, Lauder TD. Deep water running: an effective non-weightbearing exercise for the maintenance of land-based running performance. *Mil Med* 2001;166(3):253-8.
 101. Knapik JJ, Darakjy S, Hauret KG et al. Increasing the physical fitness of low-fit recruits before basic combat training: an evaluation of fitness, injuries, and training outcomes. *Mil Med* 2006;171(1):45-54.
 102. Van Mechelen W. Running injuries: a review of the epidemiological literature. *Sports Med* 1992;14(5):320-35.
 103. Yzerman JC, VanGalen WC. Blessures bij lange afstandlopers. KNAU, 1987.
 104. McHugh MP, Cosgrave CH. To stretch or not to stretch; the role of stretching in injury prevention and performance. *Scand J Med Sci Sports* 2010;20(1):169-81.
 105. Shrier I. Does stretching improve performance? A systematic and critical review of the literature. *Clin J Sport Med* 2004;14(5):267-73.
 106. Small K, McNaughton L, Matthews M. A systematic review into the efficacy of static stretching as part of a warm-up for the prevention of exercise-related injury. *Res Sports Med* 2008;16(3):213-31.
 107. Pope RP, Herbert RD, Kirwan JD et al. A randomized trial of preexercise stretching for prevention of lower-limb injury. *Med Sci Sports Exerc* 2000;32(2):271-7.
 108. Amako M, Oda T, Masuoka K et al. Effect of static stretching on prevention of injuries for military recruits. *Mil Med* 2003;168(6):442-6.
 109. Fletcher IM, Jones BH. The effect of different warm-up stretch protocols on 20 meter sprint performance in trained rugby union players. *J Strength Cond Res* 2004;18(4):885-8.
 110. Jagers JR, Swank AM, Frost KL et al. The acute effects of dynamic and ballistic stretching on vertical jump height, force, and power. *J Strength Cond Res* 2008;22(6):1844-9.
 111. Fletcher IM, Anness R. The acute effects of combined static and dynamic stretch protocols on fifty-meter sprint performance in track-and-field athletes. *J Strength Cond Res* 2007;21(3):784-7.

CHAPTER 46

1. Dvorak J, Grimm K, Schmied C et al. Development and implementation of a standardised precompetition medical assessment of international elite football players—2006 FIFA World Cup Germany. *Clin J Sport Med* 2009;19(4):316-21.
2. Ljungqvist A, Jenoure P, Engebretsen L et al. The International Olympic Committee (IOC) Consensus Statement on periodic health evaluation of elite athletes March 2009. *Br J Sports Med* 2009;43(9):631-43.
3. American Academy of Paediatrics. Preparticipation physical evaluation form. 2010. <https://www.aap.org/en-us/professional-resources/practice-support/pages/Preparticipation-Physical-Evaluation-Forms.aspx>, accessed 30 Dec 2015.
4. Bernhardt DT, Roberts WO, American Academy of Family Physicians et al. *PPE: Preparticipation Physical Evaluation*. 4th ed. Elk Grove Village, IL: American Academy of Paediatrics, 2010.
5. Maron BJ, Zipes DP. Introduction: eligibility recommendations for competitive athletes with cardiovascular abnormalities—general considerations. *J Am Coll Cardiol* 2005;45(8):1318-21.
6. Pelliccia A, Fagard R, Bjornstad HH et al. Recommendations for competitive sports participation in athletes with cardiovascular disease: a consensus document from the Study Group of Sports

- Cardiology of the Working Group of Cardiac Rehabilitation and Exercise Physiology and the Working Group of Myocardial and Pericardial Diseases of the European Society of Cardiology. *Eur Heart J* 2005;26(14):1422-45.
7. Bakken A, Targett S, Bere T et al. Health conditions detected in a comprehensive periodic health evaluation of 558 professional football players. *Br J Sports Med* 2016. Sep 6. pii: bjsports-2016-096570. doi: 10.1136/bjsports-2016-096570. [Epub ahead of print]
 8. Matheson GO, Anderson S, Robell K. Injuries and illnesses in the preparticipation evaluation data of 1693 college student-athletes. *Am J Sports Med* 2015;43(6):1518-25.
 9. National Adolescent Health Information Centre. Fact sheet on Mortality: adolescents and young adults. San Francisco: University of California, 2006.
 10. Sassi RH, Dardouri W, Yahmed MH et al. Relative and absolute reliability of a modified agility T-test and its relationship with vertical jump and straight sprint. *J Strength Cond Res* 2009;23(6):1644-51.
 11. Wilson JM, Jungner YG. Principles and practice of mass screening for disease. *Bol Oficina Sanit Panam* 1968;65(4):281-393.
 12. Bushman TT, Grier TL, Canham-Chervak MC et al. Pain on functional movement screen tests and injury risk. *J Strength Cond Res* 2015;29 Suppl 11:S65-70.
 13. Whiteley R. 'Moneyball' and time to be honest about preseason screening: it is a sham making no inroads on the 1 billion dollar injury costs in baseball. *Br J Sports Med* 2016;50(14):835-6.
 14. Bahr R. Can electrocardiographic screening prevent sudden death in athletes? No. *BMJ* 2010;341:c4914.
 15. Bahr R. Why screening tests to predict injury do not work—and probably never will . . . : a critical review. *Br J Sports Med* 2016;50(13):776-80.
 16. Drezner JA, Asif IM, Owens DS et al. Accuracy of ECG interpretation in competitive athletes: the impact of using standardised ECG criteria. *Br J Sports Med* 2012;46(5):335-40.
 17. Irwin ME, Drezner J, Price D et al. ECG screening does not cause undue anxiety in athletes undergoing pre-participation evaluation. *J Am Coll Cardiol* 2014;63(12S).
 18. Asif IM, Price D, Harmon KG et al. The psychological impact of cardiovascular screening in young athletes: perspectives across age, race and gender. *Clin J Sport Med* 2015;25(6):464-71.
 19. Drezner JA, Toresdahl BG, Rao AL et al. Outcomes from sudden cardiac arrest in US high schools: a 2-year prospective study from the National Registry for AED Use in Sports. *Br J Sports Med* 2013;47(18):1179-83.
 20. Lappe J, Cullen D, Haynatzki G et al. Calcium and vitamin D supplementation decreases incidence of stress fractures in female navy recruits. *J Bone Miner Res* 2008;23(5):741-9.
 21. Cook G, Burton L, Hoogenboom BJ et al. Functional movement screening: the use of fundamental movements as an assessment of function—part 2. *Int J Sports Phys Ther* 2014;9(4):549-63.
 22. Cook G. *Athletic Body in Balance—Optimal Movement Skills and Conditioning for Performance*. Champaign, IL: Human Kinetics, 2004.
 23. Cook G. *Movement: Functional Movement Systems: Screening, Assessment, Corrective Strategies*. Aptos, CA: On-Target Publications, 2010.
 24. United States Tennis Association. The USTA High-Performance Profile. <https://s3.amazonaws.com/ustaassets/assets/1/15/11940/highperformance.profile.lowres.pdf>
 25. Frohm A, Heijne A, Kowalski J et al. A nine-test screening battery for athletes: a reliability study. *Scand J Med Sci Sports* 2012;22(3):306-15.
 26. van Beijsterveldt AM, van de Port IG, Vereijken AJ et al. Risk factors for hamstring injuries in male soccer players: a systematic review of prospective studies. *Scand J Med Sci Sports* 2013;23(3):253-62.
 27. Dallinga JM, Benjaminse A, Lemmink KA. Which screening tools can predict injury to the lower extremities in team sports? A systematic review. *Sports Med* 2012;42(9):791-815.
 28. Opar DA, Williams MD, Timmins RG et al. Eccentric hamstring strength and hamstring injury risk in Australian footballers. *Med Sci Sports Exerc* 2015;47(4):857-65.
 29. Ruotolo C, Price E, Panchal A. Loss of total arc of motion in collegiate baseball players. *J Shoulder Elbow Surg* 2006;15(1):67-71.
 30. Myers JB, Laudner KG, Pasquale MR et al. Glenohumeral range of motion deficits and posterior shoulder tightness in throwers with pathologic internal impingement. *Am J Sports Med* 2006;34(3):385-91.
 31. Shanley E, Rauh MJ, Michener LA et al. Shoulder range of motion measures as risk factors for shoulder and elbow injuries in high school softball and baseball players. *Am J Sports Med* 2011;39(9):1997-2006.
 32. Clarsen B, Bahr R, Andersson SH et al. Reduced glenohumeral rotation, external rotation weakness and scapular dyskinesis are risk factors for shoulder injuries among elite male handball players: a prospective cohort study. *Br J Sports Med* 2014;48(17):1327-33.
 33. Cook C. Predicting future physical injury in sports: it's a complicated dynamic system. *Br J Sports Med* 2016. Jun 10. pii: bjsports-2016-096445. doi: 10.1136/bjsports-2016-096445. [Epub ahead of print]
 34. Walden M, Häggglund M, Ekstrand J. UEFA Champions League study: a prospective study of injuries in professional football during the 2001-2002 season. *Br J Sports Med* 2005;39(8):542-6.
 35. Brooks JH, Fuller CW, Kemp SP et al. Epidemiology of injuries in English professional rugby union: part 1 match injuries. *Br J Sports Med* 2005;39(10):757-66.
 36. LeFevre ML, Force USPST. Screening for hepatitis B virus infection in nonpregnant adolescents and adults: US Preventive Services Task Force recommendation statement. *Ann Intern Med* 2014;161(1):58-66.
 37. Risser WL, Hoffman HM, Bellah GG, Jr. Frequency of preparticipation sports examinations in secondary school athletes: are the University Interscholastic League guidelines appropriate? *Tex Med* 1985;81(7):35-9.



38. Carek PJ, Futrell M, Hueston WJ. The preparticipation physical examination history: who has the correct answers? *Clin J Sport Med* 1999;9(3):124-8.
39. Mountjoy M, Sundgot-Borgen J, Burke L et al. The IOC consensus statement: beyond the female athlete triad—relative energy deficiency in sport (RED-S). *Br J Sports Med* 2014;48(7):491-7.
40. Myklebust G, Skjølberg A, Bahr R. ACL injury incidence in female handball 10 years after the Norwegian ACL prevention study: important lessons learned. *Br J Sports Med* 2013;47(8):476-9.
41. Maron BJ, Levine BD, Washington RL et al. Eligibility and disqualification recommendations for competitive athletes with cardiovascular abnormalities: task force 2: preparticipation screening for cardiovascular disease in competitive athletes: a scientific statement from the American Heart Association and American College of Cardiology. *J Am Coll Cardiol* 2015;66(21):2356-61.
42. Carey JL, Huffman GR, Parekh SG et al. Outcomes of anterior cruciate ligament injuries to running backs and wide receivers in the National Football League. *Am J Sports Med* 2006;34(12):1911-7.
43. Brophy RH, Chehab EL, Barnes RP et al. Predictive value of orthopaedic evaluation and injury history at the NFL Combine. *Med Sci Sports Exerc* 2008;40(8):1368-72.
13. Dunn WR, George MS, Churchill L et al. Ethics in sports medicine. *Am J Sports Med* 2007;35(5):840-4.
14. Turbeville SD, Cowan LD, Greenfield RA. Infectious disease outbreaks in competitive sports: a review of the literature. *Am J Sports Med* 2006;34:1860-5.
15. Khan K, Freifeld CC, Wang J et al. Preparing for infectious disease threats at mass gatherings: the case of the Vancouver 2010 Olympic Winter Games. *CMAJ* 2010;182(6):579-83.
16. Maughan RJ. Contamination of dietary supplements and positive drug tests in sport. *J Sports Sci* 2005;23(9):883-9.
17. Milne CJ, Shaw MT. Travelling to China for the Beijing 2008 Olympic Games. *Br J Sports Med* 2008;42:321-6.
18. Milne C, Shaw M, Steinweg J. Medical issues relating to the Sydney Olympic Games. *Sports Med* 1999;28:287-98.
19. Blumberg LH, de Frey A, Frean J et al. The 2010 FIFA World Cup: communicable disease risks and advice for visitors to South Africa. *J Travel Med* 2010;17(3):150-2.
20. Muza SR, Beidleman BA, Fulco CS. Altitude preexposure recommendations for inducing acclimatization. *High Alt Med Biol* 2010;11(2):87-92.
21. Igreja RP. Olympics in the tropics and infectious diseases. *Clin Infect Dis* 2010;50(4):616-17.
22. Pipe AL. International travel and the elite athlete. *Clin J Sport Med* 2011;21:62-6.
23. Derman WE. Medication use by Team South Africa during the XXVIlth Olympiad: a model for quantity estimation for multi-coded team events. *S Afr J Sports Med* 2008;20(3):78-84.
24. Shaw M, Leggat PA. Traveling to Australia for the Sydney 2000 Olympic and Paralympic Games. *J Travel Med* 2000;7:200-4.
25. Reilly T, Waterhouse J, Burke LM et al. Nutrition for travel. *J Sports Sci* 2007;25 Suppl 1:S125-34.
26. Simon LM, Rubin AL. Traveling with the Team. *Curr Sports Med Rep* 2008;7:138-43.
27. Walters A. Travel medicine advice to UK based international motor sport teams. *J Travel Med* 2000;7:267-74.
28. Teichman PG, Donchin Y, Kot RJ. International aeromedical evacuation. *N Engl J Med* 2007;356:262-70.
29. Samuels C. Sleep, recovery, and performance: the new frontier in high-performance athletics. *Neurol Clin* 2009;20(1):149-59, ix.
30. Geertsema C, Williams AB, Dzendrowskyj P et al. Effect of commercial airline travel on oxygen saturation in athletes. *Br J Sports Med* 2008;42(11):577-81.
31. Sack RL. The pathophysiology of jet lag. *Travel Med Infect Dis* 2009;7(2):102-10.
32. Eastman CI, Burgess HJ. How to travel the world without jet lag. *Sleep Med Clin* 2009;4(2):241-55.
33. Winter WC, Hammond WR, Green NH et al. Measuring circadian advantage in Major League Baseball: a 10-year retrospective study. *Int J Sports Physiol Perform* 2009;4(3):394-401.
34. Reilly T, Edwards B. Altered sleep-wake cycles and physical performance in athletes. *Physiol Behav* 2007;90(2-3):274-84.
35. Arendt J. Managing jet lag: some of the problems and possible new solutions. *Sleep Med Rev* 2009;13(4):249-56.

CHAPTER 47

1. Brukner P. Surviving 30 years on the road as a team physician. *Br J Sports Med* 2013;47:10.
2. Milne C. New Zealand Olympic experience—Sydney 2000. *Br J Sports Med* 2001;35(4):281.
3. Waterhouse J, Reilly T, Edwards B. The stress of travel. *J Sports Sci* 2004;22(10):946-65; discussion 65-6.
4. Batt ME, Maryon-Davis A. Sport and exercise medicine: a timely specialty development. *Clin J Sport Med* 2007;17(2):85-6.
5. Dubey SG, Roberts C, Adebajo AO et al. Rheumatology training in the United Kingdom: the trainees' perspective. *Rheumatology* 2004;43(7):896-900.
6. Speed C, Jaques R. High-performance sports medicine: an ancient but evolving field. *Br J Sports Med* 2011;45(2):81-3.
7. Goforth M, Almquist J, Matney M et al. Understanding organization structures of the college, university, high school, clinical, and professional settings. *Clin Sports Med* 2007;26(2):201-26.
8. Menetrey J. Medical supervision of sports events and the role of the team physician. *Rev Med Suisse* 2008;4(166):1717-18, 1720-2.
9. Anderson L. Contractual obligations and the sharing of confidential health information in sport. *J Med Ethics* 2008;34(9):e6.
10. Tucker AM. Ethics and the professional team physician. *Clin Sports Med* 2004;23(2):227-41, vi.
11. Fu FH, Tjoumakaris FP, Buoncristiani A. Building a sports medicine team. *Clin Sports Med* 2007;26(2):173-9.
12. Milne C. The doctor's hip bag: advice sheet. *NZ J Sports Med* 2003;31.

36. Siedenburg J, Perry I, Stüben U. Tropical medicine and travel medicine: medical advice for aviation medical examiners concerning flight operations in tropical areas. *Aviat Space Environ Med* 2005;76(3 Suppl):A1-30.
37. Waterhouse J, Reilly T. Managing jet lag. *Sleep Med Rev* 2009;13(4):247-8.
38. Reilly T, Waterhouse J, Edwards B. Jet lag and air travel: implications for performance. *Clin Sports Med* 2005;24(2):367-80.
39. Postolache TT, Hung TM, Rosenthal RN et al. Sports chronobiology consultation: from the lab to the arena. *Clin Sports Med* 2005;24(2):415-56, xiv.
40. Waterhouse J, Reilly T, Atkinson G et al. Jet lag: trends and coping strategies. *Lancet* 2007;369(9567):1117-29.

CHAPTER 48

1. Lee IM, Shiroma EJ, Lobelo F et al. Effect of physical inactivity on major non-communicable diseases worldwide: an analysis of burden of disease and life expectancy. *Lancet* 2012;380(9838):219-29.
2. Boyd JL. Understanding the politics of being a team physician. *Clin Sports Med* 2007;26(2):161-72.