

9. Shrier I. Strategic Assessment of Risk and Risk Tolerance (StARRT) framework for return-to-play decision-making. *Br J Sports Med* 2015;49(20):1311-5.
10. Shrier I, Matheson GO, Boudier-Reveret M et al. Validating the three-step return-to-play decision model. *Scand J Med Sci Sports* 2015;25(2):e231-9.
11. Reyna VF, Rivers SE. Current theories of risk and rational decision making. *Dev Rev* 2008;28(1):1-11.
12. Ashby D, Smith AFM. Evidence-based medicine as Bayesian decision-making. *Stat Med* 2000;19(23):3291-305.
13. Shakespeare TP, Gebiski VJ, Veness MJ et al. Improving interpretation of clinical studies by use of confidence levels, clinical significance curves, and risk-benefit contours. *Lancet* 2001;357(9265):1349-53.
14. Shrier I, Safai P, Charland L. Return to play following injury: whose decision should it be? *Br J Sports Med* 2014;48(5):394-401.
15. National Heart, Lung, and Blood Institute, NIH. Risk assessment tool for estimating your 10-year risk of having a heart attack. <http://cvdrisk.nhlbi.nih.gov>.
16. Gabe J, Bury M, Elston MA. *Key concepts in medical sociology*. Thousand Oaks, CA: Sage Publications, 2004.
17. Safai P. Healing the body in the 'culture of risk': examining the negotiation of treatment between sport medicine clinicians and injured athletes in Canadian intercollegiate sport. *Sociol Sport J* 2003;20(2):127-46.
18. McFarland EG. Return to play. *Clin Sports Med* 2004;23(3): xv-xxiii.
19. Young K, Mcteer W, White P. Body talk—Male athletes reflect on sport, injury, and pain. *Sociol Sport J* 1994;11(2):175-94.
20. Podlog L, Eklund RC. A longitudinal investigation of competitive athletes' return to sport following serious injury. *J Appl Sport Psychol* 2006;18(1):44-68.
21. Podlog L, Eklund RC. The psychosocial aspects of a return to sport following serious injury: a review of the literature from a self-determination perspective. *Psychol Sport Exerc* 2007;8(4):535-66.
22. Nixon HL. Accepting the risks of pain and injury in sport—mediated cultural influences on playing hurt. *Sociol Sport J* 1993;10(2):183-96.
23. Barry MJ, Edgman-Levitan S. Shared decision making—pinnacle of patient-centered care. *N Engl J Med* 2012;366(9):780-1.
24. Nixon HL. Coaches' views of risk, pain, and injury in sport, with special reference to gender differences. *Sociol Sport J* 1994;11(1):79-87.
25. Churchill W. The Official Report, House of Commons: House of Commons, 1947:206-07.
3. McCrory P, Meeuwisse W, Johnston K et al. Consensus statement on concussion in sport: the 3rd International Conference on Concussion in Sport held in Zurich, November 2008. *Br J Sports Med* 2009;43 Suppl 1:i76-90.
4. McCrory P, Johnston K, Meeuwisse W et al. Summary and agreement statement of the 2nd International Conference on Concussion in Sport, Prague 2004. *Br J Sports Med* 2005;39(4):196-204.
5. Aubry M, Cantu R, Dvorak J et al. Summary and agreement statement of the First International Conference on Concussion in Sport, Vienna 2001. Recommendations for the improvement of safety and health of athletes who may suffer concussive injuries. *Br J Sports Med* 2002;36(1):6-10.
6. McCrory P, Meeuwisse WH, Aubry M et al. Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *Br J Sports Med* 2013;47(5):250-58.
7. Benson BW, Hamilton GM, Meeuwisse WH et al. Is protective equipment useful in preventing concussion? A systematic review of the literature. *Br J Sports Med* 2009;43 Suppl 1: i56-67.
8. Labella CR, Smith BW, Sigurdsson A. Effect of mouthguards on dental injuries and concussions in college basketball. *Med Sci Sports Exerc* 2002;34(1):41-4.
9. Barbic D, Pater J, Brison RJ. Comparison of mouth guard designs and concussion prevention in contact sports: a multicenter randomized controlled trial. *Clin J Sport Med* 2005;15(5):294-8.
10. Benson BW, McIntosh AS, Maddocks D et al. What are the most effective risk-reduction strategies in sport concussion? *Br J Sports Med* 2013;47(5):321-26.
11. Hagel B, Meeuwisse W. Risk compensation: a 'side effect' of sport injury prevention? *Clin J Sport Med* 2004;14(4):193-6.
12. Andersen TE, Arnason A, Engebretsen L et al. Mechanisms of head injuries in elite football. *Br J Sports Med* 2004;38(6):690-6.
13. Shaw NH. Bodychecking in hockey. *CMAJ* 2004;170(1):15-6; author reply 16, 18.
14. Kutcher JS, McCrory P, Davis G et al. What evidence exists for new strategies or technologies in the diagnosis of sports concussion and assessment of recovery? *Br J Sports Med* 2013;47(5):299-303.
15. Davis GA, Iverson GL, Guskiewicz KM et al. Contributions of neuroimaging, balance testing, electrophysiology and blood markers to the assessment of sport-related concussion. *Br J Sports Med* 2009;43 Suppl 1:i36-45.
16. Putukian M, Raftery M, Guskiewicz K et al. Onfield assessment of concussion in the adult athlete. *Br J Sports Med* 2013;47(5):285-88.
17. Makdissi M, Cantu RC, Johnston KM et al. The difficult concussion patient: what is the best approach to investigation and management of persistent (>10 days) postconcussive symptoms? *Br J Sports Med* 2013;47(5):308-13.
18. McCrory P, Meeuwisse WH, Kutcher JS et al. What is the evidence for chronic concussion-related changes in retired

CHAPTER 20

1. Guskiewicz KM, Register-Mihalik J, McCrory P et al. Evidence-based approach to revising the SCAT2: introducing the SCAT3. *Br J Sports Med* 2013;47(5):289-93.
2. Pocket CRT. *Br J Sports Med* 2013;47:267.

- athletes: behavioural, pathological and clinical outcomes? *Br J Sports Med* 2013;47(5):327-30.
19. Livingston SC, Goodkin HP, Hertel JN et al. Differential rates of recovery after acute sport-related concussion: electrophysiologic, symptomatic and neurocognitive indices. *J Clin Neurophysiol* 2012;29(1):23-32.
 20. Schneider KJ, Iverson GL, Emery CA et al. The effects of rest and treatment following sport-related concussion: a systematic review of the literature. *Br J Sports Med* 2013;47(5):304-7.
 21. Silverberg ND, Iverson GL. Is rest after concussion 'the best medicine?': recommendations for activity resumption following concussion in athletes, civilians and military service members. *J Head Trauma Rehabil* 2013;28(4):250-9.
 22. Makdissi M, Davis G, Jordan B et al. Revisiting the modifiers: how should the evaluation and management of acute concussions differ in specific groups? *Br J Sports Med* 2013;47(5):314-20.
 23. Echemendia RJ, Herring S, Bailes J. Who should conduct and interpret the neuropsychological assessment in sports-related concussion? *Br J Sports Med* 2009;43 Suppl 1:132-5.
 24. McCrory PR, Berkovic SF. Second impact syndrome. *Neurology* 1998;50(3):677-83.
 25. McCrory PR, Berkovic SF. Concussive convulsions. Incidence in sport and treatment recommendations. *Sports Med* 1998;25(2):131-6.
 26. McCrory P. Sports concussion and the risk of chronic neurological impairment. *Clin J Sport Med* 2011;21(1):6-12.
 27. Leddy JJ, Kozlowski K, Donnelly JP et al. A preliminary study of subsymptom threshold exercise training for refractory post-concussion syndrome. *Clin J Sport Med* 2010;20(1):21-7.
 28. Gagnon I, Grilli L, Friedman D et al. A pilot study of active rehabilitation for adolescents who are slow to recover from sport-related concussion. *Scand J Med Sci Sports* 2016;26(3):299-306.
 29. Schneider KJ, Meeuwisse WH, Nettel-Aguirre, A et al. Cervicovestibular rehabilitation in sport-related concussion: a randomized controlled trial. *Br J Sports Med* 2014;48(17):1294-8.
 5. Seifert T. Headache in sports. *Curr Pain Headache Rep* 2014;18(9):448.
 6. Williams SJ, Nukada H. Sport and exercise headache: part 1. prevalence among university students. *Br J Sports Med* 1994;28(2):90-5.
 7. Koppen H, van Veldhoven PL. Migraineurs with exercise-triggered attacks have a distinct migraine. *J Headache Pain* 2013;14:99.
 8. Williams, S. J. and H. Nukada. Sport and exercise headache: Part 2. Diagnosis and classification. *Br J Sports Med* 1994;28(2):96-100.
 9. Kernick DP, Goadsby PJ, Royal College of General Practitioners et al. Guidance for the management of headache in sport on behalf of the Royal College of General Practitioners and the British Association for the Study of Headache. *Cephalalgia* 2011;31(1):106-11.
 10. Srikiatkachorn A, le Grand SM, Supornsilpchai W et al. Pathophysiology of medication overuse headache—an update. *Headache* 2014;54(1):204-10.
 11. Sjaastad O, Bakketeig LS. Prevalence of cervicogenic headache: Vaga study of headache epidemiology. *Acta Neurol Scand* 2008;117(3):173-80.
 12. Sjaastad O, Bakketeig LS. Migraine without aura: comparison with cervicogenic headache. Vaga study of headache epidemiology. *Acta Neurol Scand* 2008;117(6):377-83.
 13. Sjaastad O, Fredriksen TA, Pfaffenrath V. Cervicogenic headache: diagnostic criteria. The Cervicogenic Headache International Study Group. *Headache* 1998;38(6):442-5.
 14. Rushton A, Rivett D, Carlesso L et al. International framework for examination of the cervical region for potential of cervical arterial dysfunction prior to orthopaedic manual therapy intervention. *Man Ther* 2014;19(3):222-8.
 15. Darabaneanu S, Overath CH, Rubin D et al. Aerobic exercise as a therapy option for migraine: a pilot study. *Int J Sports Med* 2011;32(6):455-60.
 16. Buse DC, Silberstein SD, Manack AN et al. Psychiatric comorbidities of episodic and chronic migraine. *J Neurol* 2013;260(8):1960-9.
 17. Andrasik F, Lipchik GL, McCrory DC et al. Outcome measurement in behavioral headache research: headache parameters and psychosocial outcomes. *Headache* 2005;45(5):429-37.
 18. Niere K, Robinson P. Determination of manipulative physiotherapy treatment outcome in headache patients. *Man Ther* 1997;2(4):199-205.
 19. Jacobson GP, Ramadan NM, Aggarwal SK et al. The Henry Ford Hospital Headache Disability Inventory (HDI). *Neurology* 1994;44(5):837-42.
 20. Gururaj G, Kulkarni GB, Rao GN et al. Prevalence and sociodemographic correlates of primary headache disorders: results of a population-based survey from Bangalore, India. *Indian J Pub Health* 2014;58(4):241-8.
 21. Buzzi M, Moskowitz M. The pathophysiology of migraine: year 2005. *J Headache Pain* 2005;6:105-11.

CHAPTER 21

1. Martelletti P, Steiner TJ, Bertolote JM et al. The definitive position of headache among the major public health challenges. An end to the slippery slope of disregard. *J Headache Pain* 2007;8(3):149-51.
2. Harpole LH, Samsa GP, Matchar DB et al. Burden of illness and satisfaction with care among patients with headache seen in a primary care setting. *Headache* 2005;45(8):1048-55.
3. Headache Classification Committee of the International Headache Society. The International Classification of Headache Disorders, 3rd edition (beta version). *Cephalalgia* 2013;33(9):629-808.
4. McCrory P, Heywood J, Coffey C. Prevalence of headache in Australian footballers. *Br J Sports Med* 2005;39(2):e10.

22. Goadsby PJ, Charbit AR, Andreou AP et al. Neurobiology of migraine. *Neuroscience* 2009;161(2):327-41.
23. Olesen J. Are headache disorders caused by neurobiological mechanisms? *Curr Opin Neurol* 2006;19(3):277-80.
24. Gantenbein AR, Sandor PS. Physiological parameters as biomarkers of migraine. *Headache* 2006;46(7):1069-74.
25. Goadsby PJ, Zagami AS, Lambert GA. Neural processing of craniovascular pain: a synthesis of the central structures involved in migraine. *Headache* 1991;31(6):365-71.
26. Lippi G, Mattiuzzi C, Cervellin G. Chocolate and migraine: the history of an ambiguous association. *Acta Biomed* 2014;85(3):216-21.
27. Nosedá R, Kainz V, Borsook D et al. Neurochemical pathways that converge on thalamic trigeminovascular neurons: potential substrate for modulation of migraine by sleep, food intake, stress and anxiety. *PLoS One* 2014;9(8):e103929.
28. Ahn AH. Why does increased exercise decrease migraine? *Curr Pain Headache Rep* 2013;17(12):379.
29. Varkey E, Cider A, Carlsson J et al. Exercise as migraine prophylaxis: a randomized study using relaxation and topiramate as controls. *Cephalalgia* 2011;31(14):1428-38.
30. Frich JC, Kristoffersen ES, Lundqvist C. GPs' experiences with brief intervention for medication-overuse headache: a qualitative study in general practice. *Br J Gen Pract* 2014;64(626):e525-31.
31. van der Ende-Kastelijin K, Oerlemans W, Goedegebuure S. An online survey of exercise-related headaches among cyclists. *Headache* 2012;52(10):1566-73.
32. Sjaastad O, Bakketeig LS. Exertional headache. I. Vaga study of headache epidemiology. *Cephalalgia* 2002;22(10):784-90.
33. Doepf F, Valdeuza JM, Schreiber SJ. Incompetence of internal jugular valve in patients with primary exertional headache: a risk factor? *Cephalalgia* 2008;28(2):182-5.
34. Halker RB, Vargas BB. Primary exertional headache: updates in the literature. *Curr Pain Headache Rep* 2013;17:337.
35. Kristoffersen ES, Aaseth K, Grande RB et al. Self-reported efficacy of complementary and alternative medicine: the Akershus study of chronic headache. *J Headache Pain* 2013;14(1):36.
36. Racicki S, Gerwin S, Diclaudio S et al. Conservative physical therapy management for the treatment of cervicogenic headache: a systematic review. *J Man Manip Ther* 2013;21(2):113-24.
37. Bogduk N, Govind J. Cervicogenic headache: an assessment of the evidence on clinical diagnosis, invasive tests, and treatment. *Lancet Neurol* 2009;8(10):959-68.
38. Chua NH, van Suijlekom HA, Vissers KC et al. Differences in sensory processing between chronic cervical zygapophysial joint pain patients with and without cervicogenic headache. *Cephalalgia* 2011;31(8):953-63.
39. Akerman S, Romero-Reyes M. Insights into the pharmacological targeting of the trigeminocervical complex in the context of treatments of migraine. *Expert Rev Neurother* 2013;13(9):1041-59.
40. Niere K. Central nervous system processing in cervicogenic headache. In: Selvaratnam PJ, Niere K, Zuluaga M, eds. *Headache, Orofacial Pain and Bruxism*. Edinburgh: Elsevier, 2009:95-114.
41. Cady R, Schreiber C, Farmer K et al. Primary headaches: a convergence hypothesis. *Headache* 2002;42(3):204-16.
42. Watson DH, Drummond PD. Head pain referral during examination of the neck in migraine and tension-type headache. *Headache* 2012;52(8):1226-35.
43. Bronfort G, Nilsson N, Haas M et al. Non-invasive physical treatments for chronic/recurrent headache. *Cochrane Database Syst Rev* 2004(3):CD001878.
44. Pfaffenrath V, Kaube H. Diagnostics of cervicogenic headache. *Functional Neurology* 1990;5:159-64.
45. Yi X, Cook AJ, Hamill-Ruth RJ et al. Cervicogenic headache in patients with presumed migraine: missed diagnosis or misdiagnosis? *J Pain* 2005;6(10):700-3.
46. Amiri M, Jull G, Bullock-Saxton J et al. Cervical musculoskeletal impairment in frequent intermittent headache. Part 2: subjects with concurrent headache types. *Cephalalgia* 2007;27(8):891-8.
47. Antonaci F, Ghirmal S, Bono G et al. Cervicogenic headache: evaluation of the original diagnostic criteria. *Cephalalgia* 2001;21(5):573-83.
48. Bevilacqua-Grossi D, Pegoretti KS, Goncalves MC et al. Cervical mobility in women with migraine. *Headache* 2009;49(5):726-31.
49. von Piekartz HJ, Schouten S, Aufdemkampe G. Neurodynamic responses in children with migraine or cervicogenic headache versus a control group. A comparative study. *Man Ther* 2007;12(2):153-60.
50. Hall T, Robinson K. The flexion-rotation test and active cervical mobility—a comparative measurement study in cervicogenic headache. *Man Ther* 2004;9(4):197-202.
51. Hall T, Briffa K, Hopper D. The influence of lower cervical joint pain on range of motion and interpretation of the flexion-rotation test. *J Man Manip Ther* 2010;18(3):126-31.
52. Hall TM, Briffa K, Hopper D et al. Comparative analysis and diagnostic accuracy of the cervical flexion-rotation test. *J Headache Pain* 2010;11(5):391-7.
53. Hall TM, Briffa K, Hopper D et al. The relationship between cervicogenic headache and impairment determined by the flexion-rotation test. *J Manip Physiol Ther* 2010;33(9):666-71.
54. Ogincin M, Hall T, Robinson K et al. The diagnostic validity of the cervical flexion-rotation test in C1/2-related cervicogenic headache. *Man Ther* 2007;12(3):256-62.
55. Takasaki H, Hall T, Oshiro S et al. Normal kinematics of the upper cervical spine during the flexion-rotation test—in vivo measurements using magnetic resonance imaging. *Man Ther* 2011;16(2):167-71.
56. Hall T, Briffa K, Hopper D et al. Reliability of manual examination and frequency of symptomatic cervical motion segment dysfunction in cervicogenic headache. *Man Ther* 2010;15(6):542-6.
57. Goncalves DA, Camparis CM, Speciali JG et al. Temporomandibular disorders are differentially associated

- with headache diagnoses: a controlled study. *Clin J Pain* 2011;27(7):611-5.
58. Armijo-Olivo SA, Bravo J, Magee DJ et al. The association between head and cervical posture and temporomandibular disorders: a systematic review. *J Orofacial Pain* 2006;20(1):9-23.
 59. Grondin F, Hall T, Laurentjoye M et al. Upper cervical range of motion is impaired in patients with temporomandibular disorders. *Cranio* 2015; 33(2):91-99.
 60. von Piekartz H, Hall T. Orofacial manual therapy improves cervical movement impairment associated with headache and features of temporomandibular dysfunction: a randomized controlled trial. *Man Ther* 2013;18(4):345-50.
 61. Schiffman E, Ohrbach R, Truelove E et al. Diagnostic Criteria for Temporomandibular Disorders (DC/TMD) for Clinical and Research Applications: recommendations of the International RDC/TMD Consortium Network* and Orofacial Pain Special Interest Groupdagger. *J Oral Facial Pain Headache* 2014;28(1):6-27.
 62. Jull G, Amiri M, Bullock-Saxton J et al. Cervical musculoskeletal impairment in frequent intermittent headache. Part 1: Subjects with single headaches. *Cephalalgia* 2007;27(7):793-802.
 63. Jull G, Barrett C, Magee R et al. Further clinical clarification of the muscle dysfunction in cervical headache. *Cephalalgia* 1999;19(3):179-85.
 64. Watson DH, Trott PH. Cervical headache: an investigation of natural head posture and upper cervical flexor muscle performance. *Cephalalgia* 1993;13(4):272-84; discussion 232.
 65. Zito G, Jull G, Story I. Clinical tests of musculoskeletal dysfunction in the diagnosis of cervicogenic headache. *Man Ther* 2006;11(2):118-29.
 66. Jull G, Falla D, O'Leary S et al. Cervical Spine: Idiopathic Neck Pain. In Jull G, Moore A, Falla D et al. eds. *Grieve's Modern Musculoskeletal Physiotherapy*, 4th edn. Sydney: Elsevier 2015:410-22.
 67. Jull G, Niere K. The cervical spine and headache. In: Boyling G, Jull G, eds. *Grieves Modern Manual Therapy*. 3rd edn. Edinburgh: Churchill Livingstone, 2004:291-309.
 68. Zito G, Jull G, Story I. Clinical tests of musculoskeletal dysfunction in the diagnosis of cervicogenic headache. *Man Ther* 2006;11(2):118-29.
 69. Fernandez-de-las-Penas C, Perez-de-Heredia M, Molero-Sanchez A et al. Performance of the craniocervical flexion test, forward head posture, and headache clinical parameters in patients with chronic tension-type headache: a pilot study. *J Orthop Sports Phys Ther* 2007;37(2):33-9.
 70. Roth JK, Roth RS, Weintraub JR et al. Cervicogenic headache caused by myofascial trigger points in the sternocleidomastoid: a case report. *Cephalalgia* 2007;27(4):375-80.
 71. Allison GT, Nagy BM, Hall T. A randomized clinical trial of manual therapy for cervico-brachial pain syndrome—a pilot study. *Man Ther* 2002;7(2):95-102.
 72. Fleming R, Forsythe S, Cook C. Influential variables associated with outcomes in patients with cervicogenic headache. *J Man Manip Ther* 2007;15(3):155-64.
 73. Jull GA, Stanton WR. Predictors of responsiveness to physiotherapy management of cervicogenic headache. *Cephalalgia* 2005;25(2):101-8.
 74. Kovacs FM, Seco J, Royuela A et al. Patients with neck pain are less likely to improve if they suffer from poor sleep quality. A prospective study in routine practice. *Clinical J Pain* 2015;31(8):713-21.
 75. Haas M, Spegman A, Peterson D et al. Dose response and efficacy of spinal manipulation for chronic cervicogenic headache: a pilot randomized controlled trial. *Spine J* 2010;10(2):117-28.
 76. Izquierdo Perez H, Alonso Perez JL, Gil Martinez A et al. Is one better than another? A randomized clinical trial of manual therapy for patients with chronic neck pain. *Man Ther* 2014;19(3):215-21.
 77. Hing W, Hall T, Rivett D et al. *The Mulligan Concept of Manual Therapy: Textbook of Techniques*. Sydney: Elsevier, 2015.
 78. Hall T, Chan HT, Christensen L et al. Efficacy of a C1-C2 self-sustained natural apophyseal glide (SNAG) in the management of cervicogenic headache. *J Orthop Sports Phys Ther* 2007;37(3):100-7.
 79. Borusiak P, Biedermann H, Bosserhoff S et al. Lack of efficacy of manual therapy in children and adolescents with suspected cervicogenic headache: results of a prospective, randomized, placebo-controlled, and blinded trial. *Headache* 2010;50(2): 224-30.
 80. Hopper D, Bajaj Y, Kei Choi C et al. A pilot study to investigate the short-term effects of specific soft tissue massage on upper cervical movement impairment in patients with cervicogenic headache. *J Man Manip Ther* 2013;21(1):18-23.
 81. Ylinen J, Nikander R, Nykanen M et al. Effect of neck exercises on cervicogenic headache: a randomized controlled trial. *J Rehab Med* 2010;42(4):344-9.
 82. Grondin F, Hall T, Laurentjoye M et al. Upper cervical range of motion is impaired in patients with temporomandibular disorders. *Cranio* 2015;33(2):91-9.
 83. von Piekartz H, Ludtke K. Effect of treatment of temporomandibular disorders (TMD) in patients with cervicogenic headache: a single-blind, randomized controlled study. *Cranio* 2011;29(1):43-56.
 84. Heyer GL, Young JA, Rose SC et al. Post-traumatic headaches correlate with migraine symptoms in youth with concussion. *Cephalalgia* 2016;36(4):309-16.
 85. Kontos AP, Elbin RJ, Lau B et al. Posttraumatic migraine as a predictor of recovery and cognitive impairment after sport-related concussion. *Am J Sports Med* 2013;41(7):1497-504.
 86. Heyer GL, Idris SA. Does analgesic overuse contribute to chronic post-traumatic headaches in adolescent concussion patients? *Pediatr Neurol* 2014;50(5):464-8.
 87. Matthews WB. Footballer's migraine. *Br Med J* 1972;2(5809):326-7.
 88. Mainardi F, Alicicco E, Maggioni F et al. Headache and soccer: a survey in professional soccer players of the Italian 'Series A'. *Neurol Sci* 2009;30(1):33-6.

89. Luks AM, McIntosh SE, Grissom CK et al. Wilderness Medical Society practice guidelines for the prevention and treatment of acute altitude illness: 2014 update. *Wilderness Environ Med* 2014;25(4):S4-14.
90. Cheshire WP, Jr., Ott MC. Headache in divers. *Headache* 2001;41:235-47.

CHAPTER 22

1. Romeo SJ, Hawley CJ, Romeo MW et al. Facial injuries in sports: a team physician's guide to diagnosis and treatment. *Phys Sportsmed* 2005;33(4):45-53.
2. Drolsum L. Eye injuries in sports. *Scand J Med Sci Sports* 1999;9:53-6.
3. Killer HE, Blumer BK, Rust ON. Avulsion of the optic disc after a blow to swimming goggles. *J Pediatr Ophthalmol Strabismus* 1999;36:92-3.
4. Jones NP. Eye injuries in sport: where next? *Br J Sports Med* 1998;32:197-8.
5. Labella CR, Smith BW, Sigurdsson A. Effect of mouthguards on dental injuries and concussions in college basketball. *Med Sci Sports Exerc* 2002;34:41-4.
6. Pashby TJ. Eye injuries in Canadian amateur hockey, still a concern. *Can J Ophthalmol* 1987;22:293-5.
7. Danis RP, Hu K, Bell M. Acceptability of baseball face guards and reduction of oculofacial injury in receptive youth league players. *Inj Prev* 2000;6:232-4.
8. Quarrie KL, Gianotti SM, Chalmers DJ et al. An evaluation of mouthguard requirements and dental injuries in New Zealand rugby union. *Br J Sports Med* 2005;39:650-1.
9. McCrory P, Meeuwisse WH, Aubry M et al. Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *Br J Sports Med* 2013;47:250-8.
10. Andreasen JO, Borum MK, Jacobsen HL et al. Replantation of 400 avulsed permanent incisors 4. Factors related to periodontal ligament healing. *Endod Dent Traumatol* 1995;11:76-89.
11. Blomlöf L. Milk and saliva as possible storage media for traumatically exarticulated teeth prior to replantation. *Swed Dent J* 1981;(suppl 8):1-26.
12. Sigalas E, Regan JD, Kramer PR et al. Survival of human periodontal ligament cells in media proposed for transport of avulsed teeth. *Dent Traumatol* 2004;20:21-8.
13. Pohl Y, Tekin U, Boll M et al. Investigations on a cell culture medium for storage and transportation of avulsed teeth. *Aust Endod J* 1999;25:70-5.
14. Gassner R, Ulmer H, Tuli T et al. Incidence of oral and maxillofacial skiing injuries due to different injury mechanisms. *J Oral Maxillofac Surg* 1999;57:1068-73.
15. Webster DA, Bayliss GV, Spadaro JA. Head and face injuries in scholastic women's lacrosse with and without eyewear. *Med Sci Sports Exerc* 1999;31:938-41.

16. Asplund C, Bettcher S, Borchers J. Facial protection and head injuries in ice hockey: a systematic review. *Br J Sports Med* 2009;43(13):993-9.
17. Vinger PF, Duma SM, Crandall J. Baseball hardness as a risk factor for eye injuries. *Arch Ophthalmol* 1999;117:354-8.
18. Ranalli DN, Demas PN. Orofacial injuries from sport: preventive measures for sports medicine. *Sports Med* 2002;32(7):409-18.

CHAPTER 23

1. Kamitani T, Nimura Y, Nagahiro S et al. Catastrophic head and neck injuries in judo players in Japan from 2003 to 2010. *Am J Sports Med* 2013;41(8):1915-21.
2. Swain MS, Lystad RP, Pollard H et al. Incidence and severity of neck injury in rugby union: a systematic review. *J Sci Med Sport* 2011;14(5):383-9.
3. Asplund C, Webb C, Barkdull T. Neck and back pain in bicycling. *Curr Sports Med Rep* 2005;4(5):271-4.
4. Brown J, Lambert M, Hendricks S et al. Are we currently underestimating the risk of scrum-related neck injuries in rugby union front-row players? *Br J Sports Med* 2014;48(14):1127-9.
5. Nilsson M, Häggglund M, Ekstrand J et al. Head and neck injuries in professional soccer. *Clin J Sport Med* 2013;23(4):255-60.
6. Castinel BH, Adam P, Milburn PD et al. Epidemiology of cervical spine abnormalities in asymptomatic adult professional rugby union players using static and dynamic MRI protocols: 2002 to 2006. *Br J Sports Med* 2010;44(3):194-99.
7. Tsang S, Szeto GP, Lee RY. Normal kinematics of the neck: the interplay between the cervical and thoracic spines. *Man Ther* 2013;18(5):431-7.
8. Lau KT, Cheung KY, Chan KB et al. Relationships between sagittal postures of thoracic and cervical spine, presence of neck pain, neck pain severity and disability. *Man Ther* 2010;15(5):457-62.
9. Panjabi MM, Cholewicki J, Nibu K et al. Critical load of the human cervical spine: an in vitro experimental study. *Clin Biomech (Bristol, Avon)* 1998;13:11-17.
10. Vasavada AN, Li S, Delp SL. Influence of muscle morphometry and moment arms on the moment-generating capacity of human neck muscles. *Spine (Phila Pa 1976)* 1998;23:412-22.
11. Boyd-Clark LC, Briggs CA, Galea MP. Muscle spindle distribution, morphology, and density in longus colli and multifidus muscles of the cervical spine. *Spine (Phila Pa 1976)* 2002;27(7):694-701.
12. Winters JM, Peles JD. *Neck muscle activity and 3-D head kinematics during quasi-static and dynamic tracking movements*. New York: Springer-Verlag, 1990.
13. Behrsin JF, Maguire K. Levator scapulae action during shoulder movement: a possible mechanism of shoulder pain of cervical origin. *Aust J Physiother* 1986;32:101-6.
14. Takasaki H, Hall T, Kaneko S et al. Cervical segmental motion induced by shoulder abduction assessed by magnetic resonance imaging. *Spine (Phila Pa 1976)* 2009;34(3):E122-6.

15. Dwyer A, Aprill C, Bogduk N. Cervical zygapophyseal joint pain patterns. I: a study in normal volunteers. *Spine (Phila Pa 1976)* 1990;15(6):453-7.
16. Slipman CW, Plastaras C, Patel R et al. Provocative cervical discography symptom mapping. *Spine J* 2005;5(4):381-8.
17. Cooper G, Bailey B, Bogduk N. Cervical zygapophysial joint pain maps. *Pain Med* 2007;8(4):344-53.
18. Kulkarni V, Chandy MJ, Babu KS. Quantitative study of muscle spindles in suboccipital muscles of human fetuses. *Neurol India* 2001;49:355-9.
19. Liu JX, Thornell LE, Pedrosa-Domellöf F. Muscle spindles in the deep muscles of the human neck: a morphological and immunocytochemical study. *J Histochem Cytochem* 2003;51: 175-86.
20. Røijezon U, Treleaven J. Management of the sensorimotor system: the cervical region. In: Jull G, Moore A, Falla D et al. (eds). *Grieve's modern musculoskeletal physiotherapy*. 4th ed. UK: Elsevier, 2015.
21. Carroll LJ, Holm LW, Hogg-Johnson S et al. Course and prognostic factors for neck pain in whiplash-associated disorders (WAD). *Eur Spine J* 2008;33(4S):S83-92.
22. Hogg-Johnson S, van der Velde G, Carroll LJ et al. The burden and determinants of neck pain in the general population: results of the bone and joint decade 2000-2010 task force on neck pain and its associated disorders. *Spine (Phila Pa 1976)* 2008;33(4):S39-51.
23. Hoy D, March L, Woolf A et al. The global burden of neck pain: estimates from the global burden of disease 2010 study. *Ann Rheum Dis* 2014;73(7):1309-15.
24. Walton DM, Eilon-Avigdor Y, Wonderham M et al. Exploring the clinical course of neck pain in physical therapy: a longitudinal study. *Arch Phys Med Rehabil* 2014;95(2):303-8.
25. Walton DM, Macdermid JC, Giorgianni AA et al. Risk factors for persistent problems following acute whiplash injury: update of a systematic review and meta-analysis. *J Orthop Sports Phys Ther* 2013;43(2):31-43.
26. Dabette S, Leys D. Cervical-artery dissections: predisposing factors, diagnosis, and outcome. *Lancet Neurol* 2009;8(7):668-78.
27. Thomas LC. Cervical arterial dissection: an overview and implications for manipulative therapy practice. *Man Ther* 2016;21:2-9.
28. Li Q, Shen H, Li M. Magnetic resonance imaging signal changes of alar and transverse ligaments not correlated with whiplash-associated disorders: a meta-analysis of case-control studies. *Eur Spine J* 2013;22(1):14-20.
29. Vetti N, Kråkenes J, Ask T et al. Follow-up MR imaging of the alar and transverse ligaments after whiplash injury: a prospective controlled study. *Am J Neuroradiol* 2011;32(10):1836-41.
30. Chien A, Eliav E, Sterling M. Whiplash (grade II) and cervical radiculopathy share a similar sensory presentation: an investigation using quantitative sensory testing. *Clin J Pain* 2008;24(7):595-603.
31. Chien A, Sterling M. Sensory hypoaesthesia is a feature of chronic whiplash but not chronic idiopathic neck pain. *Man Ther* 2010;15(1):48-53.
32. Antonaci F, Ghirmai S, Bono G et al. Cervicogenic headache: evaluation of the original diagnostic criteria. *Cephalalgia* 2001;21(5):573-83.
33. Ligthart L, Gerrits MM, Boomsma DI et al. Anxiety and depression are associated with migraine and pain in general: an investigation of the interrelationships. *J Pain* 2013;14:363-70.
34. Jensen MP, Karoly P, Braver S. The measurement of clinical pain intensity: a comparison of six methods. *Pain* 1986;27(1):117-26.
35. Vernon H, Mior S. The neck disability index: a study of reliability and validity. *J Manipulative Physiol Ther* 1991;14(7):409-15.
36. Westaway MD, Stratford PW, Binkley JM. The patient-specific functional scale: validation of its use in persons with neck dysfunction. *J Orthop Sports Phys Ther* 1998;27(5):331-8.
37. Jaeschke R, Singer J, Guyatt GH. Measurement of health status. Ascertaining the minimal clinically important difference. *Control Clin Trials* 1989;10(4):407-15.
38. Beaton DE. Understanding the relevance of measured change through studies of responsiveness. *Spine (Phila Pa 1976)* 2000;25(24):3192-9.
39. Cleland JA, Childs JD, Whitman JM. Psychometric properties of the neck disability index and numeric pain rating scale in patients with mechanical neck pain. *Arch Phys Med Rehabil* 2008;89(1):69-74.
40. Stratford PW, Spadoni G. The reliability, consistency, and clinical application of a numeric pain rating scale. *Physiother Can* 2001;53(2):88-91.
41. Cleland JA, Fritz JM, Whitman JM et al. The reliability and construct validity of the neck disability index and patient specific functional scale in patients with cervical radiculopathy. *Spine (Phila Pa 1976)* 2006;31(5):598-602.
42. Duane TM, Wilson SP, Mayglothling J et al. Canadian cervical spine rule compared with computed tomography: a prospective analysis. *J Trauma* 2011;71(2):352-5.
43. Michaleff ZA, Maher CG, Verhagen AP et al. Accuracy of the Canadian C-spine rule and NEXUS to screen for clinically important cervical spine injury in patients following blunt trauma: a systematic review. *CMAJ* 2012;184(16):E867-76.
44. Knackstedt H, Kråkenes J, Bansevicius D et al. Magnetic resonance imaging of craniocervical structures: clinical significance in cervicogenic headaches. *J Headache Pain* 2012; 13(1):39-44.
45. Nakashima H, Yukawa Y, Suda K et al. Abnormal findings on magnetic resonance images of the cervical spines in 1211 asymptomatic subjects. *Spine (Phila Pa 1976)* 2015; 40(6):392-8.
46. Elvey RL. Physical evaluation of the peripheral nervous system in disorders of pain and dysfunction. *J Hand Ther* 1997;10:122-9.
47. Calhoun AH, Ford S, Millen C et al. The prevalence of neck pain in migraine. *Headache* 2010;50(8):1273-7.

48. Bartsch T. Migraine and the neck: new insights from basic data. *Curr Pain Headache Rep* 2005;9:191-6.
49. Jull G, Amiri M, Bullock-Saxton J et al. Cervical musculoskeletal impairment in frequent intermittent headache. Part 1: subjects with single headaches. *Cephalalgia* 2007;27(7):793-802.
50. Snodgrass SJ, Cleland JA, Haskins R et al. The clinical utility of cervical range of motion in diagnosis, prognosis, and evaluating the effects of manipulation: a systematic review. *Physiotherapy* 2014;100(4):290-304.
51. Hall TM, Robinson KW, Fujinawa O et al. Interrater reliability and diagnostic validity of the cervical flexion-rotation test. *J Manipulative Physiol Ther* 2008;31(4):293-300.
52. Takasaki H, Hall T, Oshiro S et al. Normal kinematics of the upper cervical spine during the flexion-rotation test—in vivo measurements using magnetic resonance imaging. *Man Ther* 2011;16(2):167-71.
53. Hall T, Briffa K, Hopper D et al. Long-term stability and minimal detectable change of the cervical flexion-rotation test. *J Orthop Sports Phys Ther* 2010;40(4):225-9.
54. Hall T, Briffa K, Hopper D et al. The relationship between cervicogenic headache and impairment determined by the flexion-rotation test. *J Manipulative Physiol Ther* 2010;33(9):666-71.
55. Hall TM, Briffa K, Hopper D et al. Comparative analysis and diagnostic accuracy of the cervical flexion-rotation test. *J Headache Pain* 2010;11(5):391-7.
56. Schneider GM, Jull G, Thomas K et al. Derivation of a clinical decision guide in the diagnosis of cervical facet joint pain. *Arch Phys Med Rehabil* 2014;95(9):1695-701.
57. Chua NH, Suijlekom HA, Vissers KC et al. Differences in sensory processing between chronic cervical zygapophysial joint pain patients with and without cervicogenic headache. *Cephalalgia* 2011;31(8):953-63.
58. Sterling M, Jull G, Vicenzino B et al. Sensory hypersensitivity occurs soon after whiplash injury and is associated with poor recovery. *Pain* 2003;104:509-17.
59. Maxwell S, Sterling M. An investigation of the use of a numeric pain rating scale with ice application to the neck to determine cold hyperalgesia. *Man Ther* 2013;18(2):172-4.
60. Hall T, Briffa K, Hopper D et al. Reliability of manual examination and frequency of symptomatic cervical motion segment dysfunction in cervicogenic headache. *Man Ther* 2010;15(6):542-6.
61. Moffett JK, Jackson DA, Gardiner ED et al. Randomized trial of two physiotherapy interventions for primary care neck and back pain patients: 'McKenzie' vs brief physiotherapy pain management. *Rheumatol (Oxford)* 2006;45(12):1514-21.
62. Stamos I, Heneghan NR, McCarthy C et al. Inter-examiner reliability of active combined movements assessment of subjects with a history of mechanical neck problems. *Man Ther* 2012;17(5):438-44.
63. Takasaki H, Hall T, Jull G et al. The influence of cervical traction, compression, and Spurling test on cervical intervertebral foramen size. *Spine (Phila Pa 1976)* 2009;34(16):1658-62.
64. Cook C, Brismée JM, Fleming R et al. Identifiers suggestive of clinical cervical spine instability: a Delphi study of physical therapists. *Phys Ther* 2005;85:895-906.
65. Uitvlugt G, Indenbaum S. Clinical assessment of atlantoaxial instability using the Sharp-Purser test. *Arthritis Rheum* 1988;31(7):918-22.
66. Osmotherly PG, Rivett D, Rowe L. Construct validity of clinical tests for alar ligament integrity: an evaluation using magnetic resonance imaging. *Phys Ther* 2012;92(5):718-25.
67. Nee RJ, Jull GA, Vicenzino B et al. The validity of upper-limb neurodynamic tests for detecting peripheral neuropathic pain. *J Orthop Sports Phys Ther* 2012;42(5):413-24.
68. Schmid AB, Brunner F, Luomajoki H et al. Reliability of clinical tests to evaluate nerve function and mechanosensitivity of the upper limb peripheral nervous system. *BMC Musculoskelet Disord* 2009;21(10):11.
69. Wainner RS, Fritz JM, Irrgang JJ et al. Reliability and diagnostic accuracy of the clinical examination and patient self-report measures for cervical radiculopathy. *Spine (Phila Pa 1976)* 2003;28(1):52-62.
70. Cagnie B, Dolphens M, Peeters I et al. Use of muscle functional magnetic resonance imaging to compare cervical flexor activity between patients with whiplash-associated disorders and people who are healthy. *Phys Ther* 2010;90(8):1157-64.
71. Falla DL, Jull GA, Hodges PW. Patients with neck pain demonstrate reduced electromyographic activity of the deep cervical flexor muscles during performance of the craniocervical flexion test. *Spine (Phila Pa 1976)* 2004;29(19):2108-14.
72. Jull G, Kristjansson E, Dall'Alba P. Impairment in the cervical flexors: a comparison of whiplash and insidious onset neck pain patients. *Man Ther* 2004;9(2):89-94.
73. O'Leary S, Cagnie B, Reeve A et al. Is there altered activity of the extensor muscles in chronic mechanical neck pain? A functional magnetic resonance imaging study. *Arch Phys Med Rehabil* 2011;92(6):929-34.
74. Schomacher J, Farina D, Lindstroem R et al. Chronic trauma-induced neck pain impairs the neural control of the deep semispinalis cervicis muscle. *Clin Neurophysiol* 2012;123(7):1403-8.
75. Johnston V, Jull G, Souvlis T et al. Neck movement and muscle activity characteristics in female office workers with neck pain. *Spine (Phila Pa 1976)* 2008;33(5):555-63.
76. Lindstrøm R, Schomacher J, Farina D et al. Association between neck muscle coactivation, pain, and strength in women with neck pain. *Man Ther* 2011;16(1):80-6.
77. Falla D, Bilenkij G, Jull G. Patients with chronic neck pain demonstrate altered patterns of muscle activation during performance of a functional upper limb task. *Spine (Phila Pa 1976)* 2004;29(13):1436-40.
78. Falla D, Jull G, Hodges PW. Feedforward activity of the cervical flexor muscles during voluntary arm movements is delayed in chronic neck pain. *Exper Brain Res* 2004;157(1):43-8.
79. Jull G, Falla D, Vicenzino B et al. The effect of therapeutic exercise on activation of the deep cervical flexor muscles

- in people with chronic neck pain. *Man Ther* 2009;14(6): 696-701.
80. Boudreau SA, Falla D. Chronic neck pain alters muscle activation patterns to sudden movements. *Exp Brain Res* 2014;232(6):2011-20.
 81. Edmondston S, Björnsdóttir G, Pálsson T et al. Endurance and fatigue characteristics of the neck flexor and extensor muscles during isometric tests in patients with postural neck pain. *Man Ther* 2011;16(4):332-8.
 82. Falla D, Rainoldi A, Merletti R et al. Myoelectric manifestations of sternocleidomastoid and anterior scalene muscle fatigue in chronic neck pain patients. *Clin Neurophysiol* 2003;114(3): 488-95.
 83. O'Leary S, Jull G, Kim M et al. Cranio-cervical flexor muscle impairment at maximal, moderate, and low loads is a feature of neck pain. *Man Ther* 2007;12(1):34-9.
 84. Uhlig Y, Weber BR, Grob D et al. Fiber composition and fiber transformations in neck muscles of patients with dysfunction of the cervical spine. *J Orthop Res* 1995;13:240-9.
 85. Elliott JM, Pedler AR, Jull GA et al. Differential changes in muscle composition exist in traumatic and nontraumatic neck pain. *Spine (Phila Pa 1976)* 2014;39(1):39-47.
 86. Kristjansson E. Reliability of ultrasonography for the cervical multifidus muscle in asymptomatic and symptomatic subjects. *Man Ther* 2004;9(2):83-88.
 87. McPartland JM, Brodeur RR, Hallgren RC. Chronic neck pain, standing balance, and suboccipital muscle atrophy—a pilot study. *J Manipulative Physiol Ther* 1997;20(1):24-9.
 88. Elliott J, Pedler A, Kenardy J et al. The temporal development of fatty infiltrates in the neck muscles following whiplash injury: an association with pain and posttraumatic stress. *PLoS One* 2011;6(6):e21194.
 89. De Loose V, Van den Oord M, Keser I et al. MRI study of the morphometry of the cervical musculature in F-16 pilots. *Aviat Space Environ Med* 2009;80(8):727-31.
 90. Elliott J, Sterling M, Noteboom JT et al. Fatty infiltrate in the cervical extensor muscles is not a feature of chronic, insidious-onset neck pain. *Clin Radiol* 2008;63(6):681-7.
 91. Falla D, Farina D. Muscle fiber conduction velocity of the upper trapezius muscle during dynamic contraction of the upper limb in patients with chronic neck pain. *Pain* 2005;116(1-2):138-45.
 92. Nederhand MJ, Hermens HJ, Ijzerman MJ et al. Cervical muscle dysfunction in chronic whiplash-associated disorder grade 2: the relevance of the trauma. *Spine (Phila Pa 1976)* 2002;27(10): 1056-61.
 93. Falla D, Arendt-Nielsen L, Farina D. Gender-specific adaptations of upper trapezius muscle activity to acute nociceptive stimulation. *Pain* 2008;138(1):217-25.
 94. Falla D, Arendt-Nielsen L, Farina D. The pain-induced change in relative activation of upper trapezius muscle regions is independent of the site of noxious stimulation. *Clin Neurophysiol* 2009;120(1):150-7.
 95. Helgadóttir H, Kristjánsson E, Einarsson E et al. Altered activity of the serratus anterior during unilateral arm elevation in patients with cervical disorders. *J Electromyogr Kinesiol* 2011;21:947-53.
 96. Christensen S, Hirata RP, Graven-Nielsen T. The effect of experimental neck pain on pressure pain sensitivity and axio-shoulder motor control. *J Pain* 2015; 16(4):367-79.
 97. Jull GA, O'Leary SP, Falla DL. Clinical assessment of the deep cervical flexor muscles: the craniocervical flexion test. *J Manip Physiol Ther* 2008;31(7):525-33.
 98. Falla D, Jull G, Dall'Alba P et al. An electromyographic analysis of the deep cervical flexor muscles in performance of craniocervical flexion. *Phys Ther* 2003;83(10):899-906.
 99. O'Leary S, Falla D, Jull G et al. Muscle specificity in tests of cervical flexor muscle performance. *J Electromyogr Kinesiol* 2007;17(1):35-40.
 100. Harris KD, Heer DM, Roy TC et al. Reliability of a measurement of neck flexor muscle endurance. *Phys Ther* 2005;85(12):1349-55.
 101. Jull G, Sterling M, Falla D et al. *Whiplash, headache and neck pain: research-based directions for physical therapies*. Edinburgh: Elsevier UK, 2008.
 102. Elliott J, O'Leary S, Cagnie B et al. Muscle functional magnetic resonance imaging of cervical extensor muscles during different cervical extension exercises. *Arch Phys Med Rehabil* 2010;91 1418-22.
 103. Schomacher J, Boudreau SA, Petzke F et al. Localized pressure pain sensitivity is associated with lower activation of the semispinalis cervicis muscle in patients with chronic neck pain. *Clin J Pain* 2013;29(10):898-906.
 104. Lee H, Nicholson LL, Adams RD. Cervical range of motion associations with subclinical neck pain. *Spine (Phila Pa 1976)* 2004;29(1):33-40.
 105. Sahrmann SA. *Diagnosis and treatment of movement impairment syndromes*. St. Louis: Mosby, 2002.
 106. Van Dillen LR, McDonnell MK, Susco TM et al. The immediate effect of passive scapular elevation on symptoms with active neck rotation in patients with neck pain. *Clin J Pain* 2007;23:641-7.
 107. McClure PW, Michener LA, Sennett BJ et al. Direct 3-dimensional measurement of scapular kinematics during dynamic movements in vivo. *J Shoulder Elbow Surg* 2001;10(3):269-77.
 108. Kendall F, McGeary E, Provance P. *Muscle testing and function*. 4th ed. Baltimore: Williams and Wilkins, 1993.
 109. Treleaven J, Takasaki H. Characteristics of visual disturbances reported by subjects with neck pain. *Man Ther* 2014;19(3): 203-7.
 110. Clark NC, Röijezon U, Treleaven J. Proprioception in musculoskeletal rehabilitation. Part 2: clinical assessment and intervention. *Man Ther* 2015;20(3):378-87.
 111. Röijezon U, Clark NC, Treleaven J. Proprioception in musculoskeletal rehabilitation. Part 1: Basic science and

- principles of assessment and clinical interventions. *Man Ther* 2015;20(3):368-77.
112. L'Heureux-Lebeau B, Godbout A, Berbiche D et al. Evaluation of paraclinical tests in the diagnosis of cervicogenic dizziness. *Otol Neurotol* 2014;35(10):1858-65.
 113. Treleaven J, LowChoy N, Darnell R et al. Comparison of sensorimotor disturbance between subjects with persistent whiplash-associated disorder and subjects with vestibular pathology associated with acoustic neuroma. *Arch Phys Med Rehabil* 2008;89(3):522-30.
 114. Della Casa E, Helbling JA, Meichtry A et al. Head-eye movement control tests in patients with chronic neck pain; inter-observer reliability and discriminative validity. *BMC Musculoskel Dis* 2014;15(16).
 115. Treleaven J, Jull G, Lowchoy N. Standing balance in chronic whiplash: comparison between subjects with and without dizziness. *J Rehabil Med* 2005;37(4):219-23.
 116. Treleaven J, Jull G, LowChoy N. The relationship of cervical joint position error to balance and eye movement disturbances in persistent whiplash. *Man Ther* 2006;11(2):99-106.
 117. Yahia A, Ghroubi S, Jribi S et al. Chronic neck pain and vertigo: is a true balance disorder present? *Ann Phys Rehabil Med* 2009;52:556-67.
 118. Roren A, Mayoux-Benhamou MA, Fayad F et al. Comparison of visual and ultrasound based techniques to measure head repositioning in healthy and neck-pain subjects. *Man Ther* 2009;14(3):270-7.
 119. Kristjansson E, Oddsdottir G. 'The Fly': a new clinical assessment and treatment method for deficits of movement control in the cervical spine: reliability and validity. *Spine (Phila Pa 1976)* 2010;35(23):E1298-305.
 120. Oddsdottir G, Kristjansson E, Gislason M. Database of movement control in the cervical spine: reference normal of 182 asymptomatic persons. *Man Ther* 2013;18(3):206-10.
 121. Woodhouse A, Stavdahl Ø, Vasseljen O. Irregular head movement patterns in whiplash patients during a trajectory task. *Exp Brain Res* 2010;201(2):261-70.
 122. Yu LJ, Stokell R, Treleaven J. The effect of neck torsion on postural stability in subjects with persistent whiplash. *Man Ther* 2011;16(4):339-43.
 123. Kristjansson E, Treleaven J. Sensorimotor function and dizziness in neck pain: implications for assessment and management. *J Orthop Sports Phys Ther* 2009;39(5):364-77.
 124. Stokell R, Yu A, Williams K et al. Dynamic and functional balance tasks in subjects with persistent whiplash: a pilot trial. *Man Ther* 2011;16(4):394-8.
 125. Tjell C, Rosenhall U. Smooth pursuit neck torsion test: a specific test for cervical dizziness. *Am J Otol* 1998;19:76-81.
 126. Rushton A, Rivett D, Carlesso L et al. International framework for examination of the cervical region for potential cervical arterial dysfunction prior to orthopaedic manual therapy intervention. *Man Ther* 2014;19:222-8.
 127. Thomas LC, Rivett DA, Bolton PS. Pre-manipulative testing and the use of the velocimeter. *Man Ther* 2008;13(1):29-36.
 128. Zaina C, Grant R, Johnson C et al. The effect of cervical rotation on blood flow in the contralateral vertebral artery. *Man Ther* 2003;8(2):103-9.
 129. Thomas L. Cervical arterial dissection: an overview and implications for manipulative therapy practice. *Man Ther* 2016; 21:2-9.
 130. Gross A, Forget M, St George K et al. Patient education for neck pain. *Cochrane Database Syst Rev* 2012;3:CD005106.
 131. Gross AR, Kaplan F, Huang S et al. Psychological care, patient education, orthotics, ergonomics and prevention strategies for neck pain: an systematic overview update as part of the ICON project. *Open Orthop J* 2013;7:530-61.
 132. Gross A, Kay TM, Paquin JP et al. Exercises for mechanical neck disorders. *Cochrane Database Syst Rev* 2015;1:CD004250.
 133. Gross A, Miller J, D'Sylva J et al. Manipulation or mobilisation for neck pain: a Cochrane review. *Man Ther* 2010;15(4):315-33.
 134. Kroeling P, Gross A, Goldsmith CH et al. Electrotherapy for neck pain. *Cochrane Database Syst Rev* 2013;8:CD004251.
 135. Liu L, Huang QM, Liu QG et al. Effectiveness of dry needling for myofascial trigger points associated with neck and shoulder pain: a systematic review and meta-analysis. *Arch Phys Med Rehabil* 2015;96(5):944-55.
 136. Miller J, Gross A, D'Sylva J et al. Manual therapy and exercise for neck pain: a systematic review. *Man Ther* 2010;15:334-54.
 137. Edmondston S, Sharp M, Symes A et al. Changes in mechanical load and extensor muscle activity in the cervico-thoracic spine induced by sitting posture modification. *Ergonomics* 2011;54(2):179-86.
 138. Grip H, Sundelin G, Gerdle B et al. Cervical helical axis characteristics and its center of rotation during active head and upper arm movements—comparisons of whiplash-associated disorders, non-specific neck pain and asymptomatic individuals. *J Biomech* 2008;41:2799-805.
 139. Eckner JT, Oh YK, Joshi MS et al. Effect of neck muscle strength and anticipatory cervical muscle activation on the kinematic response of the head to impulsive loads. *Am J Sports Med* 2014;42(3):566-76.
 140. King DA, Hume PA, Gianotti S et al. Neck back and spine injuries in amateur rugby league: a review of nine years of Accident Compensation Corporation injury entitlement claims and costs. *J Sci Med Sport* 2011;14(2):126-9.
 141. Mihalik JP, Guskiewicz KM, Marshall SW et al. Does cervical muscle strength in youth ice hockey players affect head impact biomechanics? *Clin J Sport Med* 2011;21(5):416-21.
 142. Jull G, Kenardy J, Hendrikz J et al. Management of acute whiplash: a randomized controlled trial of multidisciplinary stratified treatments. *Pain* 2013;154:1798-806.
 143. Jull G, Sterling M, Kenardy J et al. Does the presence of sensory hypersensitivity influence outcomes of physical rehabilitation for chronic whiplash? A preliminary RCT. *Pain* 2007;129(1-2):28-34.

144. Coronado RA, Gay CW, Bialosky JE et al. Changes in pain sensitivity following spinal manipulation: a systematic review and meta-analysis. *J Electromyogr Kinesiol.* 2012 Oct;22(5):752-67.
145. D'Sylva J, Miller J, Gross A et al. Manual therapy with or without physical medicine modalities for neck pain: a systematic review. *Man Ther* 2010;15(5):415-33.
146. Cagnie B, Dewitte V, Coppieters I et al. Effect of ischemic compression on trigger points in the neck and shoulder muscles in office workers: a cohort study. *J Manipulative Physiol Ther* 2013;36(8):482-9.
147. Furlan AD, Yazdi F, Tsertsvadze A et al. A systematic review and meta-analysis of efficacy, cost-effectiveness, and safety of selected complementary and alternative medicine for neck and low-back pain. *Evid Based Complement Alternat Med* 2012;2012:953139.
148. Llamas-Ramos R, Pecos-Martín D, Gallego-Izquierdo T et al. Comparison of the short-term outcomes between trigger point dry needling and trigger point manual therapy for the management of chronic mechanical neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther* 2014;44(11):852-61.
149. Hing WA, Reid DA, Monaghan M. Manipulation of the cervical spine. *Man Ther* 2003;8(1):2-9.
150. Leaver AM, Maher CG, Herbert RD et al. A randomized controlled trial comparing manipulation with mobilization for recent onset neck pain. *Arch Phys Med Rehabil* 2010;91:1313-18.
151. Hing W, Hall T, Rivett D et al. *The Mulligan concept of manual therapy.* Sydney: Churchill Livingstone, Elsevier, 2015.
152. Kaltenborn FM, Evjenth O, Kaltenborn TB et al. *Manual mobilization of the joints, vol. 2: The spine.* 4th ed. Oslo: Norli, 2003.
153. Maitland GD, Hengeveld E, Banks K et al. *Maitland's vertebral manipulation.* 7th ed. London: Butterworth, 2005.
154. Izquierdo Pérez H, Alonso Perez J, Gil Martinez AG et al. Is one better than another? A randomized clinical trial of manual therapy for patients with chronic neck pain. *Man Ther* 2014;19:215-28.
155. Reid SA, Rivett DA, Katekar MG et al. Comparison of Mulligan sustained natural apophyseal glides and Maitland mobilizations for treatment of cervicogenic dizziness: a randomized controlled trial. *Phys Ther* 2014;94(4):466-76.
156. Tuttle N. Do changes within a manual therapy treatment session predict between-session changes for patients with cervical spine pain? *Aust J Physiother* 2005;51(1):43-8.
157. González-Iglesias J, Fernández-de-las-Peñas C, Cleland J et al. Thoracic spine manipulation for the management of patients with neck pain: a randomized clinical trial. *J Orthop Sports Phys Ther* 2009;39(1):20-7.
158. Lau HM, Wing Chiu TT, Lam TH. The effectiveness of thoracic manipulation on patients with chronic mechanical neck pain—a randomized controlled trial. *Man Ther* 2011;16(2):141-7.
159. Edwards BC. *Manual of combined movements.* 2nd ed. Edinburgh: Churchill Livingstone, 1999.
160. Stamos I, Heneghan N, McCarthy C et al. Inter-examiner reliability of active combined movements assessment of subjects with a history of mechanical neck problems. *Man Ther* 2012;17(5):438-44.
161. Balster SM, Jull GA. Upper trapezius muscle activity during the brachial plexus tension test in asymptomatic subjects. *Man Ther* 1997;2:144-9.
162. Coppieters M, Stappaerts K, Wouters L et al. Aberrant protective force generation during neural provocation testing and the effect of treatment in patients with neurogenic cervicobrachial pain. *J Manipulative Physiol Ther* 2003;26(2):99-106.
163. Coppieters MW, Butler DS. Do 'sliders' slide and 'tensioners' tension? An analysis of neurodynamic techniques and considerations regarding their application. *Man Ther* 2008;13(3):213-21.
164. Allison GT, Nagy BM, Hall T. A randomized clinical trial of manual therapy for cervico-brachial pain syndrome—a pilot study. *Man Ther* 2002;7(2):95-102.
165. Coppieters MW, Stappaerts KH, Wouters LL et al. The immediate effects of a cervical lateral glide treatment technique in patients with neurogenic cervicobrachial pain. *J Orthop Sports Phys Ther* 2003;33(7):369-78.
166. Nee RJ, Vicenzino B, Jull GA et al. Neural tissue management provides immediate clinically relevant benefits without harmful effects for patients with nerve-related neck and arm pain: a randomised trial. *J Physiother* 2012;58(1):23-31.
167. Sterling M, Pedler A, Chan C et al. Cervical lateral glide increases nociceptive flexion reflex threshold but not pressure or thermal pain thresholds in chronic whiplash associated disorders: a pilot randomised controlled trial. *Man Ther* 2010;15(2):149-53.
168. Coppieters MW, Hough AD, Dilley A. Different nerve-gliding exercises induce different magnitudes of median nerve longitudinal excursion: an in vivo study using dynamic ultrasound imaging. *J Orthop Sports Phys Ther* 2009;39(3):164-71.
169. Nee RJ, Vicenzino B, Jull GA et al. A novel protocol to develop a prediction model that identifies patients with nerve-related neck and arm pain who benefit from the early introduction of neural tissue management. *Contemp Clin Trials* 2011;32(5):760-70.
170. Boudreau SA, Farina D, Falla D. The role of motor learning and neuroplasticity in designing rehabilitation approaches for musculoskeletal pain disorders. *Man Ther* 2010;15:410-14.
171. Falla D, Jull G, Hodges P et al. An endurance-strength training regime is effective in reducing myoelectric manifestations of cervical flexor muscle fatigue in females with chronic neck pain. *Clin Neurophysiol* 2006;117(4):828-37.
172. Falla D, Jull G, Russell T et al. Effect of neck exercise on sitting posture in patients with chronic neck pain. *Phys Ther* 2007;87(4):408-17.
173. O'Leary S, Jull G, Kim M et al. Training mode-dependent changes in motor performance in neck pain. *Arch Phys Med Rehabil* 2012;93(7):1225-33.

174. Boudreau S, Romaniello A, Wang K et al. The effects of intra-oral pain on motor cortex neuroplasticity associated with short-term novel tongue-protrusion training in humans. *Pain* 2007;132:169-78.
175. Falla D, O'Leary S, Fagan A et al. Recruitment of the deep cervical flexor muscles during a postural-correction exercise performed in sitting. *Man Ther* 2007;12(2):139-43.
176. Fountain FP, Minear WL, Allison PD. Function of longus colli and longissimus cervicis muscles in man. *Arch Phys Med Rehabil* 1966;47:665-9.
177. Beer A, Treleaven J, Jull G. Can a functional postural exercise improve performance in the cranio-cervical flexion test? A preliminary study. *Man Ther* 2012;17(3):219-24.
178. Schomacher J, Petzke F, Falla D. Localised resistance selectively activates the semispinalis cervicis muscle in patients with neck pain. *Man Ther* 2012;17(6):544-8.
179. Hellebrandt F, Houtz S. Mechanisms of muscle training in man: experimental demonstration of the overload principle. *Phys Ther Rev* 1956;36(6):371-83.
180. Jull G, Falla D, Treleaven J et al. Retraining cervical joint position sense: the effect of two exercise regimes. *J Orthop Res* 2007;25(3):404-12.
181. Revel M, Minguet M, Gregoy P et al. Changes in cervicocephalic kinesthesia after a proprioceptive rehabilitation program in patients with neck pain: a randomized controlled study. *Arch Phys Med Rehabil* 1994;75:895-9.
182. Sarig Bahat H, Takasaki H, Chen X et al. Cervical kinematic training with and without interactive VR training for chronic neck pain—a randomized clinical trial. *Man Ther* 2015;20(1):68-78.
183. Fattori B, Borsari C, Vannucci G et al. Acupuncture treatment for balance disorders following whiplash injury. *Acupunct Electrother Res* 1996;21(3-4):207-17.
184. Kristjansson E, Hardardottir L, Asmundardottir M et al. A new clinical test for cervicocephalic kinesthetic sensibility: 'the fly'. *Arch Phys Med Rehabil* 2004;85(3):490-5.
185. Hurley MV, Newham DJ. The influence of arthrogenous muscle inhibition on quadriceps rehabilitation of patients with early, unilateral osteoarthritic knees. *Br J Rheumatol* 1993;32:127-31.
186. Headache Classification Committee of the International Headache Society (IHS). The international classification of headache disorders, 3rd edition. *Cephalalgia* 2013;33(9):629-808.
187. Bogduk N. The neck and headaches. *Neurol Clin* 2014;32(2):471-7.
188. Amiri M, Jull G, Bullock-Saxton J et al. Cervical musculoskeletal impairment in frequent intermittent headache. Part 2: subjects with concurrent headache types. *Cephalalgia* 2007;27(8):891-98.
189. Mercer S, Bogduk N. Intra-articular inclusions of the cervical synovial joints. *Br J Rheumatol* 1993;32:705-10.
190. Roche CJ, O'Malley M, Dorgan JC et al. A pictorial review of atlanto-axial rotatory fixation: key points for the radiologist. *Clin Radiol* 2001;56(12):947-58.
191. Sobolewski BA, Mittiga MR, Reed JL. Atlantoaxial rotary subluxation after minor trauma. *Pediatr Emerg Care* 2008;24(12):852-6.
192. Engquist M, Löfgren H, Öberg B et al. Surgery versus nonsurgical treatment of cervical radiculopathy: a prospective, randomized study comparing surgery plus physiotherapy with physiotherapy alone with a 2-year follow-up. *Spine* 2013;38(20):1715-22.
193. Altaf F, Mannan K, Bharania P et al. Severe brachial plexus injuries in rugby. *Injury* 2012;43(4):272-3.
194. Safran MR. Nerve injury about the shoulder in athletes, part 2: long thoracic nerve, spinal accessory nerve, burners/stingers, thoracic outlet syndrome. *Am J Sports Med* 2004;32(4):1063-76.
195. Standaert CJ, Herring SA. Expert opinion and controversies in musculoskeletal and sports medicine: stingers. *Arch Phys Med Rehabil* 2009;90(3):402-6.
196. Jull G, Kenardy J, Hendrikz J et al. Management of acute whiplash: a randomized controlled trial of multidisciplinary stratified treatments. *Pain* 2013;154:1798-806.
197. Ritchie C, Hendrikz J, Jull G et al. External validation of a clinical prediction rule to predict full recovery and ongoing moderate/severe disability following acute whiplash injury. *J Orthop Sports Phys Ther* 2015;45(4):242-50.
198. Sterling M, Hendrik J, Kenardy J et al. Assessment and validation of prognostic models for poor functional recovery 12 months following whiplash injury: a multicentre inception cohort study. *Pain* 2012;153(8):1727-34.

CHAPTER 24

1. Pagnani MJ, Warren RF. Stabilizers of the glenohumeral joint. *J Shoulder Elb Surg* 1994;3(3):173-90.
2. Wattanaprakornkul D, Halaki M, Boettcher C et al. A comprehensive analysis of muscle recruitment patterns during shoulder flexion: an electromyographic study. *Clin Anat* 2011;24(5):619-26.
3. Clarsen B, Bahr R, Andersson SH et al. Reduced glenohumeral rotation, external rotation weakness and scapular dyskinesis are risk factors for shoulder injuries among elite male handball players: a prospective cohort study. *Br J Sports Med* 2014;48(17):1327-33.
4. Kawasaki T, Yamakawa J, Kaketa T et al. Does scapular dyskinesis affect top rugby players during a game season? *J Shoulder Elbow Surg* 2012;21(6):709-14.
5. Wassinger CA, Sole G, Osborne H. Clinical measurement of scapular upward rotation in response to acute subacromial pain. *J Orthop Sports Phys Ther* 2013;43(4):199-203.
6. Seitz AL, McClure PW, Finucane S et al. Mechanisms of rotator cuff tendinopathy: intrinsic, extrinsic, or both? *Clin Biomech (Bristol, Avon)* 2011;26(1):1-12.
7. Wilk KE, Obma P, Simpson CD et al. Shoulder injuries in the overhead athlete. *J Orthop Sports Phys Ther* 2009;39(2):38-54.

8. Balke M, Schmidt C, Dedy N et al. Correlation of acromial morphology with impingement syndrome and rotator cuff tears. *Acta Orthop* 2013;84(2):178-83.
9. McClure PW, Michener LA. Staged approach for rehabilitation classification: shoulder disorders (STAR-Shoulder). *Phys Ther* 2015;95(5):791-800.
10. Cools AM, Cambier D, Witvrouw EE. Screening the athlete's shoulder for impingement symptoms: a clinical reasoning algorithm for early detection of shoulder pathology. *Br J Sports Med* 2008;42(8):628-35.
11. Hegedus EJ, Goode AP, Cook CE et al. Which physical examination tests provide clinicians with the most value when examining the shoulder? Update of a systematic review with meta-analysis of individual tests. *Br J Sports Med* 2012;46(14):964-78.
12. Hanchard NC, Lenza M, Handoll HH et al. Physical tests for shoulder impingements and local lesions of bursa, tendon or labrum that may accompany impingement. *Cochrane Database Syst Rev* 2013(4):Cd007427.
13. Hughes PC, Taylor NF, Green RA. Most clinical tests cannot accurately diagnose rotator cuff pathology: a systematic review. *Aust J Physiother* 2008;54(3):159-70.
14. Lewis J. Rotator cuff related shoulder pain: assessment, management and uncertainties. *Man Ther* 2016;23:57-68.
15. Lewis J, McCreesh K, Roy JS et al. Rotator cuff tendinopathy: navigating the diagnosis-management conundrum. *J Orthop Sports Phys Ther* 2015;45(11):923-37.
16. Lewis JS. Rotator cuff tendinopathy/subacromial impingement syndrome: is it time for a new method of assessment? *Br J Sports Med* 2009;43(4):259-64.
17. Frohm A, Heijne A, Kowalski J et al. A nine-test screening battery for athletes: a reliability study. *Scand J Med Sci Sports* 2012;22(3):306-15.
18. Cools AM, De Wilde L, Van Tongel A et al. Measuring shoulder external and internal rotation strength and range of motion: comprehensive intra-rater and inter-rater reliability study of several testing protocols. *J Shoulder Elbow Surg* 2014;23(10):1454-61.
19. Cools AM, Vanderstucken F, Vereecken F et al. Eccentric and isometric shoulder rotator cuff strength testing using a hand-held dynamometer: reference values for overhead athletes. *Knee Surg Sports Traumatol Arthrosc* 2015 Aug 21 [Epub ahead of print].
20. Cools AM, Johansson FR, Cambier DC et al. Descriptive profile of scapulothoracic position, strength and flexibility variables in adolescent elite tennis players. *Br J Sports Med* 2010;44(9):678-84.
21. Borstad JD, Ludewig PM. The effect of long versus short pectoralis minor resting length on scapular kinematics in healthy individuals. *J Orthop Sports Phys Ther* 2005;35(4):227-38.
22. Borms D. Are upper quadrant field tests related to isokinetic upper limb strength in overhead athletes? *J Athl Train* 2016; in press.
23. Kennedy CA, Beaton DE, Solway S et al. *The DASH Outcome Measure User's Manual*. 3rd ed. Toronto, Ontario: Institute for Work & Health, 2001.
24. Beaton DE, Katz JN, Fossel AH et al. Measuring the whole or the parts? Validity, reliability, and responsiveness of the Disabilities of the Arm, Shoulder and Hand outcome measure in different regions of the upper extremity. *J Hand Ther* 2001;14(2):128-46.
25. Raven EE, Haverkamp D, Sierevelt IN et al. Construct validity and reliability of the Disability of Arm, Shoulder and Hand questionnaire for upper extremity complaints in rheumatoid arthritis. *J Rheumatol* 2008;35(12):2334-8.
26. Schmitt JS, Di Fabio RP. Reliable change and minimum important difference (MID) proportions facilitated group responsiveness comparisons using individual threshold criteria. *J Clin Epidemiol* 2004;57(10):1008-18.
27. Slobogean GP, Noonan VK, O'Brien PJ. The reliability and validity of the Disabilities of Arm, Shoulder and Hand, EuroQol-5D, Health Utilities Index, and Short Form-6D outcome instruments in patients with proximal humeral fractures. *J Shoulder Elbow Surg* 2010;19(3):342-8.
28. Angst F, Pap G, Mannion AF et al. Comprehensive assessment of clinical outcome and quality of life after total shoulder arthroplasty: usefulness and validity of subjective outcome measures. *Arthritis Rheum* 2004;51(5):819-28.
29. Staples MP, Forbes A, Green S et al. Shoulder-specific disability measures showed acceptable construct validity and responsiveness. *J Clin Epidemiol* 2010;63(2):163-70.
30. Roy JS, MacDermid JC, Woodhouse LJ. Measuring shoulder function: a systematic review of four questionnaires. *Arthritis Rheum* 2009;61(5):623-32.
31. Beaton DE, Wright JG, Katz JN. Development of the QuickDASH: comparison of three item-reduction approaches. *J Bone Joint Surg Am* 2005;87(5):1038-46.
32. Gummesson C, Ward MM, Atroshi I. The shortened Disabilities of the Arm, Shoulder and Hand questionnaire (QuickDASH): validity and reliability based on responses within the full-length DASH. *BMC Musculoskelet Disord* 2006;7:44.
33. Mintken PE, Glynn P, Cleland JA. Psychometric properties of the shortened Disabilities of the Arm, Shoulder, and Hand questionnaire (QuickDASH) and Numeric Pain Rating Scale in patients with shoulder pain. *J Shoulder Elbow Surg* 2009;18(6):920-6.
34. Angst F, Goldhahn J, Drerup S et al. How sharp is the short QuickDASH? A refined content and validity analysis of the short form of the Disabilities of the Shoulder, Arm and Hand questionnaire in the strata of symptoms and function and specific joint conditions. *Qual Life Res* 2009;18(8):1043-51.
35. Beaton DE, Richards RR. Measuring function of the shoulder. A cross-sectional comparison of five questionnaires. *J Bone Joint Surg Am* 1996;78(6):882-90.
36. Beaton D, Richards RR. Assessing the reliability and responsiveness of 5 shoulder questionnaires. *J Shoulder Elbow Surg* 1998;7(6):565-72.
37. Godfrey J, Hamman R, Lowenstein S et al. Reliability, validity, and responsiveness of the Simple Shoulder Test: psychometric

- properties by age and injury type. *J Shoulder Elbow Surg* 2007;16(3):260-7.
38. Roddey TS, Olson SL, Cook KF et al. Comparison of the University of California-Los Angeles Shoulder Scale and the Simple Shoulder Test with the Shoulder Pain and Disability Index: single-administration reliability and validity. *Phys Ther* 2000;80(8):759-68.
 39. MacDermid JC, Drosdowech D, Faber K. Responsiveness of self-report scales in patients recovering from rotator cuff surgery. *J Shoulder Elbow Surg* 2006;15(4):407-14.
 40. Angst F, Schwyzer HK, Aeschlimann A et al. Measures of adult shoulder function: Disabilities of the Arm, Shoulder, and Hand questionnaire (DASH) and its short version (QuickDASH), Shoulder Pain and Disability Index (SPADI), American Shoulder and Elbow Surgeons (ASES) Society standardized shoulder assessment form, Constant (Murley) Score (CS), Simple Shoulder Test (SST), Oxford Shoulder Score (OSS), Shoulder Disability Questionnaire (SDQ), and Western Ontario Shoulder Instability index (WOSI). *Arthritis Care Res (Hoboken)* 2011;63 Suppl 11:S174-88.
 41. Angst F, Goldhahn J, Pap G et al. Cross-cultural adaptation, reliability and validity of the German Shoulder Pain and Disability Index (SPADI). *Rheumatology* 2007;46(1):87-92.
 42. Tveita EK, Ekeberg OM, Juel NG et al. Responsiveness of the Shoulder Pain and Disability Index in patients with adhesive capsulitis. *BMC Musculoskelet Disord* 2008;9:161.
 43. Ekeberg OM, Bautz-Holter E, Keller A et al. A questionnaire found disease-specific WORC index is not more responsive than SPADI and OSS in rotator cuff disease. *J Clin Epidemiol* 2010;63(5):575-84.
 44. Ekeberg OM, Bautz-Holter E, Tveita EK et al. Agreement, reliability and validity in 3 shoulder questionnaires in patients with Rotator Cuff disease. *BMC Musculoskelet Disord* 2008;9:68.
 45. de Witte PB, Henseler JF, Nagels J et al. The Western Ontario rotator cuff index in rotator cuff disease patients: a comprehensive reliability and responsiveness validation study. *Am J Sports Med* 2012;40(7):1611-9.
 46. Kirkley A, Alvarez C, Griffin S. The development and evaluation of a disease-specific quality-of-life questionnaire for disorders of the rotator cuff: the Western Ontario Rotator Cuff index. *Clin J Sport Med* 2003;13(2):84-92.
 47. Kirkley A, Griffin S, Dainty K. Scoring systems for the functional assessment of the shoulder. *Arthroscopy* 2003;19(10):1109-20.
 48. Kirkley A, Griffin S, McLintock H et al. The development and evaluation of a disease-specific quality of life measurement tool for shoulder instability. The Western Ontario Shoulder Instability index (WOSI). *Am J Sports Med* 1998;26(6):764-72.
 49. Drerup S, Angst F, Griffin S et al. [Western Ontario Shoulder Instability index (WOSI): translation and cross-cultural adaptation for use by German speakers]. *Orthopade* 2010;39(7):711-8.
 50. Hofstaetter JG, Hanslik-Schnabel B, Hofstaetter SG et al. Cross-cultural adaptation and validation of the German version of the Western Ontario Shoulder Instability index. *Arch Orthop Trauma Surg* 2010;130(6):787-96.
 51. Salomonsson B, Ahlstrom S, Dalen N et al. The Western Ontario Shoulder Instability index (WOSI): validity, reliability, and responsiveness retested with a Swedish translation. *Acta Orthop* 2009;80(2):233-8.
 52. van der Linde JA, Willems WJ, van Kampen DA et al. Measurement properties of the Western Ontario Shoulder Instability index in Dutch patients with shoulder instability. *BMC Musculoskelet Disord* 2014;15:211.
 53. Alberta FG, ElAttrache NS, Bissell S et al. The development and validation of a functional assessment tool for the upper extremity in the overhead athlete. *Am J Sports Med* 2010;38(5):903-11.
 54. Wilk KE, Hooks TR. Rehabilitation of the throwing athlete: where we are in 2014. *Clin Sports Med* 2015;34(2):247-61.
 55. Coombes BK, Bisset L, Brooks P et al. Effect of corticosteroid injection, physiotherapy, or both on clinical outcomes in patients with unilateral lateral epicondylalgia: a randomized controlled trial. *JAMA* 2013;309(5):461-9.
 56. Littlewood C, Ashton J, Chance-Larsen K et al. Exercise for rotator cuff tendinopathy: a systematic review. *Physiotherapy* 2012;98(2):101-9.
 57. Reinold MM, Escamilla RF, Wilk KE. Current concepts in the scientific and clinical rationale behind exercises for glenohumeral and scapulohoracic musculature. *J Orthop Sports Phys Ther* 2009;39(2):105-17.
 58. Boettcher CE, Ginn KA, Cathers I. Which is the optimal exercise to strengthen supraspinatus? *Med Sci Sports Exerc* 2009;41(11):1979-83.
 59. Maenhout AG, Mahieu NN, De Muynck M et al. Does adding heavy load eccentric training to rehabilitation of patients with unilateral subacromial impingement result in better outcome? A randomized, clinical trial. *Knee Surg Sports Traumatol Arthrosc* 2013;21(5):1158-67.
 60. Holmgren T, Bjornsson Hallgren H, Oberg B et al. Effect of specific exercise strategy on need for surgery in patients with subacromial impingement syndrome: randomised controlled study. *BMJ* 2012;344:e787.
 61. Speed C. A systematic review of shockwave therapies in soft tissue conditions: focusing on the evidence. *Br J Sports Med* 2014;48(21):1538-42.
 62. Namdari S, Baldwin K, Ahn A et al. Performance after rotator cuff tear and operative treatment: a case-control study of major league baseball pitchers. *J Athl Train* 2011;46(3):296-302.
 63. Kuhn JE, Dunn WR, Sanders R et al. Effectiveness of physical therapy in treating atraumatic full-thickness rotator cuff tears: a multicenter prospective cohort study. *J Shoulder Elbow Surg* 2013;22(10):1371-9.
 64. Ainsworth R, Lewis JS. Exercise therapy for the conservative management of full thickness tears of the rotator cuff: a systematic review. *Br J Sports Med* 2007;41(4):200-10.
 65. Uhl TL, Muir TA, Lawson L. Electromyographical assessment of passive, active assistive, and active shoulder rehabilitation exercises. *PM&R* 2010;2(2):132-41.

66. Levy O, Mullett H, Roberts S et al. The role of anterior deltoid re-education in patients with massive irreparable degenerative rotator cuff tears. *J Shoulder Elbow Surg* 2008;17(6):863-70.
67. Smits-Engelsman B, Klerks M, Kirby A. Beighton score: a valid measure for generalized hypermobility in children. *J Pediatr* 2011;158(1):119-23, 123.e1-4.
68. Gagey OJ, Gagey N. The hyperabduction test. *J Bone Joint Surg Br* 2001;83(1):69-74.
69. Kibler WB, Kuhn JE, Wilk K et al. The disabled throwing shoulder: spectrum of pathology-10-year update. *Arthroscopy* 2013;29(1):141-61.e26.
70. Braun S, Kokmeyer D, Millett PJ. Shoulder injuries in the throwing athlete. *J Bone Joint Surg Am* 2009;91(4):966-78.
71. Edwards SL, Lee JA, Bell JE et al. Nonoperative treatment of superior labrum anterior posterior tears: improvements in pain, function, and quality of life. *Am J Sports Med* 2010;38(7):1456-61.
72. Fedorow WW, Ramkumar P, McCulloch PC et al. Return to play after treatment of superior labral tears in professional baseball players. *Am J Sports Med* 2014;42(5):1155-60.
73. Gorantla K, Gill C, Wright RW. The outcome of type II SLAP repair: a systematic review. *Arthroscopy* 2010;26(4):537-45.
74. Wilk KE, Reinold MM, Andrews JR. *The Athlete's Shoulder*. 2nd ed. Philadelphia: Churchill, Livingston, 2009.
75. Cools AM, Borms D, Cottens S et al. Rehabilitation exercises for athletes with biceps disorders and SLAP lesions: a continuum of exercises with increasing loads on the biceps. *Am J Sports Med* 2014;42(6):1315-22.
76. Borsa PA, Laudner KG, Sauers EL. Mobility and stability adaptations in the shoulder of the overhead athlete: a theoretical and evidence-based perspective. *Sports Med* 2008;38(1):17-36.
77. Cools AM, Declercq G, Cagnie B et al. Internal impingement in the tennis player: rehabilitation guidelines. *Br J Sports Med* 2008;42(3):165-71.
78. Grossman MG, Tibone JE, McGarry MH et al. A cadaveric model of the throwing shoulder: a possible etiology of superior labrum anterior-to-posterior lesions. *J Bone Joint Surg Am* 2005;87(4):824-31.
79. Maenhout AG, Van Eessel V, Van Dyck L et al. Quantifying acromioclavicular distance in overhead athletes with glenohumeral internal rotation loss and the influence of a stretching program. *Am J Sports Med* 2012;40(9):2105-12.
80. Wilk KE, Macrina LC, Fleisig GS et al. Correlation of glenohumeral internal rotation deficit and total rotational motion to shoulder injuries in professional baseball pitchers. *Am J Sports Med* 2011;39(2):329-35.
81. Shanley E, Rauh MJ, Michener LA et al. Shoulder range of motion measures as risk factors for shoulder and elbow injuries in high school softball and baseball players. *Am J Sports Med* 2011;39(9):1997-2006.
82. McClure PW, Balaicuis J, Heiland D et al. A randomized controlled comparison of stretching procedures for posterior shoulder tightness. *J Orthop Sports Phys Ther* 2007;37(3):108-14.
83. Manske RC, Meschke M, Porter A et al. A randomized controlled single-blinded comparison of stretching versus stretching and joint mobilization for posterior shoulder tightness measured by internal rotation motion loss. *Sports Health* 2010;2(2):94-100.
84. Cools AM, Johansson FR, Cagnie B et al. Stretching the posterior shoulder structures in subjects with internal rotation deficit: comparison of two stretching techniques. *Shoulder & Elbow* 2012;4(1):56-63.
85. Moore SD, Laudner KG, McLoda TA et al. The immediate effects of muscle energy technique on posterior shoulder tightness: a randomized controlled trial. *J Orthop Sports Phys Ther* 2011;41(6):400-7.
86. Tyler TF, Nicholas SJ, Lee SJ et al. Correction of posterior shoulder tightness is associated with symptom resolution in patients with internal impingement. *Am J Sports Med* 2010;38(1):114-9.
87. McConnell J, McIntosh B. The effect of tape on glenohumeral rotation range of motion in elite junior tennis players. *Clin J Sport Med* 2009;19(2):90-4.
88. Cools AM, Struyf F, De Mey K et al. Rehabilitation of scapular dyskinesis: from the office worker to the elite overhead athlete. *Br J Sports Med* 2014;48(8):692-7.
89. Borstad JD, Ludewig PM. Comparison of three stretches for the pectoralis minor muscle. *J Shoulder Elbow Surg* 2006;15(3):324-30.
90. Mottram SL, Woledge RC, Morrissey D. Motion analysis study of a scapular orientation exercise and subjects' ability to learn the exercise. *Man Ther* 2009;14(1):13-8.
91. Cools AM, Dewitte V, Lanszweert F et al. Rehabilitation of scapular muscle balance: which exercises to prescribe? *Am J Sports Med* 2007;35(10):1744-51.
92. De Mey K, Danneels L, Cagnie B et al. Scapular muscle rehabilitation exercises in overhead athletes with impingement symptoms: effect of a 6-week training program on muscle recruitment and functional outcome. *Am J Sports Med* 2012;40(8):1906-15.
93. Merolla G, De Santis E, Sperling JW et al. Infraspinatus strength assessment before and after scapular muscles rehabilitation in professional volleyball players with scapular dyskinesis. *J Shoulder Elbow Surg* 2010;19(8):1256-64.
94. Robertson GA, Wood AM. Return to sport following clavicle fractures: a systematic review. *Br Med Bull* 2016;119(1):111-28.
95. Beitzel K, Mazzocca AD, Bak K et al. ISAKOS upper extremity committee consensus statement on the need for diversification of the Rockwood classification for acromioclavicular joint injuries. *Arthroscopy* 2014;30(2):271-8.
96. Canadian Orthopaedic Trauma Society. Multicenter Randomized Clinical Trial of Nonoperative Versus Operative Treatment of Acute Acromio-Clavicular Joint Dislocation. *J Orthop Trauma* 2015;29(11):479-87.
97. Woodmass JM, Esposito JG, Ono Y et al. Complications following arthroscopic fixation of acromioclavicular separations: a systematic review of the literature. *Open Access J Sports Med* 2015;6:97-107.

98. Pensak M, Grumet RC, Slabaugh MA et al. Open versus arthroscopic distal clavicle resection. *Arthroscopy* 2010;26(5):697-704.
99. ElMaraghy AW, Devereaux MW. A systematic review and comprehensive classification of pectoralis major tears. *J Shoulder Elbow Surg* 2012;21(3):412-22.
100. Balazs GC, Brelin AM, Donohue MA et al. Incidence Rate and Results of the Surgical Treatment of Pectoralis Major Tendon Ruptures in Active-Duty Military Personnel. *Am J Sports Med* 2016;44(7):1837-43.
101. Wong CK, Levine WN, Deo K et al. Natural history of frozen shoulder: fact or fiction? A systematic review. *Physiotherapy* 2016.
102. Xiao RC, Walley KC, DeAngelis JP et al. Corticosteroid Injections for Adhesive Capsulitis: A Review. *Clin J Sport Med* 2016.
103. Grant JA, Schroeder N, Miller BS et al. Comparison of manipulation and arthroscopic capsular release for adhesive capsulitis: a systematic review. *J Shoulder Elbow Surg* 2013;22(8):1135-45.
104. Eljabu W, Klinger HM, von Knoch M. Prognostic factors and therapeutic options for treatment of frozen shoulder: a systematic review. *Arch Orthop Trauma Surg* 2016;136(1):1-7.
105. Hill LJ, Jelsing EJ, Terry MJ et al. Evaluation, treatment, and outcomes of suprascapular neuropathy: a 5-year review. *PMR* 2014;6(9):774-80.
106. Ferretti A, De Carli A, Fontana M. Injury of the suprascapular nerve at the spinoglenoid notch. The natural history of infraspinatus atrophy in volleyball players. *Am J Sports Med* 1998;26(6):759-63.
107. Reeser JC, Fleisig GS, Cools AM et al. Biomechanical insights into the aetiology of infraspinatus syndrome. *Br J Sports Med* 2013;47(4):239-44.
108. Hughes PC, Green RA, Taylor NF. Isolation of infraspinatus in clinical test positions. *J Sci Med Sport* 2014;17(3):256-60.
109. Lee S, Saetia K, Saha S et al. Axillary nerve injury associated with sports. *Neurosurg Focus* 2011;31(5):E10.
110. Galvin JW, Eichinger JK. Outcomes Following Closed Axillary Nerve Injury: A Case Report and Review of the Literature. *Mil Med* 2016;181(3):e291-7.
111. Povlsen B, Hansson T, Povlsen SD. Treatment for thoracic outlet syndrome. *Cochrane Database Syst Rev* 2014(11):Cd007218.
112. Peek J, Vos CG, Unlu C et al. Outcome of surgical treatment for thoracic outlet syndrome: systematic review and meta-analysis. *Ann Vasc Surg* 2016.
113. Klitfod L, Broholm R, Baekgaard N. Deep venous thrombosis of the upper extremity. A review. *Int Angiol* 2013;32(5):447-52.
114. Slobogean GP, Johal H, Lefavre KA et al. A scoping review of the proximal humerus fracture literature. *BMC Musculoskeletal Disord* 2015;16:112.
115. Warth RJ, Spiegl UJ, Millett PJ. Scapulothoracic bursitis and snapping scapula syndrome: a critical review of current evidence. *Am J Sports Med* 2015;43(1):236-45.
116. Gaskill T, Millett PJ. Snapping scapula syndrome: diagnosis and management. *J Am Acad Orthop Surg* 2013;21(4):214-24.
117. De Mey K, Danneels L, Cagnie B et al. Kinetic chain influences on upper and lower trapezius muscle activation during eight variations of a scapular retraction exercise in overhead athletes. *J Sci Med Sport* 2013;16(1):65-70.
118. Maenhout A, Van Praet K, Pizzi L et al. Electromyographic analysis of knee push up plus variations: what is the influence of the kinetic chain on scapular muscle activity? *Br J Sports Med* 2010;44(14):1010-5.
119. Creighton DW, Shrier I, Shultz R et al. Return-to-play in sport: a decision-based model. *Clin J Sport Med* 2010;20(5):379-85.
120. McCarty EC, Ritchie P, Gill HS et al. Shoulder instability: return to play. *Clin Sports Med* 2004;23(3):335-51, vii-viii.
121. Wilk KE, Yenchak AJ, Arrigo CA et al. The Advanced Throwers Ten Exercise Program: a new exercise series for enhanced dynamic shoulder control in the overhead throwing athlete. *Phys Sportsmed* 2011;39(4):90-7.

CHAPTER 25

1. Heales LJ, Broadhurst N, Mellor R et al. Diagnostic ultrasound imaging for lateral epicondylalgia: a case-control study. *Med Sci Sports Exerc* 2014;46(11):2070-6.
2. Clarke AW, Ahmad M, Curtis M et al. Lateral elbow tendinopathy: correlation of ultrasound findings with pain and functional disability. *Am J Sports Med* 2010;38(6):1209-14.
3. Qi L, Zhang YD, Yu RB et al. Magnetic Resonance Imaging of Patients With Chronic Lateral Epicondylitis: Is There a Relationship Between Magnetic Resonance Imaging Abnormalities of the Common Extensor Tendon and the Patient's Clinical Symptom? *Medicine (Baltimore)* 2016;95(5):e2681.
4. Fairbank SM, Corlett RJ. The role of the extensor digitorum communis muscle in lateral epicondylitis. *J Hand Surg Br* 2002;27B(5):405-9.
5. Milz S, Tischer T, Buettner A et al. Molecular composition and pathology of entheses on the medial and lateral epicondyles of the humerus: a structural basis for epicondylitis. *Ann Rheum Dis* 2004;63(9):1015-21.
6. Coonrad RW, Hooper WR. Tennis elbow: its course, natural history, conservative and surgical management. *J Bone Joint Surg Am* 1973;55(6):1177-82.
7. Nirschl RP, Pettrone FA. Tennis elbow. The surgical treatment of lateral epicondylitis. *J Bone Joint Surg Am* 1979;61(6):832-9.
8. Kraushaar BS, Nirschl RP. Tendinosis of the elbow (tennis elbow). Clinical features and findings of histological, immunohistochemical, and electron microscopy studies. *J Bone Joint Surg Am* 1999;81A(2):259-78.
9. Alfredson H, Ohberg L, Forsgren S. Is vasculo-neural ingrowth the cause of pain in chronic Achilles tendinosis? An investigation using ultrasonography and colour Doppler, immunohistochemistry, and diagnostic injections. *Knee Surg Sports Traumatol Arthrosc* 2003;11(5):334-38.

10. Regan W, Wold LE, Coonrad R et al. Microscopic histopathology of chronic refractory lateral epicondylitis. *Am J Sports Med* 1992;20(6):746-9.
11. Heales LJ, Vicenzino B, MacDonald DA et al. Forearm muscle activity is modified bilaterally in unilateral lateral epicondylalgia: a case-control study. *Scand J Med Sci Sports* 2015 [Dec 1. doi: 10.1111/sms.12584 epub ahead of print].
12. Bisset LM, Russell T, Bradley S et al. Bilateral sensorimotor abnormalities in unilateral lateral epicondylalgia. *Arch Phys Med Rehabil* 2006;87(4):490-5.
13. Alizadehkhayat O, Fisher AC, Kemp GJ et al. Upper limb muscle imbalance in tennis elbow: a functional and electromyographic assessment. *J Orthop Res* 2007;25(12):1651-7.
14. Andersson G, Forsgren S, Scott A et al. Tenocyte hypercellularity and vascular proliferation in a rabbit model of tendinopathy: contralateral effects suggest the involvement of central neuronal mechanisms. *Br J Sports Med* 2011;45(5):399-406.
15. Pienimäki TT, Kauranen K, Vanharanta H. Bilaterally decreased motor performance of arms in patients with chronic tennis elbow. *Arch Phys Med Rehabil* 1997;78(10):1092-5.
16. Waugh EJ, Jaglal SB, Davis AM. Computer use associated with poor long-term prognosis of conservatively managed lateral epicondylalgia. *J Orthop Sports Phys Ther* 2004;34(12):770-80.
17. Coombes BK, Bisset L, Vicenzino B. Bilateral cervical dysfunction in patients with unilateral lateral epicondylalgia without concomitant cervical or upper limb symptoms: a cross-sectional case-control study. *J Manipulative Physiol Ther* 2014;37(2):79-86.
18. Coombes BK, Bisset L, Vicenzino B. A new integrative model of lateral epicondylalgia. *Br J Sports Med* 2009;43(4):252-8.
19. Cleland JA, Flynn TW, Palmer JA. Incorporation of manual therapy directed at the cervicotoracic spine in patients with lateral epicondylalgia: a pilot clinical trial. *J Man Manip Ther* 2005;13(3):143-51.
20. Smidt N, Lewis M, van der Windt D et al. Lateral epicondylitis in general practice: course and prognostic indicators of outcome. *J Rheumatol* 2006;33(10):2053-9.
21. Pienimäki T, Karinen P, Kemilä T et al. Long-term follow-up of conservatively treated chronic tennis elbow patients. A prospective and retrospective analysis. *Scand J Rehabil Med* 1998;30(3):159-66.
22. Coombes BK, Bisset L, Connelly LB et al. Optimising corticosteroid injection for lateral epicondylalgia with the addition of physiotherapy: a protocol for a randomised control trial with placebo comparison. *BMC Musculoskel Disord* 2009;10(1):76.
23. Pienimäki TT, Tarvainen TK, Siira PT et al. Progressive strengthening and stretching exercises and ultrasound for chronic lateral epicondylitis. *Physiotherapy* 1996;82(9):522-30.
24. Svernlöv B, Adolfsson L. Non-operative treatment regime including eccentric training for lateral humeral epicondylalgia. *Scand J Med Sci Sports* 2001;11(6):328-34.
25. Bjordal JM, Lopes-Martins RA, Joensen J et al. A systematic review with procedural assessments and meta-analysis of low level laser therapy in lateral elbow tendinopathy (tennis elbow). *BMC Musculoskel Disord* 2008;9:75.
26. Staples MP, Forbes A, Ptasznik R et al. A randomized controlled trial of extracorporeal shock wave therapy for lateral epicondylitis (tennis elbow). *J Rheumatol* 2008;35(10):2038-46.
27. Chesterton LS, van der Windt DA, Sim J et al. Transcutaneous electrical nerve stimulation for the management of tennis elbow: a pragmatic randomized controlled trial: the TATE trial (ISRCTN 87141084). *BMC Musculoskel Disord* 2009;10:156.
28. Bjordal JM, Johnson MI, Ljunggreen AE. Transcutaneous electrical nerve stimulation (TENS) can reduce postoperative analgesic consumption. A meta-analysis with assessment of optimal treatment parameters for postoperative pain. *Eur J Pain* 2003;7(2):181-8.
29. van der Windt DA, van der Heijden GJ, van den Berg SG et al. Ultrasound therapy for musculoskeletal disorders: a systematic review. *Pain* 1999;81(3):257-71.
30. Oken O, Kahraman Y, Ayhan F et al. The short-term efficacy of laser, brace, and ultrasound treatment in lateral epicondylitis: a prospective, randomized, controlled trial. *J Hand Ther* 2008;21(1):63-8.
31. Vicenzino B, Hing W, Rivett D et al. *Mobilisation With Movement: The Art and the Science*. Sydney: Churchill Livingstone, 2011.
32. Bisset L, Beller E, Jull G et al. Mobilisation with movement and exercise, corticosteroid injection, or wait and see for tennis elbow: randomised trial. *BMJ* 2006;333(7575):939-41.
33. Vicenzino B, Cleland JA, Bisset L. Joint manipulation in the management of lateral epicondylalgia: a clinical commentary. *J Man Manip Ther* 2007;15(1):50-6.
34. Stasinopoulos D, Stasinopoulos I. Comparison of effects of Cyriax physiotherapy, a supervised exercise programme and polarized polychromatic non-coherent light (Biopton light) for the treatment of lateral epicondylitis. *Clin Rehabil* 2006;20(1):12-23.
35. Verhaar JA, Walenkamp GH, van Mameren H et al. Local corticosteroid injection versus Cyriax-type physiotherapy for tennis elbow. *J Bone Joint Surg Br* 1996;78B(1):128-32.
36. Fernández-Carnero J, Fernández-de-las-Peñas C, de la Llave-Rincon AI et al. Bilateral myofascial trigger points in the forearm muscles in patients with chronic unilateral lateral epicondylalgia: a blinded, controlled study. *Clin J Pain* 2008;24(9):802-7.
37. Bisset L, Paungmali A, Vicenzino B et al. A systematic review and meta-analysis of clinical trials on physical interventions for lateral epicondylalgia. *Br J Sports Med* 2005;39(7):411-22.
38. Ng G. The effects of forearm brace tension on neuromuscular performance in subjects with lateral humeral epicondylitis: a review. *Int J Sports Med* 2005;6(2):124-9.
39. Vicenzino B, Brooksbank J, Minto J et al. Initial effects of elbow taping on pain-free grip strength and pressure pain threshold. *J Orthop Sports Phys Ther* 2003;33(7):400-7.
40. Coombes BK, Bisset L, Vicenzino B. Efficacy and safety of corticosteroid injections and other injections for management

- of tendinopathy: a systematic review of randomised controlled trials. *Lancet* 2010;376(9754):1751-67.
41. Coombes BK, Bisset L, Brooks P et al. Effect of corticosteroid injection, physiotherapy, or both on clinical outcomes in patients with unilateral lateral epicondylalgia: a randomized controlled trial. *JAMA* 2013;309(5):461-9.
 42. Bisset L, Smidt N, van der Windt DA et al. Conservative treatments for tennis elbow—do subgroups of patients respond differently? *Rheumatology (Oxford)* 2007;46(10):1601-5.
 43. Murrell GAC. Using nitric oxide to treat tendinopathy. *Br J Sports Med* 2007;41(4):227-31.
 44. Paoloni JA, Murrell GA, Burch RM et al. Randomised, double blind, placebo-controlled clinical trial of a new topical glyceryl trinitrate patch for chronic lateral epicondylitis. *Br J Sports Med* 2009;43(4):299-302.
 45. Espandar R, Heidari P, Rasouli MR et al. Use of anatomic measurement to guide injection of botulinum toxin for the management of chronic lateral epicondylitis: a randomized controlled trial. *CMAJ* 2010;182(8):768-73.
 46. Vicenzino B, Coombes BK. A single botulinum toxin injection at a precise anatomic point on the forearm reduces pain at rest, compared to placebo injection in patients with chronic refractory lateral epicondylitis. *Evid Based Med* 2010;15(5):149-50.
 47. Hall MP, Ward JP, Cardone DA. Platelet rich placebo? Evidence for platelet rich plasma in the treatment of tendinopathy and augmentation of tendon repair. *Bull Hosp Jt Dis (2013)* 2013;71(1):54-9.
 48. de Mos M, van der Windt AE, Jahr H et al. Can platelet-rich plasma enhance tendon repair? A cell culture study. *Am J Sports Med* 2008;36(6):1171-8.
 49. Jo CH, Kim JE, Yoon KS et al. Platelet-rich plasma stimulates cell proliferation and enhances matrix gene expression and synthesis in tenocytes from human rotator cuff tendons with degenerative tears. *Am J Sports Med* 2012;40(5):1035-45.
 50. de Vos RJ, Windt J, Weir A. Strong evidence against platelet-rich plasma injections for chronic lateral epicondylar tendinopathy: a systematic review. *Br J Sports Med* 2014;48(12):952-6.
 51. Moraes VY, Lenza M, Tamaoki MJ et al. Platelet-rich therapies for musculoskeletal soft tissue injuries. *Cochrane Database Syst Rev* 2014(4):Cd010071.
 52. Gosens T, Mishra AK. Editorial in response to the systematic review by de Vos et al: 'Strong evidence against platelet-rich plasma injections for chronic lateral epicondylar tendinopathy: a systematic review'. *Br J Sports Med* 2014;48(12):945-6.
 53. de Vos RJ, Weir A, Brasher PM et al. Platelet-rich plasma for chronic tennis elbow: letters to the editor. *Am J Sports Med* 2014;42(1):Np3-5.
 54. Wang A, Breidahl W, Mackie KE et al. Autologous tenocyte injection for the treatment of severe, chronic resistant lateral epicondylitis: a pilot study. *Am J Sports Med* 2013;41(12):2925-32.
 55. Karkhanis S, Frost A, Maffulli N. Operative management of tennis elbow: a quantitative review. *Br Med Bull* 2008;88(1):171-88.
 56. Ciccotti MC, Schwartz MA, Ciccotti MG. Diagnosis and treatment of medial epicondylitis of the elbow. *Clin Sports Med* 2004;23(4):693-705.
 57. Park GY, Lee SM, Lee MY. Diagnostic value of ultrasonography for clinical medial epicondylitis. *Arch Phys Med Rehabil* 2008;89(4):738-42.
 58. Rossy WH, Oh LS. Pitcher's elbow: medial elbow pain in the overhead-throwing athlete. *Curr Rev Musculoskelet Med* 2016;9(2):207-14.
 59. Jobe FW, Kvitne RS. Elbow instability in the athlete. *Instr Course Lect* 1991;40:17-23.
 60. King JW, Brelsford HJ, Tullos HS. Analysis of the pitching arm of the professional baseball pitcher. *Clin Orthop Relat Res* 1969;67:116-23.
 61. Vitale MA, Ahmad CS. The outcome of elbow ulnar collateral ligament reconstruction in overhead athletes: a systematic review. *Am J Sports Med* 2008;36(6):1193-205.
 62. Frostick SP, Mohammad M, Ritchie DA. Sport injuries of the elbow. *Br J Sports Med* 1999;33(5):301-11.
 63. Jacobson JA, Jebson PJ, Jeffers AW et al. Ulnar nerve dislocation and snapping triceps syndrome: diagnosis with dynamic sonography—report of three cases. *Radiology* 2001;220(3):601-5.
 64. Appelboom A, Reuben AD, Bengel JR et al. Elbow extension test to rule out elbow fracture: multicentre, prospective validation and observational study of diagnostic accuracy in adults and children. *BMJ* 2008;337:a2428.
 65. Shapiro MS, Wang JC. Elbow fractures: treating to avoid complications. *Phys Sportsmed* 1995;23(4):39-50.
 66. Ellerin BE, Helfet D, Parikh S et al. Current therapy in the management of heterotopic ossification of the elbow: a review with case studies. *Am J Phys Med Rehabil* 1999;78(3):259-71.
 67. Matar HE, Ali AA, Buckley S et al. Surgical interventions for treating fractures of the olecranon in adults. *Cochrane Database Syst Rev* 2014(11):Cd010144.
 68. Mason ML. Some observations on fractures of the head of the radius with a review of one hundred cases. *Br J Surg* 1954;42(172):123-32.
 69. Popovic N, Gillet P, Rodriguez A et al. Fracture of the radial head with associated elbow dislocation: results of treatment using a floating radial head prosthesis. *J Orthop Trauma* 2000;14(3):171-7.
 70. Kodde IF, Kaas L, Flipsen M et al. Current concepts in the management of radial head fractures. *World J Orthop* 2015;6(11):954-60.
 71. Popovic N, Lemaire R, Georis P et al. Midterm results with a bipolar radial head prosthesis: radiographic evidence of loosening at the bone-cement interface. *J Bone Joint Surg Am* 2007;89(11):2469-76.
 72. Regan W, Morrey BF. Classification and treatment of coronoid process fractures. *Orthopedics* 1992;15(7):845-8.
 73. O'Driscoll SW, Morrey BF, Korinek S et al. Elbow subluxation and dislocation. A spectrum of instability. *Clin Orthop Relat Res* 1992;280:186-97.

74. O'Driscoll SW, Jupiter JB, King GJ et al. The unstable elbow. *Instr Course Lect* 2001;50:89-102.
75. Ross G, McDevitt ER, Chronister R et al. Treatment of simple elbow dislocation using an immediate motion protocol. *Am J Sports Med* 1999;27(3):308-11.
76. Verrall GM. Return to Australian rules football after acute elbow dislocation: a report of three cases and review of the literature. *J Sci Med Sport* 2001;4(2):245-50.
77. O'Driscoll SW, Bell DF, Morrey BF. Posterolateral rotatory instability of the elbow. *J Bone Joint Surg Am* 1991;73(3):440-6.
78. Regan W, Lapner PC. Prospective evaluation of two diagnostic apprehension signs for posterolateral instability of the elbow. *J Shoulder Elbow Surg* 2006;15(3):344-6.
79. Anakwenze OA, Kwon D, O'Donnell E et al. Surgical treatment of posterolateral rotatory instability of the elbow. *Arthroscopy* 2014;30(7):866-71.
80. Fedorka CJ, Oh LS. Posterolateral rotatory instability of the elbow. *Curr Rev Musculoskelet Med* 2016;9(2):240-6.
81. Kenter K, Behr CT, Warren RF et al. Acute elbow injuries in the National Football League. *J Shoulder Elbow Surg* 2000;9(1):1-5.
82. Tyrdal S, Bahr R. High prevalence of elbow problems among goalkeepers in European team handball-'handball goalie's elbow'. *Scand J Med Sci Sports* 1996;6(5):297-302.
83. Popovic N, Lemaire R. Hyperextension trauma to the elbow: radiological and ultrasonographic evaluation in handball goalkeepers. *Br J Sports Med* 2002;36(6):452-6.
84. Konjengbam M, Elangbam J. Radial nerve in the radial tunnel: anatomic sites of entrapment neuropathy. *Clin Anat* 2004;17(1):21-5.
85. Sarhadi NS, Korday SN, Bainbridge LC. Radial tunnel syndrome: diagnosis and management. *J Hand Surg Br* 1998;23(5):617-9.
86. Lee JC, Malara FA, Wood T et al. MRI of stress reaction of the distal humerus in elite tennis players. *AJR Am J Roentgenol* 2006;187(4):901-4.
7. Ellis JA, Kierulf JC, Klassen TP. Injuries associated with in-line skating from the Canadian hospitals injury reporting and prevention program database. *Can J Public Health* 1995;86(2):133-6.
8. Aldridge JM, 3rd, Mallon WJ. Hook of the hamate fractures in competitive golfers: results of treatment by excision of the fractured hook of the hamate. *Orthopedics* 2003;26(7):717-9.
9. Lajoie AS, McCabe SJ, Thomas B et al. Determining the sensitivity and specificity of common diagnostic tests for carpal tunnel syndrome using latent class analysis. *Plast Reconstr Surg* 2005;116(2):502-07.
10. Valdes K, LaStayo P. The value of provocative tests for the wrist and elbow: a literature review. *J Hand Ther* 2013;26(1):32-42; quiz 43.
11. Wolfe SW, Crisco JJ. Mechanical evaluation of the scaphoid shift test. *J Hand Surg Am* 1994;19(5):762-8.
12. La Stayo P, Howell J. Clinical provocative tests used in evaluating wrist pain: a descriptive study. *J Hand Ther* 1995;8:10-17.
13. Tay SC, Tomita K, Berger RA. The 'ulnar fovea sign' for defining ulnar wrist pain: an analysis of sensitivity and specificity. *J Hand Surg Am* 2007;32(4):438-44.
14. Lester B, Halbrecht J, Levy IM et al. 'Press test' for office diagnosis of triangular fibrocartilage complex tears of the wrist. *Ann Plast Surg* 1995;35(1):41-5.
15. Joy CM, Robert SR, Allan D et al. Responsiveness of the short form-36, disability of the arm, shoulder, and hand questionnaire, patient-rated wrist evaluation, and physical impairment measurements in evaluating recovery after a distal radius fracture. *J Hand Surg Am* 2000;25(2):330-40.
16. MacDermid JC, Turgeon T, Richards RS et al. Patient rating of wrist pain and disability: a reliable and valid measurement tool. *J Orthop Trauma* 1998;12(8):577-86.
17. Beaton DE, Katz JN, Fossel AH et al. Measuring the whole or the parts? Validity, reliability, and responsiveness of the Disabilities of the Arm, Shoulder and Hand outcome measure in different regions of the upper extremity. *J Hand Ther* 2001;14(2):128-46.
18. The DASH Outcome Measure. <http://www.dash.iwh.on.ca> Accessed 25 May 2016.
19. Buijze GA, Ring D. Clinical impact of united versus nonunited fractures of the proximal half of the ulnar styloid following volar plate fixation of the distal radius. *J Hand Surg Am* 2010;35(2):223-7.
20. Walenkamp MM, de Muinck Keizer RJ, Goslings JC et al. The minimum clinically important Difference of the Patient-rated Wrist Evaluation Score for Patients with distal radius fractures. *Clin Orthop Relat Res* 2015;473(10):3235-41.
21. Mintken PE, Glynn P, Cleland JA. Psychometric properties of the shortened disabilities of the Arm, Shoulder, and Hand Questionnaire (QuickDASH) and Numeric Pain Rating Scale in patients with shoulder pain. *J Shoulder Elbow Surg* 2009;18(6):920-6.

CHAPTER 26

1. Rettig AC. Athletic injuries of the wrist and hand. Part I: traumatic injuries of the wrist. *Am J Sports Med* 2003;31(6):1038-48.
2. Hill C, Riaz M, Mozzam A et al. A regional audit of hand and wrist injuries. A study of 4873 injuries. *J Hand Surg Br* 1998;23(2):196-200.
3. Parmelee-Peters K, Eathorne SW. The wrist: common injuries and management. *Prim Care* 2005;32(1):35-70.
4. Dekker R, Groothoff JW, van der Sluis CK et al. Long-term disabilities and handicaps following sports injuries: outcome after outpatient treatment. *Disabil Rehabil* 2003;25(20):1153-7.
5. Chow TK, Corbett SW, Farstad DJ. Spectrum of injuries from snowboarding. *J Trauma* 1996;41(2):321-5.
6. Heller DR, Routley V, Chambers S. Rollerblading injuries in young people. *J Paediatr Child Health* 1996;32(1):35-8.

22. Beaton DE, Wright JG, Katz JN et al. Development of the QuickDASH: comparison of three item-reduction approaches. *J Bone Joint Surg Am* 2005;87(5):1038-46.
23. Vendittoli PA, Sonia J, Davison KS et al. Descriptive study of osteoporotic fractures and hip fracture risk evaluation of subjects with past minor fractures. *Orthopedic Proceedings* 2008;90-B(Suppl 1 71).
24. Lichtman DM, Bindra RR, Boyer MI et al. Treatment of distal radius fractures. *J Am Acad Orthop Surg* 2010;18(3):180-9.
25. Downing ND, Karantana A. A revolution in the management of fractures of the distal radius? *J Bone Joint Surg Br* 2008;90(10):1271-5.
26. Tang JB. Distal radius fracture: diagnosis, treatment, and controversies. *Clin Plast Surg* 2014;41(3):481-99.
27. Schnependahl J, Windolf J, Kaufmann RA. Distal radius fractures: current concepts. *J Hand Surg Am* 2012;37(8):1718-25.
28. Lozano-Calderon SA, Souer S, Mudgal C et al. Wrist mobilization following volar plate fixation of fractures of the distal part of the radius. *J Bone Joint Surg Am* 2008;90(6):1297-304.
29. Glickel SZ, Catalano LW, Raia FJ et al. Long-term outcomes of closed reduction and percutaneous pinning for the treatment of distal radius fractures. *J Hand Surg Am* 2008;33(10):1700-5.
30. Handoll HH, Huntley JS, Madhok R. Different methods of external fixation for treating distal radial fractures in adults. *Cochrane Database Syst Rev* 2008(1):CD006522.
31. Matzkin E, Singer DI. Scaphoid stress fracture in a 13-year-old gymnast: a case report. *J Hand Surg Am* 2000;25(4):710-3.
32. Banks LN, Trail IA. Local vascular bone graft reconstruction. In: Trail IA, Fleming ANM, eds. *Disorders of the Hand*. London: Springer, 2015:153-60.
33. Handoll HH, Watts AC. Bone grafts and bone substitutes for treating distal radial fractures in adults. *Cochrane Database Syst Rev* 2008(2):CD006836.
34. Suh N, Ek ET, Wolfe SW. Carpal fractures. *J Hand Surg Am* 2014;39(4):785-91; quiz 791.
35. DiFiori JP, Puffer JC, Aish B et al. Wrist pain, distal radial physeal injury, and ulnar variance in young gymnasts: does a relationship exist? *Am J Sports Med* 2002;30(6):879-85.
36. DiFiori JP, Caine DJ, Malina RM. Wrist pain, distal radial physeal injury, and ulnar variance in the young gymnast. *Am J Sports Med* 2006;34(5):840-9.
37. Grant-Ford M, Sitler MR, Kozin SH et al. Effect of a prophylactic brace on wrist and ulnocarpal joint biomechanics in a cadaveric model. *Am J Sports Med* 2003;31(5):736-43.
38. Peters-Veluthamaningal C, van der Windt DA, Winters JC et al. Corticosteroid injection for de Quervain's tenosynovitis. *Cochrane Database Syst Rev* 2009(3):CD005616.
39. McNally E, Wilson D, Seiler S. Rowing injuries. *Semin Musculoskelet Radiol* 2005;9(4):379-96.
40. Williams JG. Surgical management of traumatic non-infective tenosynovitis of the wrist extensors. *J Bone Joint Surg Br* 1977;59-B(4):408-10.
41. Markiewicz AD, Merryman J. Radial nerve compression in the upper extremity. *J Am Soc Surg Hand* 2005;5(2):87-99.
42. Couzens G, Daunt N, Crawford R et al. Positive magnetic resonance imaging findings in the asymptomatic wrist. *ANZ J Surg* 2014;84(7-8):528-32.
43. Slade JF, 3rd, Milewski MD. Management of carpal instability in athletes. *Hand Clin* 2009;25(3):395-408.
44. Jones DB, Jr., Kakar S. Perilunate dislocations and fracture dislocations. *J Hand Surg Am* 2012;37(10):2168-73; quiz 2174.
45. Vitale MA, Seetharaman M, Ruchelsman DE. Perilunate dislocations. *J Hand Surg Am* 2015;40(2):358-62; quiz 362.
46. Forli A, Courvoisier A, Wimsey S et al. Perilunate dislocations and transscaphoid perilunate fracture-dislocations: a retrospective study with minimum ten-year follow-up. *J Hand Surg Am* 2010;35(1):62-8.
47. Talwalkar SC, Edwards AT, Hayton MJ et al. Results of tri-ligament tenodesis: a modified Brunelli procedure in the management of scapholunate instability. *J Hand Surg Br* 2006;31(1):110-7.
48. Ross M, Loveridge J, Cutbush K et al. Scapholunate ligament reconstruction. *J Wrist Surg* 2013;2(2):110-5.
49. Herzberg G, Mercier S, Charbonnier JP et al. Kienböck's disease in a 14-year-old gymnast: a case report. *J Hand Surg Am* 2006;31(2):264-8.
50. Wollstein R, Wollstein A, Rodgers J et al. A hand therapy protocol for the treatment of lunare overload or early Kienböck's disease. *J Hand Ther* 2013;26(3):255-59; quiz 260.
51. Viljakka T, Tallroth K, Vastamaki M. Long-term outcome (20 to 33 years) of radial shortening osteotomy for Kienböck's lunatomalacia. *J Hand Surg Eur Vol* 2014;39(7):761-9.
52. Beredjikian PK. Kienböck's disease. *J Hand Surg Am* 2009;34(1):167-75.
53. Schmitt R, Heinze A, Fellner F et al. Imaging and staging of avascular osteonecroses at the wrist and hand. *Eur J Radiol* 1997;25(2):92-103.
54. Innes L, Strauch RJ. Systematic review of the treatment of Kienböck's disease in its early and late stages. *J Hand Surg Am* 2010;35(5):713-7, 17 e1-4.
55. Sammer DM, Chung KC. Management of the distal radioulnar joint and ulnar styloid fracture. *Hand Clin* 2012;28(2):199-206.
56. Wysocki RW, Ruch DS. Ulnar styloid fracture with distal radius fracture. *J Hand Surg Am* 2012;37(3):568-9.
57. Daneshvar P, Chan R, MacDermid J et al. The effects of ulnar styloid fractures on patients sustaining distal radius fractures. *J Hand Surg Am* 2014;39(10):1915-20.
58. Jacobson JA, Miller BS, Morag Y. Golf and racquet sports injuries. *Semin Musculoskelet Radiol* 2005;9(4):346-59.
59. Theriault G, Lachance P. Golf injuries. An overview. *Sports Med* 1998;26(1):43-57.
60. Scheufler O, Radmer S, Erdmann D et al. Therapeutic alternatives in nonunion of hamate hook fractures: personal experience in 8 patients and review of literature. *Ann Plast Surg* 2005;55(2):149-54.

61. Marchessault J, Conti M, Baratz ME. Carpal fractures in athletes excluding the scaphoid. *Hand Clin* 2009;25(3):371-88.
62. Suh N, Ek ET, Wolfe SW. Carpal fractures. *J Hand Surg Am* 2014;39(4):785-91.
63. Atkinson CT, Watson J. Lunotriquetral ligament tears. *J Hand Surg Am* 2012;37(10):2142-4.
64. Goldberg SH, Strauch RE, Rosenwasser MP. Scapholunate and lunotriquetral instability in the athlete: diagnosis and management. *Oper Techn Sport Med* 2006;14(2):108-21.
65. Leon-Lopez MM, Salva-Coll G, Garcia-Elias M et al. Role of the extensor carpi ulnaris in the stabilization of the lunotriquetral joint. An experimental study. *J Hand Ther* 2013;26(4):312-7; quiz 317.
66. Ko JH, Wiedrich TA. Triangular fibrocartilage complex injuries in the elite athlete. *Hand Clin* 2012;28(3):307-21, viii.
67. Berna-Serna JD, Martinez F, Reus M et al. Wrist arthrography: a simple method. *Eur Radiol* 2006;16(2):469-72.
68. De Smet L. Magnetic resonance imaging for diagnosing lesions of the triangular fibrocartilage complex. *Acta Orthop Belg* 2005;71(4):396-8.
69. Bachoura A, Wroblewski A, Jacoby SM et al. Hook of hamate fractures in competitive baseball players. *Hand (N Y)* 2013;8(3):302-7.
70. Keogh CF, Wong AD, Wells NJ et al. High-resolution sonography of the triangular fibrocartilage: initial experience and correlation with MRI and arthroscopic findings. *AJR Am J Roentgenol* 2004;182(2):333-6.
71. Tomaino MM, Weiser RW. Combined arthroscopic TFCC debridement and wafer resection of the distal ulna in wrists with triangular fibrocartilage complex tears and positive ulnar variance. *J Hand Surg Am* 2001;26(6):1047-52.
72. Ozer K. A new surgical technique for the ligament reconstruction of the trapeziometacarpal joint. *Tech Hand Up Extrem Surg* 2006;10(3):181-6.
73. Salva-Coll G, Garcia-Elias M, Leon-Lopez MM et al. Role of the extensor carpi ulnaris and its sheath on dynamic carpal stability. *J Hand Surg Eur Vol* 2012;37(6):544-8.
74. Iorio ML, Bayomy AF, Huang JI. Morphology of the extensor carpi ulnaris groove and tendon. *J Hand Surg Am* 2014;39(12):2412-6.
75. Iorio ML, Huang JI. Extensor carpi ulnaris subluxation. *J Hand Surg Am* 2014;39(7):1400-2.
76. Rettig AC. Athletic injuries of the wrist and hand: part II: overuse injuries of the wrist and traumatic injuries to the hand. *Am J Sports Med* 2004;32(1):262-73.
77. Montalvan B, Parier J, Brasseur JL et al. Extensor carpi ulnaris injuries in tennis players: a study of 28 cases. *Br J Sports Med* 2006;40(5):424-9; discussion 429.
78. Stroia K, Baudo, M., Martin, K. Taping techniques for TFCC and ECU injuries on the Sony Ericsson WTA tour. *Med Sci Tennis* 2009;14(1):15-19.
79. Graham TJ. Pathologies of the extensor carpi ulnaris (ECU) tendon and its investments in the athlete. *Hand Clin* 2012;28(3):345-56, ix.
80. Schrijver HM, Gerritsen AA, Strijers RL et al. Correlating nerve conduction studies and clinical outcome measures on carpal tunnel syndrome: lessons from a randomized controlled trial. *J Clin Neurophysiol* 2005;22(3):216-21.
81. Dincer U, Cakar E, Kiralp MZ et al. The effectiveness of conservative treatments of carpal tunnel syndrome: splinting, ultrasound, and low-level laser therapies. *Photomed Laser Surg* 2009;27(1):119-25.
82. Milo R, Kalichman L, Volchek L et al. Local corticosteroid treatment for carpal tunnel syndrome: a 6-month clinical and electrophysiological follow-up study. *J Back Musculoskelet Rehabil* 2009;22(2):59-64.
83. Wong SM, Hui AC, Lo SK et al. Single vs. two steroid injections for carpal tunnel syndrome: a randomised clinical trial. *Int J Clin Pract* 2005;59(12):1417-21.
84. Hui AC, Wong S, Leung CH et al. A randomized controlled trial of surgery vs steroid injection for carpal tunnel syndrome. *Neurology* 2005;64(12):2074-8.
85. Thoma A, Veltri K, Haines T et al. A systematic review of reviews comparing the effectiveness of endoscopic and open carpal tunnel decompression. *Plast Reconstr Surg* 2004;113(4):1184-91.
86. Capitani D, Beer S. Handlebar palsy—a compression syndrome of the deep terminal (motor) branch of the ulnar nerve in biking. *J Neurol* 2002;249(10):1441-5.
87. Ginn TA, Smith AM, Snyder JR et al. Vascular changes of the hand in professional baseball players with emphasis on digital ischemia in catchers. *J Bone Joint Surg Am* 2005;87(7):1464-9.
88. Buijze GA, Goslings JC, Rhemrev SJ et al. Cast immobilization with and without immobilization of the thumb for nondisplaced and minimally displaced scaphoid waist fractures: a multicenter, randomized, controlled trial. *J Hand Surg Am* 2014;39(4):621-7.
89. Kitay A, Wolfe SW. Scapholunate instability: current concepts in diagnosis and management. *J Hand Surg Am* 2012;37(10):2175-96.

CHAPTER 27

1. Hill C, Riaz M, Mozzam A et al. A regional audit of hand and wrist injuries. A study of 4873 injuries. *J Hand Surg Br* 1998;23(2):196-200.
2. Feddermann-Demont N, Junge A, Edouard P et al. Injuries in 13 international athletics championships between 2007-12. *Br J Sports Med* 2014;48(7):513-22.
3. Chung KC, Pillsbury MS, Walters MR et al. Reliability and validity testing of the Michigan Hand Outcomes Questionnaire. *J Hand Surg Am* 1998;23A:575-87.
4. Beaton DE, Katz JN, Fossel AH et al. Measuring the whole or the parts? Validity, reliability, and responsiveness of the Disabilities of the Arm, Shoulder and Hand outcome measure in different regions of the upper extremity. *J Hand Ther* 2001;14(2):128-46.
5. Mintken PE, Glynn P, Cleland JA. Psychometric properties of the shortened disabilities of the Arm, Shoulder, and Hand Questionnaire (QuickDASH) and Numeric Pain Rating

- Scale in patients with shoulder pain. *J Shoulder Elbow Surg* 2009;18(6):920-6.
6. Beaton DE, Wright JG, Katz JN et al. Development of the QuickDASH: comparison of three item-reduction approaches. *J Bone Joint Surg* 2005;87-A(5):1038-46.
 7. Ridha H, Crerar-Gilbert A, Fleming A. The use of Tegaderm for finger buddy-strapping: a dressing with many advantages. *Ann R Coll Surg Engl* 2008;90(6):525.
 8. Oetgen ME, Dodds SD. Non-operative treatment of common finger injuries. *Curr Rev Musculoskelet Med* 2008;1(2):97-102.
 9. Kennedy S, Peck F, Stone J. Balancing contracture management and function: the treatment of interphalangeal joint flexion contractures with reinforced Lycra finger sleeves. *J Hand Ther* 2000;13(1):52-5.
 10. Singletary S, Geissler WB. Bracing and rehabilitation for wrist and hand injuries in collegiate athletes. *Hand Clin* 2009;25(3):443-8.
 11. Paulius K, Maguina P, Mejia A. Ultrasound-guided management of hand fractures. *Orthopedics* 2008;31(12). pii: orthosupersite.com/view.aspx?rID=34709.
 12. Body R, Ferguson CJ. Best evidence topic report. Early mobilisation for volar plate avulsion fractures. *Emerg Med J* 2005;22(7):505.
 13. Nazarian N, Page RS, Hoy GA et al. Combined joint fusion for index and middle carpometacarpal instability in elite boxers. *J Hand Surg Eur* 2014;39(3):242-8.
 14. Bindra RR, Foster BJ. Management of proximal interphalangeal joint dislocations in athletes. *Hand Clin* 2009;25(3):423-35.
 15. Thayer DT. Distal interphalangeal joint injuries. *Hand Clin* 1988;4:1-4.
 16. Banerji S, Bullocks J, Cole P et al. Irreducible distal interphalangeal joint dislocation: a case report and literature review. *Ann Plast Surg* 2007;58(6):683-5.
 17. Calfee RP, Sommerkamp TG. Fracture-dislocation about the finger joints. *J Hand Surg Am* 2009;34(6):1140-7.
 18. Lee AT, Carlson MG. Thumb metacarpophalangeal joint collateral ligament injury management. *Hand Clin* 2012;28(3):361-70, ix-x.
 19. Carlson MG. Commentary on RCL/UCL injury in basketball. *Hand Clin* 2012;28(3):373-5.
 20. Werner BC, Hadeed MM, Lyons ML et al. Return to football and long-term clinical outcomes after thumb ulnar collateral ligament suture anchor repair in collegiate athletes. *J Hand Surg Am* 2014;39(10):1992-8.
 21. Chhor KS, Culp RW. Baseball commentary 'thumb ligament injuries: RCL and UCL'. *Hand Clin* 2012;28(3):371-2.
 22. Williams CS. Thumb metacarpophalangeal joint ligament injury: football commentary. *Hand Clin* 2012;28(3):377-8.
 23. Taylor KF, Lanzi JT, Cage JM et al. Radial collateral ligament injuries of the thumb metacarpophalangeal joint: epidemiology in a military population. *J Hand Surg Am* 2013;38(3):532-6.
 24. Edelstein DM, Kardashian G, Lee SK. Radial collateral ligament injuries of the thumb. *J Hand Surg Am* 2008;33(5):760-70.
 25. Chinchalkar SJ, Gan BS. Management of proximal interphalangeal joint fractures and dislocations. *J Hand Ther* 2003;16:117-28.
 26. Konecny J, Martinkova J, Kabela M et al. Laser in treatment of finger sprains. *Proceedings of IALMS 2007 conference: Laser Florence: the laser medicine world* 2007:106-8.
 27. Moradi A, Kachooei AR, Mudgal CS. Mallet fracture. *J Hand Surg Am* 2014;39(10):2067-9.
 28. Yeh PC, Shin SS. Tendon ruptures: mallet, flexor digitorum profundus. *Hand Clin* 2012;28(3):425-30, xi.
 29. Elui VM, Rodrigues MC, Purquerio BM et al. Swan neck deformity orthosis development: AOA (articulated ring orthosis). *J Hand Ther* 2012;25(4):e9-e10.
 30. Lourie GM. Boutonnière and pulley rupture football commentary. *Hand Clin* 2012;28(3):451-2.
 31. Weiland AJ. Boutonnière and pulley rupture in elite baseball players. *Hand Clin* 2012;28(3):447.
 32. Marino JT, Lourie GM. Boutonnière and pulley rupture in elite athletes. *Hand Clin* 2012;28(3):437-45, xii.
 33. Smith DW. Boutonnière and pulley rupture in elite basketball. *Hand Clin* 2012;28(3):449-50.
 34. Evans RB. Early active short arc motion for the closed central slip injury. *J Hand Ther* 2010;23(4):e15-e16.
 35. Schoffl VR, Schoffl I. Finger pain in rock climbers: reaching the right differential diagnosis and therapy. *J Sports Med Phys Fitness* 2007;47(1):70-78.
 36. Schoffl VR, Schoffl I. Injuries to the finger flexor pulley system in rock climbers: current concepts. *J Hand Surg Am* 2006;31A(4):647-54.
 37. Schweizer A. Lumbrical tears in rock climbers. *J Hand Surg Br* 2003;28(2):187-9.
 38. Peters-Veluthamaningal C, Winters JC, Groenier KH et al. Corticosteroid injections effective for trigger finger in adults in general practice: a double-blinded randomised placebo controlled trial. *Ann Rheum Dis* 2008;67(9):1262-6.
 39. Nimigan AS, Ross DC, Gan BS. Steroid injections in the management of trigger fingers. *Am J Phys Med Rehabil* 2006;85(1):36-43.

CHAPTER 28

1. Benson MK, Byrnes DP. The clinical syndromes and surgical treatment of thoracic intervertebral disc prolapse. *J Bone Joint Surg Br* 1975;57(4):471-7.
2. Singer K, Edmondston J. Introduction. The enigma of the thoracic spine. In: Singer K, Giles L, eds. *Clinical Anatomy and Management of Thoracic Spine Pain*. Oxford: Butterworth-Heinemann, 2000:3-13.
3. Papadakos N, Georges H, Sibtain N et al. Thoracic disc prolapse presenting with abdominal pain: case report and review of the literature. *Ann R Coll Surg Engl* 2009;91(5):W4-6. doi: 10.1308/147870809X401038.
4. Groen G, Stolker R. Thoracic neural anatomy. In: Singer K, Giles L, eds. *Clinical Anatomy and Management of Thoracic Spine Pain*. Oxford: Butterworth-Heinemann, 2000:114-41.
5. Maigne J. Cervicothoracic and thoracolumbar spinal pain syndromes. In: Singer K, Giles L, eds. *Clinical Anatomy and*

- Management of Thoracic Spine Pain*. Oxford: Butterworth-Heinemann, 2000:157-68.
6. Boyle J, Singer K, Milne N. Clinical anatomy of the cervicothoracic junction. In: Giles L, Singer K, eds. *Clinical Anatomy and Management of Cervical Spine Pain*. Oxford: Butterworth-Heinemann, 1998:100-13.
 7. Edwards B. *Manual of Combined Movements*. 2nd ed. Oxford: Butterworth-Heinemann, 1999.
 8. Singer K, Edmondston J. Pathoanatomical characteristics of the thoracolumbar junction region. In: Singer K, Giles L, eds. *Clinical Anatomy and Management of Thoracic Spine Pain*. Oxford: Butterworth-Heinemann, 2000:100-13.
 9. Reinhold M, Knop C, Beisse R et al. Operative treatment of 733 patients with acute thoracolumbar spinal injuries: comprehensive results from the second, prospective, internet-based multicenter study of the Spine Study Group of the German Association of Trauma Surgery. *Eur Spine J* 2010;19(10):1657-76. doi: 10.1007/s00586-010-1451-5.
 10. Bogduk N, Jull G. The theoretical pathology of acute locked back: a basis for manipulative therapy. *Manual Medicine* 1985;1:78-82.
 11. Ashton-Miller JA. Thoracic hyperkyphosis in the young athlete: a review of the biomechanical issues. *Curr Sports Med Rep* 2004;3(1):47-52.
 12. Wood KB, Garvey TA, Gundry C et al. Magnetic resonance imaging of the thoracic spine. Evaluation of asymptomatic individuals. *J Bone Joint Surg Am* 1995;77(11):1631-8.
 13. Whitcomb DC, Martin SP, Schoen RE et al. Chronic abdominal pain caused by thoracic disc herniation. *Am J Gastroenterol* 1995;90(5):835-7.
 14. Singer K. Zeit für einen Paradigmenwechsel? Vom T4-Syndrom zur exakten Diagnose. *Manuelle Therapie* 2013;17:103-7.
 15. Boyle JJ. Is the pain and dysfunction of shoulder impingement lesion really second rib syndrome in disguise? Two case reports. *Man Ther* 1999;4(1):44-8.
 16. Farfan H. The tired neck syndrome. In: Singer K, Giles L, eds. *Clinical Anatomy and Management of Thoracic Spine Pain*. Oxford: Butterworth-Heinemann, 2000:171-6.
 17. Bosomworth NJ. Atrial fibrillation and physical activity: should we exercise caution? *Can Fam Physician* 2015;61(12):1061-70.
 18. Bicos J, Nicholson GP. Treatment and results of sternoclavicular joint injuries. *Clin Sports Med* 2003;22(2):359-70.
 19. Garretson RB, 3rd, Williams GR, Jr. Clinical evaluation of injuries to the acromioclavicular and sternoclavicular joints. *Clin Sports Med* 2003;22(2):239-54.
 20. Wroble RR. Sternoclavicular injuries: managing damage to an overlooked joint. / Blessures de la ceinture scapulaire: soigner l'articulation en cause. *Phys Sportsmed* 1995;23(9):19-20;21-24;26.
 21. Williams CC. Posterior sternoclavicular joint dislocation. *Phys Sportsmed* 1999;27(2):105-13. doi: 10.3810/psm.1999.02.678.
 22. Kiroff GK, McClure DN, Skelley JW. Delayed diagnosis of posterior sternoclavicular joint dislocation. *Med J Aust* 1996;164(4):242-3.
 23. Mirza AH, Alam K, Ali A. Posterior sternoclavicular dislocation in a rugby player as a cause of silent vascular compromise: a case report. *Br J Sports Med* 2005;39(5):e28.
 24. Gregory PL, Biswas AC, Batt ME. Musculoskeletal problems of the chest wall in athletes. *Sports Med* 2002;32(4):235-50.
 25. Brukner P, Bennell K, Matheson G. *Stress Fractures*. Melbourne: Blackwell Scientific, Asia, 1999.
 26. Hopper MA, Tirman P, Robinson P. Muscle injury of the chest wall and upper extremity. *Semin Musculoskelet Radiol* 2010;14(2):122-30. doi: 10.1055/s-0030-1253156.
 27. Evans G, Redgrave A. Great Britain rowing team guideline for diagnosis and management of rib stress injury: part 2—the guideline itself. *Br J Sports Med* 2016;50(5):270-2. doi: 10.1136/bjsports-2015-095928.
 28. Evans G, Redgrave A. Great Britain rowing team guideline for diagnosis and management of rib stress injury: part 1. *Br J Sports Med* 2016;50(5):266-9. doi: 10.1136/bjsports-2015-095126.
 29. Vinther A, Kanstrup IL, Christiansen E et al. Exercise-induced rib stress fractures: potential risk factors related to thoracic muscle co-contraction and movement pattern. *Scand J Med Sci Sports* 2006;16(3):188-96.
 30. Vinther A, Kanstrup IL, Christiansen E et al. Exercise-induced rib stress fractures: influence of reduced bone mineral density. *Scand J Med Sci Sports* 2005;15(2):95-9.
 31. Connell DA, Jhamb A, James T. Side strain: a tear of internal oblique musculature. *AJR Am J Roentgenol* 2003;181(6):1511-1517.
 32. Conte SA, Thompson MM, Marks MA et al. Abdominal muscle strains in professional baseball 1991-2010. *Am J Sports Med* 2012;40(3):650-656.
 33. Humphries D, Jamison M. Clinical and magnetic resonance imaging features of cricket bowler's side strain. *Br J Sports Med* 2004;38(5):e21.
 34. Boyce Cam NJ, Muthukumar N, Boyle S et al. Rib impingement in first class cricketers: case reports of two patients who underwent rib resection. *Br J Sports Med* 2006;40(8):732-733.
 35. Obaid H, Nealon A, Connell D. Sonographic appearance of side strain injury. *AJR Am J Roentgenol* 2008;191(6):W264-267.
 36. O'Neal ML, McKown K, Poulis GC. Complex strain injury involving an intercostal hematoma in a professional baseball player. *Clin J Sport Med* 2008;18(4):372-373.
 37. Stevens KJ, Crain JM, Akizuki KH et al. Imaging and ultrasound-guided steroid injection of internal oblique muscle strains in baseball pitchers. *Am J Sports Med* 2010;38(3):581-585.
 38. Maquirriain J, Ghisi JP. Uncommon abdominal muscle injury in a tennis player: internal oblique strain. *Br J Sports Med* 2006;40(5):462-463.
 39. Nealon AR, Kountouris A, Cook JL. Side strain in sport: a narrative review of pathomechanics, diagnosis, imaging and management for the clinician. *J Sci Med Sport* 2016 Aug 28. pii: S1440-2440(16)30159-1. doi: 10.1016/j.jsams.2016.08.016. [Epub ahead of print]
 40. Orchard J, Kountouris A, Portus M et al. Cricket Australia 2014 Injury Report. *Sport Health* 2014;32(4):14.

41. Ranson C, King M, Burnett A et al. The effect of coaching intervention on elite fast bowling technique over a two year period. *Sports Biomech* 2009;8(4):261-274.
42. Humphries D, Orchard J, Kountouris A. Abdominal wall injuries at the elite level in Australian male professional cricketers. *J Postgrad Med Edu Res* 2015;49(4): 155-158.

CHAPTER 29

1. O'Sullivan P. It's time for change with the management of non-specific chronic low back pain. *Br J Sports Med* 2012;46:224-7.
2. Deyo R, Mirza S, Turner J. Over treating chronic back pain: time to back off? *J Am Board Fam Med* 2009;22:62-8.
3. Gatchel R, Peng Y, Peters M et al. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychol Bull* 2007;133:581-624.
4. Deyo R. Diagnostic evaluation of LBP. Reaching a specific diagnosis is often impossible. *Arch Int Med* 2002;162(13): 1444-7.
5. Brinjikji W, Luetmer P, Comstock B et al. Systematic literature review of imaging features of spinal degeneration in asymptomatic populations. *Am J Neuroradiol* 2015;36(4):811-6.
6. Jarvik JG, Hollingworth W, Heagerty PJ et al. Three-year incidence of low back pain in an initially asymptomatic cohort: clinical and imaging risk factors. *Spine* 2005;30:1541-8.
7. Webster BS, Cifuentes M. Relationship of early magnetic resonance imaging for work-related acute low back pain with disability and medical utilization outcomes. *J Occ Environ Med* 2010;52:900-7.
8. Newlands C, Reid D, Parmar P. The prevalence, incidence and severity of low back pain among international-level rowers. *Br J Sports Med* 2015;49(14):951-6.
9. Smith A, O'Sullivan P, Straker L. Classification of sagittal thoraco-lumbo-pelvic alignment of the adolescent spine in standing and its relationship to low back pain. *Spine* 2008;1(33):2101-7.
10. O'Sullivan P, Grahamslaw K, Kendell M et al. The effect of different standing and sitting postures on trunk muscle activity in a pain free population. *Spine* 2002;27:1238-44.
11. Wade M, Campbell A, Norcott J et al. Are static lumbar postures a signature for predicting landing postures in elite female gymnasts? *J App Biomech* 2012;28(6):677-86.
12. O'Sullivan P. Diagnosis and classification of chronic low back pain disorders: maladaptive movement and motor control impairments as underlying mechanism. *Man Ther* 2005;10:242-55.
13. Dankaerts W, O'Sullivan PB, Burnett AF et al. Discriminating healthy controls and two clinical subgroups of nonspecific chronic low back pain patients using trunk muscle activation and lumbosacral kinematics of postures and movements: a statistical classification model. *Spine* 2009;34:1610-8.
14. O'Sullivan PB, Mitchell T, Bulch P et al. The relationship between posture, lumbar muscle endurance and low back pain in industrial workers. *Man Ther* 2006;11(4):264-71.
15. Popovich JM, Jr., Kulig K. Lumbopelvic landing kinematics and EMG in women with contrasting hip strength. *Med Sci Sports Exerc* 2012;44(1):146-53.
16. Ng L, Caneiro JP, Campbell A et al. Cognitive functional approach to manage low back pain in male adolescent rowers: a randomised controlled trial. *Br J Sports Med* 2015;49(17):1125-31.
17. Dijken C, Fjellman-Wiklund A, Hildingsson C. Low back pain, lifestyle factors and physical activity: a population based-study. *J Rehabil Med* 2008;40:864-9.
18. Mitchell T, O'Sullivan P, Burnett A et al. Identification of modifiable personal factors that predict new-onset low back pain: a prospective study of female nursing students. *Clin J Pain* 2009;26:275-83.
19. Dean E, Soderlund A. What is the role of lifestyle behaviour change associated with non-communicable disease risk in managing musculoskeletal health conditions with special reference to chronic pain? *BMC Musculoskelet Disord* 2015;13(16):18.
20. Briggs A, Jordan J, Buchbinder R et al. Health literacy and beliefs among a community cohort with and without chronic low back pain. *Pain* 2010;150:275-83.
21. O'Sullivan P, Lin I. Acute low back pain: beyond drug therapy. *Pain Management Today* 2014;1(1):1-13.
22. Bunzli S, O'Sullivan P, Smith A et al. What do people who score highly on the Tampa Scale of Kinesiophobia really believe? A mixed methods investigation in people with chronic non specific low back pain. *Clin J Pain* 2014;31(7):621-32.
23. Linton SJ, Shaw WS. Impact of psychological factors in the experience of pain. *Phys Ther* 2011;91(5):700-11.
24. Hasenbring M, Hallner D, Klasen B et al. Pain-related avoidance versus endurance in primary care patients with subacute back pain: psychological characteristics and outcome at a 6-month follow-up. *Pain* 2012;153:211-7.
25. Geisser M, Haig A, Wallbom A et al. Pain-related fear, lumbar flexion, and dynamic EMG among persons with chronic musculoskeletal low back pain. *Clin J Pain* 2004;20:61-9.
26. Linton S, Boersma K. Early identification of patients at risk of developing a persistent back problem: the predictive validity of the Orebro Musculoskeletal Pain Questionnaire. *Clin J Pain* 2003;19:80-6.
27. Galambos SA, Terry PC, Moyle GM et al. Psychological predictors of injury among elite athletes. *Br J Sports Med* 2005;39(6):351-4; discussion 354.
28. O'Sullivan P, Waller R, Gardner J et al. Sensory characteristics of chronic non-specific low back pain: a subgroup investigation. *Man Ther* 2014;19(4):311-8.
29. Genevay S, Finckh A, Payer M et al. Elevated levels of tumor necrosis factor-alpha in periradicular fat tissue in patients with radiculopathy from herniated disc. *Spine* 2008;33(19):2041-6.
30. Cahalan R, O'Sullivan P, Purtill H et al. Inability to perform because of pain/injury in elite adult Irish dance: a prospective investigation of contributing factors. *Scand J Med Sci Sports* 2015; 26(6):694-702.

31. O'Keeffe M, Cullinane P, Hurley J et al. What influences patient-therapist interactions in musculoskeletal physiotherapy? A qualitative systematic review and meta-synthesis. *Phys Ther* 2016;96(5):609-22.
32. Hill JC, Dunn KM, Main CJ et al. Subgrouping low back pain: a comparison of the STarT Back Tool with the Orebro Musculoskeletal Pain Screening Questionnaire. *Eur J Pain* 2010;14(1):83-9.
33. Maughan EF, Lewis JS. Outcome measures in chronic low back pain. *Eur Spine J* 2010;19(9):1484-94.
34. Roland M, Morris R. A study of the natural history of back pain. Part I: development of a reliable and sensitive measure of disability in low-back pain. *Spine* 1983;8(2):141-4.
35. Kopec JA, Esdaile JM, Abrahamowicz M et al. The Quebec Back Pain Disability Scale: measurement properties. *Spine* 1995;20(3):341-52.
36. Stratford PW, Finch E, Solomon P et al. Using the Roland-Morris Questionnaire to make decisions about individual patients. *Physio Canada* 1996;48:107-10.
37. Turner JA, Fulton-Kehoe D, Franklin G et al. Comparison of the Roland-Morris Disability Questionnaire and generic health status measures: a population-based study of workers' compensation back injury claimants. *Spine* 2003;28(10):1061-7; discussion 1067.
38. Davidson M, Keating JL. A comparison of five low back disability questionnaires: reliability and responsiveness. *Phys Ther* 2002;82(1):8-24.
39. Linton SJPD, Boersma KMA. Early identification of patients at risk of developing a persistent back problem: the predictive validity of the Orebro Musculoskeletal Pain Questionnaire. *Clin J Pain* 2003;19(2):80-6.
40. Linton SJ, Hallden K. Can we screen for problematic back pain? A screening questionnaire for predicting outcome in acute and subacute back pain. *Clin J Pain* 1998;14(3):209-15.
41. Hockings RL, McAuley JH, Maher CG. A systematic review of the predictive ability of the Orebro Musculoskeletal Pain Questionnaire. *Spine* 2008;33(15):E494-500.
42. Fritz JM, Beneciuk JM, George SZ. Relationship between categorization with the STarT Back Screening Tool and prognosis for people receiving physical therapy for low back pain. *Phys Ther* 2011;91(5):722-32.
43. Hill JC, Dunn KM, Lewis M et al. A primary care back pain screening tool: identifying patient subgroups for initial treatment. *Arth Care Res* 2008;59(5):632-41.
44. Brown TA, Chorpita BF, Korotitsch W et al. Psychometric properties of the Depression Anxiety Stress Scales (DASS) in clinical samples. *Behav Res Ther* 1997;35(1):79-89.
45. George SZ, Valencia C, Beneciuk JM. A psychometric investigation of fear-avoidance model measures in patients with chronic low back pain. *J Orthop Sports Phys Ther* 2010;40(4):197-205.
46. Swinkels-Meewisse EJ, Swinkels RA, Verbeek AL et al. Psychometric properties of the Tampa Scale for kinesiophobia and the fear-avoidance beliefs questionnaire in acute low back pain. *Man Ther* 2003;8(1):29-36.
47. Ostelo RW, Swinkels-Meewisse IJ, Knol DL et al. Assessing pain and pain-related fear in acute low back pain: what is the smallest detectable change? *Int J Behav Med* 2007;14(4):242-8.
48. Freynhagen R, Baron R, Gockel U et al. painDETECT: a new screening questionnaire to identify neuropathic components in patients with back pain. *Curr Med Res Opin* 2006;22(10):1911-20.
49. De Andres J, Perez-Cajaraville J, Lopez-Alarcon MD et al. Cultural adaptation and validation of the painDETECT scale into Spanish. *Clin J Pain* 2012;28(3):243-53.
50. Aina A, May S, Clare H. The centralization phenomenon of spinal symptoms—a systematic review. *Man Ther* 2004;9(3):134-43.
51. Al Nezari NH, Schneiders AG, Hendrick PA. Neurological examination of the peripheral nervous system to diagnose lumbar spinal disc herniation with suspected radiculopathy: a systematic review and meta-analysis. *Spine J* 2013;13(6):657-74.
52. van der Windt DA, Simons E, Riphagen II et al. Physical examination for lumbar radiculopathy due to disc herniation in patients with low-back pain. *Cochrane Database Syst Rev* 2010(2):Cd007431.
53. Majlesi J, Togay H, Unalan H et al. The sensitivity and specificity of the slump and the straight leg raising tests in patients with lumbar disc herniation. *J Clin Rheumatol* 2008;14(2):87-91.
54. Cleland J, Koppenhaver S. *Netter's Orthopaedic Clinical Examination: An Evidence-Based Approach*. 2nd ed. London: Elsevier Health Sciences, 2010.
55. Alqarni AM, Schneiders AG, Cook CE et al. Clinical tests to diagnose lumbar spondylolysis and spondylolisthesis: a systematic review. *Phys Ther Sport* 2015;16(3):268-75.
56. Laslett M, Aprill CN, McDonald B et al. Clinical predictors of lumbar provocation discography: a study of clinical predictors of lumbar provocation discography. *Eur Spine J* 2006;15(10):1473-84.
57. Boden SD, Davis DO, Dina TS et al. Abnormal magnetic-resonance scans of the lumbar spine in asymptomatic subjects. A prospective investigation. *J Bone Joint Surg Am* 1990;72(3):403-8.
58. Babar S, Saifuddin A. MRI of the post-discectomy lumbar spine. *Clin Radiol* 2002;57(11):969-81.
59. Breger RK, Williams AL, Daniels DL et al. Contrast enhancement in spinal MR imaging. *AJR Am J Roentgenol* 1989;153(2):387-91.
60. Wiltse LL, Widell EH, Jackson DW. Fatigue fracture: the basic lesion in the isthmic spondylolisthesis. *J Bone Joint Surg Am* 1975;57-a:17-22.
61. Kobayashi A, Kobayashi T, Kato K et al. Diagnosis of radiographically occult lumbar spondylolysis in young athletes by magnetic resonance imaging. *Am J Sports Med* 2013;41(1):169-76.
62. Herman MJ, Pizzutillo PD. Spondylolysis and spondylolisthesis in the child and adolescent: a new classification. *Clin Orthop Relat Res* 2005;434:46-54.
63. Miyagi R, Sairyo K, Sakai T et al. Two types of laminolysis in adolescent athletes. *J Orthop Traumatol* 2012;13(4):225-8.
64. Standaert CJ, Herring SA. Spondylolysis: a critical review. *Br J Sports Med* 2000;34(6):415-22.

65. Borg B, Modic MT, Obuchowski N et al. Pedicle marrow signal hyperintensity on short tau inversion recovery- and t2-weighted images: prevalence and relationship to clinical symptoms. *Am J Neuroradiol* 2011;32(9):1624-31.
66. Pascal-Mousellard H, Broizat M, Cursolles JC et al. Association of unilateral isthmic spondylolysis with lamina fracture in an athlete: case report and literature review. *Am J Sports Med* 2005;33(4):591-5.
67. Fredrickson B, Baker D, McHolick W et al. The natural history of spondylolysis and spondylolithesis. *J Bone Joint Surg Am* 1984;66A(5):699-707.
68. Leone A, Cianfoni A, Cerase A et al. Lumbar spondylolysis: a review. *Skel Radiol* 2011;40:683-700.
69. Brooks B, Southam S, Mlady G et al. Lumbar spine spondylolysis in the adult population: using computed tomography to evaluate the possibility of adult onset lumbar spondylolysis as a cause of back pain. *Skel Radiol* 2010;(7):669-73.
70. Been E, Ling L, Hunter D et al. Geometry of the vertebral bodies and the intervertebral discs in lumbar segments adjacent to spondylolysis and spondylolisthesis: pilot study. *Eur Spine J* 2011;20(7):1159-65.
71. Klein G, Mehlman CT, McCarty M. Nonoperative treatment of spondylolysis and grade I spondylolisthesis in children and young adults: a meta-analysis of observational studies. *J Pediatr Orthop* 2009;29(2):146-56.
72. Micheli LJ, Wood R. Back pain in young athletes. Significant differences from adults in causes and patterns. *Arch Ped Adol Med* 1995;149(1):15-8.
73. Ruiz-Cotorro A, Balius-Matas R, Estruch-Massana A et al. Spondylolysis in young tennis players. *Br J Sports Med* 2006;40(5):441-6.
74. Ciullo JV, Jackson DW. Pars interarticularis stress reaction, spondylolysis and spondylolisthesis in gymnasts. *Clin Sports Med* 1985;4:95-110.
75. Fellander-Tsai L, Micheli L. Treatment of spondylolysis with external electrical stimulation and bracing in adolescent athletes: a report of two cases. *Clin J Sport Med* 1998;8(3):232-4.
76. Gregory P, Batt M, Kerslake R. Comparing spondylolysis in cricketers and soccer players. *Br J Sports Med* 2004;38(6):737-42.
77. Loud K, Micheli L, Bristol S et al. Family history predicts stress fracture in active female adolescents. *Pediatrics* 2007;120(2):364-72.
78. Rossi F. Spondylolysis, spondylolisthesis and sports. *J Sports Med Phys Fitness* 1988;18(4):317-40.
79. Schmitt H, Brocai D, Carstens C. Long-term review of the lumbar spine in javelin throwers. *J Bone Joint Surg Br* 2001;83(3):324-7.
80. Elliott B, Hardcastle P, Burnett A et al. The influence of fast bowling and physical factors on radiological features in high performance young fast bowlers. *J Sports Med Train Rehabil* 1992;3(2):113-30.
81. Foster D, John D, Elliott B et al. Back injuries to fast bowlers in cricket: a prospective study. *Br J Sports Med* 1989;23(3):150-4.
82. Engstrom C, Walker D. Pars interarticularis stress lesions in the lumbar spine of cricket fast bowlers. *Med Sci Sports Exerc* 2007;39(1):28-33.
83. Ranson C, Burnett A, Kerslake R. Injuries to the lower back in elite fast bowlers. *J Bone Joint Surg Br* 2010;92:1664-8.
84. Kwon Y-H, Shim J, Shim J et al., eds. Acute lumbar stress injury, trunk kinematics, lumbar MRI and paraspinal morphology in cricket fast bowlers in cricket. 26 *International Conference on Biomechanics in Sport*; 2008; Seoul, Korea.
85. Millson H, Gray J, Stretch R et al. Dissociation between back pain and bone stress reaction as measured by CT scan in young cricket fast bowlers. *Br J Sports Med* 2004;38:586-91.
86. Sys J, Michielsens J, Bracke P et al. Nonoperative treatment of active spondylolysis in elite athletes with normal X-ray findings: literature review and results of conservative treatment. *Eur Spine J* 2001;10(6):498-504.
87. Kountouris A, Portus M, Cook J. Quadratus lumborum asymmetry and lumbar spine injury in adolescent cricket fast bowlers. *J Sci Med Sport* 2012;15:393-7.
88. Kurd MF, Patel D, Norton R et al. Nonoperative treatment of symptomatic spondylolysis. *J Spinal Dis Techniques* 2007;20(8):560-4.
89. d'Hemecourt P, Zurakowski D, Kriemler S et al. Spondylolysis: returning the athlete to sports participation with brace treatment. *Orthopedics* 2002;25(6):653-7.
90. Wu SS, Lee CH, Chen PQ. Operative repair of symptomatic spondylolysis following a positive response to diagnostic pars injection. *J Spinal Dis* 1999;12(1):10-6.
91. Sairyo K, Sakai T, Mase Y et al. Painful lumbar spondylolysis among pediatric sports players: a pilot MRI study. *Arch Orth Trauma Surg* 2011;131(11):1485-9.
92. Vrable A, Sherman AL. Elite male adolescent gymnast who achieved union of a persistent bilateral pars defect. *Am J Phys Med Rehabil* 2009;88(2):156-60.
93. Masci L, Pike J, Malara F et al. Use of the one-legged hyperextension test and magnetic resonance imaging in the diagnosis of active spondylolysis. *Br J Sports Med* 2006;40:940-6.
94. Crewe H, Elliott B, Couanis G et al. The lumbar spine of the young cricket fast bowler: an MRI study. *J Sci Med Sport* 2012;15(3):190-4.
95. Hollenberg GM, Beattie PF, Meyers SP et al. Stress reactions of the lumbar pars interarticularis: the development of a new MRI classification system. *Spine* 2002;27(2):181-6.
96. Campbell R, Grainger A, Hide I et al. Juvenile spondylolysis: a comparative analysis of CT, SPECT and MRI. *Skel Radiol* 2005;34:63-73.
97. Read MT. Single photon emission computed tomography (SPECT) scanning for adolescent back pain. A sine qua non? *Br J Sports Med* 1994;28(1):56-7.
98. van der Wall H, Storey G, Magnussen J et al. Distinguishing scintigraphic features of spondylolysis. *J Ped Orthop* 2002;22(3):308-11.

99. Zukotynski K, Curtis C, Grant FD et al. The value of SPECT in the detection of stress injury to the pars interarticularis in patients with low back pain. *J Orthop Surg Res* 2010;5:13.
100. Sakai T, Sairyo K, Mima S et al. Significance of magnetic resonance imaging signal change in the pedicle in the management of pediatric lumbar spondylolysis. *Spine* 2010;35(14):641-5.
101. Morita T, Ikata T, Katoh S et al. Lumbar spondylolysis in children and adolescents. *J Bone Joint Surg Br* 1995;77-B(4): 620-5.
102. Sairyo K, Sakai T, Yasui N. Conservative treatment of lumbar spondylolysis in childhood and adolescence: the radiological signs which predict healing. *J Bone Joint Surg Br* 2009;91(2): 206-9.
103. Hardcastle P, Annear P, Foster DH, et al. Spinal abnormalities in young fast bowlers. *J Bone Joint Surg Br* 1992;74:421-5.
104. Sakai T, Sairyo K, Suzue N et al. Incidence and etiology of lumbar spondylolysis: review of the literature. *J Orthop Sci* 2010;15(3):281-8.
105. Ward CV, Latimer B. Human evolution and the development of spondylolysis. *Spine* 2005;30(16):1808-14.
106. Glazier PS. Is the 'crunch factor' an important consideration in the aetiology of lumbar spine pathology in cricket fast bowlers? *Sports Med* 2010;40(10):809-15.
107. de Visser H, Adam C, Crozier S et al. The role of quadratus lumborum asymmetry in the occurrence of lesions in the lumbar vertebrae of cricket fast bowlers. *Med Eng Physics* 2007;29:877-85.
108. Cyron BM, Hutton WC. The fatigue strength of the lumbar neural arch in spondylolysis. *J Bone Joint Surg Br* 1978;60-B(2):234-8.
109. Portus M, Mason B, Elliot B et al. Technique factors related to ball release speed and trunk injuries in high performance cricket fast bowlers. *Sports Biomech* 2004;3(2):263-84.
110. Cyron BM, Hutton WC. Variations in the amount and distribution of cortical bone across the partes interarticulares of L5. A predisposing factor in spondylolysis? *Spine* 1979;4(2):163-7.
111. Chung S-B, Lee S, Kim H et al. Significance of interfacet distance, facet joint orientation, and lumbar lordosis in spondylolysis. *Clin Anat* 2012;25(3):391-7.
112. Rankine JJ, Dickson RA. Unilateral spondylolysis and the presence of facet joint tropism. *Spine* 2010;35(21):E1111-4.
113. Bugg WG, Lewis M, Juetta A et al. Lumbar lordosis and pars interarticularis fractures: a case-control study. *Skel Radiol* 2012;41(7):817-22.
114. Bajwa NS, Toy JO, Ahn NU. L5 pedicle length is increased in subjects with spondylolysis: an anatomic study of 1072 cadavers. *Clin Orthop Relat Res* 2012;470(11):3202-6.
115. Miyauchi A, Baba I, Sumida T et al. Relationship between the histological findings of spondylolytic tissue, instability of the loose lamina, and low back pain. *Spine* 2008;33(6): 687-93.
116. Blanda J, Bethem D, Moats W et al. Defects of pars interarticularis in athletes: a protocol for nonoperative treatment. *J Spinal Dis* 1993;6(5):406-11.
117. Micheli L, Hall J, Miller M. Use of modified Boston brace for back injuries in athletes. *Am J Sports Med* 1980;8(5):351-6.
118. Sairyo K, Katoh S, Takata Y et al. MRI signal changes of the pedicle as an indicator for early diagnosis of spondylolysis in children and adolescents: a clinical and biomechanical study. *Spine* 2006;31(2):206-11.
119. Trainor TJ, Wiesel SW. Epidemiology of back pain in the athlete. *Clin Sports Med* 2002;21(1):93-103.
120. Orchard J, James T, Portus M. Injuries to elite male cricketers in Australia over a 10-year period. *J Sci Med Sport* 2006;9: 459-67.
121. Sairyo K, Sakai T, Yasui N et al. Conservative treatment for pediatric lumbar spondylolysis to achieve bone healing using a hard brace: what type and how long? *J Neurosurg Spine* 2012;16(6):610-4.
122. Debnath U, Freeman B, Grevitt M et al. Clinical outcome of symptomatic unilateral stress injuries of the lumbar pars interarticularis. *Spine* 2007;32(9):995-1000.
123. Alvarez-Diaz P, Alentorn-Geli E, Steinbacher G et al. Conservative treatment of lumbar spondylolysis in young soccer players. *Knee Surg Sports Traumatol Arthrosc* 2011;19(12):2111-4.
124. Congeni J, McCulloch J, Swanson K. Lumbar spondylolysis: a study of natural progression in athletes. *Am J Sports Med* 1997;25(2):248-53.
125. El Rassi G, Takemitsu M, Woratanarat P et al. Lumbar spondylolysis in pediatric and adolescent soccer players. *Am J Sports Med* 2005;33(11):1688-93.
126. Bayne H, Elliott B, Campbell A et al. Lumbar load in adolescent fast bowlers: a prospective injury study. *J Sci Med Sport* 2016; 19(2):117-22.
127. Khan K, Scott A. Mechanotherapy: how physical therapists' prescription of exercise promotes tissue repair. *Br J Sports Med* 2009;43:247-52.
128. Hulin B, Gabbett T, Blanch P et al. Spikes in acute workload are more predictive of injury than absolute workload in Australian cricket fast bowlers. *Br J Sports Med* 2014;48(8):708-12.
129. Rogalski B, Dawson B, Heasman J et al. Training and game loads and injury in elite Australian footballers. *J Sci Med Sport* 2013; 16(6):499-503.
130. Kountouris A, Cook J. Rehabilitation of Achilles and patellar tendinopathies. *Best Prac Clin Rheumatol* 2007;21(2):295-316.
131. Andrade N, Ashton C, Wray N et al. Systematic review of observational studies reveals no association between low back pain and lumbar spondylolysis with or without isthmic spondylolisthesis. *Eur Spine J* 2015;24:1289-95.
132. O'Sullivan PB, Twomey L, Allison G. Evaluation of specific stabilising exercise in the treatment of chronic low back pain with radiological diagnosis of spondylolysis and spondylolisthesis. *Spine* 1997;22(24):2959-67.

133. Tsirikos AI, Garrido EG. Spondylolysis and spondylolisthesis in children and adolescents. *J Bone Joint Surg Br* 2010;92(6): 751-9.
134. Benson R, Tavares S, Robertson S et al. Conservatively treated massive prolapsed discs: a 7-year follow-up. *Ann R Coll Surg Engl* 2010;92:147-53.
135. Petron DJ, Prideaux CC, Lickness L. Interventional spine procedures in athletes. *Curr Sports Med Rep* 2012;11(6):335-40.
136. Manchikanti L, Buenaventura RM, Manchikanti KN et al. Effectiveness of therapeutic lumbar transforaminal epidural steroid injections in managing lumbar spinal pain. *Pain Physician* 2012;15(3):E199-245.
137. Albert H, Sorensen J, Christensen B et al. Antibiotic treatment in patients with chronic low back pain and vertebral bone edema (Modic type 1 changes): a double-blind randomized clinical controlled trial of efficacy. *Eur Spine J* 2013;22(4): 697-707.
138. Vibe-Fersum K, O'Sullivan P, Kvale A et al. Efficacy of classification based cognitive functional therapy in patients with non-specific chronic low back pain: a randomized controlled trial. *Eur Pain J* 2013;17(6):916-28.
139. Deyo RA, Diehl AK, Rosenthal M. How many days of bed rest for acute low back pain? A randomized clinical trial. *N Engl J Med* 1986;315(17):1064-70.
140. Kuijpers T, van Middelkoop M, Rubinstein SM et al. A systematic review on the effectiveness of pharmacological interventions for chronic non-specific low-back pain. *Eur Spine J* 2011;20(1):40-50.
141. Chou R, Qaseem A, Snow V et al. Diagnosis and treatment of low back pain: a joint clinical practice guideline from the American College of Physicians and the American Pain Society. *Ann Int Med* 2007;147(7):478-91.
142. Lord SM, Barnsley L, Bogduk N. The utility of comparative local anesthetic blocks versus placebo-controlled blocks for the diagnosis of cervical zygapophysial joint pain. *Clin J Pain* 1995;11(3):208-13.
143. Dreyfuss P, Halbrook B, Pauza K et al. Efficacy and validity of radiofrequency neurotomy for chronic lumbar zygapophysial joint pain. *Spine* 2000;25(10):1270-7.
144. Carragee EJ, Lincoln T, Parmar VS et al. A gold standard evaluation of the 'discogenic pain' diagnosis as determined by provocative discography. *Spine* 2006;31(18):2115-23.
145. Carragee EJ, Don AS, Hurwitz EL et al. 2009 ISSLS Prize Winner: Does discography cause accelerated progression of degeneration changes in the lumbar disc: a ten-year matched cohort study. *Spine (Phila Pa 1976)* 2009;34(21):2338-45.
146. Willems PC. Provocative diskography: safety and predictive value in the outcome of spinal fusion or pain intervention for chronic low-back pain. *J Pain Res* 2014;7:699-705.
147. Yin W, Willard F, Carreiro J et al. Sensory stimulation-guided sacroiliac joint radiofrequency neurotomy: technique based on neuroanatomy of the dorsal sacral plexus. *Spine* 2003;28(20):2419-25.
148. Hill J, Whitehurst D, Lewis M et al. Comparison of stratified primary care management for low back pain with current best practice (STarT Back): a randomised controlled trial. *Lancet* 2011;Oct 29(378):1560-71.
149. O'Sullivan P. Lumbar segmental 'instability': clinical presentation and specific stabilizing exercise management. *Man Ther* 2000;5(1):2-12.
150. Dankaerts W, O'Sullivan P, Burnett A et al. The inter-examiner reliability of a classification method for non specific chronic low back pain patients with motor control impairment. *Man Ther* 2006;1:28-39.
151. Fersum K, Kvale A, O'Sullivan P et al. Interexaminer reliability of a classification system for patients with non-specific low back pain. *Man Ther* 2009;14:555-61.
152. Dankaerts W, O'Sullivan P. The validity of O'Sullivan's classification system (CS) for a sub-group of NS-CLBP with motor control impairment (MCI): overview of a series of studies and review of the literature. *Man Ther* 2011;16:9-14.
153. Sullivan M, Thibault P, Andrikonyte J et al. Psychological influences on repetition-induced summation of activity-related pain in patients with chronic low back pain. *Pain* 2009;141:70-8.
154. Rabey M, Beales D, Slater H et al. Multidimensional pain profiles in four cases of chronic non-specific axial low back pain: an examination of the limitations of contemporary classification systems. *Man Ther* 2015;20(1):138-47.
155. O'Sullivan K, Dankaerts W, O'Sullivan L et al. Cognitive functional therapy for disabling nonspecific chronic low back pain: multiple case-cohort study. *Phys Ther* 2015;95(11):1478-88.
156. Smith BE, Littlewood C, May S. An update of stabilisation exercises for low back pain: a systematic review with meta-analysis. *BMC Musculoskelet Disord* 2014;15:416.

CHAPTER 30

1. Hausswirth C, Louis J, Aubry A et al. Evidence of disturbed sleep and increased illness in overreached endurance athletes. *Med Sci Sports Exerc* 2014;46(5):1036-45.
2. Hannibal KE, Bishop MD. Chronic stress, cortisol dysfunction and pain: a psychoneuroendocrine rationale for stress management in pain rehabilitation. *Phys Ther* 2014;94(12):1816-25.
3. van der Does HT, Brink MS, Otter RT et al. Injury risk is increased by changes in perceived recovery of team sport players. *Clin J Sport Med* 2016. Mar 3. [Epub ahead of print]
4. Otter RT, Brink MS, Diercks RL et al. A negative life event impairs psychosocial stress, recovery and running economy of runners. *Int J Sports Med* 2016;37(3):224-9.
5. Lentz TA, Zeppieri G, Jr., George SZ et al. Comparison of physical impairment, functional and psychosocial measures based on fear of reinjury/lack of confidence and return-to-sport status after ACL reconstruction. *Am J Sports Med* 2015;43(2):345-53.
6. Tjong VK, Devitt BM, Murnaghan ML et al. A qualitative investigation of return to sport after arthroscopic bankart repair: beyond stability. *Am J Sports Med* 2015;43(8):2005-11.

7. van der Wurff P, Buijs EJ, Groen GJ. Intensity mapping of pain referral areas in sacroiliac joint pain patients. *J Manipulative Physiol Ther* 2006;29(3):190-5.
8. Donelson R, Aprill C, Medcalf R et al. A prospective study of centralization of lumbar and referred pain. A predictor of symptomatic discs and anular competence. *Spine (Phila Pa 1976)* 1997;22(10):1115-22.
9. Laslett M, Aprill CN, McDonald B et al. Diagnosis of sacroiliac joint pain: validity of individual provocation tests and composites of tests. *Man Ther* 2005;10(3):207-18.
10. Murrie VL, Dixon AK, Hollingworth W et al. Lumbar lordosis: study of patients with and without low back pain. *Clin Anat* 2003;16(2):144-7.
11. Preece SJ, Willan P, Nester CJ et al. Variation in pelvic morphology may prevent the identification of anterior pelvic tilt. *J Man Manip Ther* 2008;16(2):113-7.
12. Herrington L. Assessment of the degree of pelvic tilt within a normal asymptomatic population. *Man Ther* 2011;16(6):646-8.
13. Tsai L, Wredmark T. Spinal posture, sagittal mobility and subjective rating of back problems in former female elite gymnasts. *Spine (Phila Pa 1976)* 1993;18(7):872-5.
14. Tuzun C, Yorulmaz I, Cindas A et al. Low back pain and posture. *Clin Rheumatol* 1999;18(4):308-12.
15. Sarikaya S, Ozdolap S, Gumustass S et al. Low back pain and lumbar angles in Turkish coal miners. *Am J Ind Med* 2007;50(2):92-6.
16. Christensen ST, Hartvigsen J. Spinal curves and health: a systematic critical review of the epidemiological literature dealing with associations between sagittal spinal curves and health. *J Manipulative Physiol Ther* 2008;31(9):690-714.
17. Laird RA, Gilbert J, Kent P et al. Comparing lumbo-pelvic kinematics in people with and without back pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord* 2014;15:229.
18. Kibsgard TJ, Roise O, Sturesson B et al. Radiostereometric analysis of movement in the sacroiliac joint during a single-leg stance in patients with long-lasting pelvic girdle pain. *Clin Biomech (Bristol, Avon)* 2014;29(4):406-11.
19. Riddle DL, Freburger JK. Evaluation of the presence of sacroiliac joint region dysfunction using a combination of tests: a multicenter intertester reliability study. *Phys Ther* 2002;82(8):772-81.
20. Robinson HS, Brox JJ, Robinson R et al. The reliability of selected motion- and pain provocation tests for the sacroiliac joint. *Man Ther* 2007;12(1):72-9.
21. Harms MC, Bader DL. Variability of forces applied by experienced therapists during spinal mobilisation. *Clin Biomech (Bristol, Avon)* 1997;12(6):393-9.
22. Snodgrass SJ, Rivett DA, Robertson VJ. Manual forces applied during posterior-to-anterior spinal mobilisation: a review of the evidence. *J Manipulative Physiol Ther* 2006;29(4):316-29.
23. Koppenhaver SL, Hebert JJ, Kawchuk GN et al. Criterion validity of manual assessment of spinal stiffness. *Man Ther* 2014;19(6):589-94.
24. Scaia V, Baxter D, Cook C. The pain provocation-based straight leg raise test for diagnosis of lumbar disc herniation, lumbar radiculopathy, and/or sciatica: a systematic review of clinical utility. *J Back Musculoskelet Rehabil* 2012;25(4):215-23.
25. Cacchio A, De Paulis F, Maffulli N. Development and validation of a new visa questionnaire (VISA-H) for patients with proximal hamstring tendinopathy. *Br J Sports Med* 2014;48(6):448-52.
26. Alyas F, Turner M, Connell D. MRI findings in the lumbar spines of asymptomatic, adolescent, elite tennis players. *Br J Sports Med* 2007;41(11):836-41; discussion 841.
27. Campbell J, Colvin LA. Management of low back pain. *BMJ* 2013;347:f3148.
28. Ekizos A, Papatzika F, Charcharis G et al. Ultrasound does not provide reliable results for the measurement of the patellar tendon cross sectional area. *J Electromyogr Kinesiol* 2013;23(6):1278-82.
29. Trainor TJ, Trainor MA. Aetiology of low back pain in athletes. *Curr Sports Med Rep* 2004;3(1):41-6.
30. Simons DG, Travell J. Myofascial trigger points, a possible explanation. *Pain* 1981;10(1):106-9.
31. Hardcastle P, Nade S. The significance of the Trendelenburg test. *J Bone Joint Surg Br* 1985;67(5):741-6.
32. Grimaldi A. Assessing lateral stability of the hip and pelvis. *Man Ther* 2011;16(1):26-32.
33. Arnold CM, Warkentin KD, Chilibeck PD et al. The reliability and validity of handheld dynamometry for the measurement of lower-extremity muscle strength in older adults. *J Strength Cond Res* 2010;24(3):815-24.
34. Thorborg K, Bandholm T, Schick M et al. Hip strength assessment using handheld dynamometry is subject to intertester bias when testers are of different sex and strength. *Scand J Med Sci Sports* 2013;23(4):487-93.
35. Widler KS, Glatthorn JF, Bizzini M et al. Assessment of hip abductor muscle strength. A validity and reliability study. *J Bone Joint Surg Am* 2009;91(11):2666-72.
36. Selkowitz DM, Beneck GJ, Powers CM. Which exercises target the gluteal muscles while minimising activation of the tensor fascia lata? Electromyographic assessment using fine-wire electrodes. *J Orthop Sports Phys Ther* 2013;43(2):54-64.
37. MacAskill MJ, Durant TJ, Wallace DA. Gluteal muscle activity during weightbearing and non-weightbearing exercise. *Int J Sports Phys Ther* 2014;9(7):907-14.
38. Reiman MP, Bolgia LA, Loudon JK. A literature review of studies evaluating gluteus maximus and gluteus medius activation during rehabilitation exercises. *Physiother Theory Pract* 2012;28(4):257-68.
39. Huang Q, Zhang X, Wang J. The diagnoses and treatments of myofascial trigger point pains company with referred pains of leg. *Chinese J Rehab Med* 2005;3:193-5.
40. Schwarzer AC, Derby R, Aprill CN et al. Pain from the lumbar zygapophysial joints: a test of two models. *J Spinal Disord* 1994;7(4):331-6.
41. Laslett M, Aprill CN, McDonald B et al. Clinical predictors of lumbar provocation discography: a study of clinical predictors of

- lumbar provocation discography. *Eur Spine J* 2006;15(10):1473-84.
42. Majlesi J, Togay H, Unalan H et al. The sensitivity and specificity of the slump and the straight leg raising tests in patients with lumbar disc herniation. *J Clin Rheumatol* 2008;14(2):87-91.
 43. Maffulli N, Khan KM, Puddu G. Overuse tendon conditions: time to change a confusing terminology. *Arthroscopy* 1998;14(8):840-3.
 44. Lee HH, Lin CW, Wu HW et al. Changes in biomechanics and muscle activation in injured ballet dancers during a jump-land task with turnout (Sissonne Fermee). *J Sports Sci* 2012;30(7):689-97.
 45. Puranen J, Orava S. The hamstring syndrome. A new diagnosis of gluteal sciatic pain. *Am J Sports Med* 1988;16(5):517-21.
 46. Fredericson M, Moore W, Guillet M et al. High hamstring tendinopathy in runners: meeting the challenges of diagnosis, treatment and rehabilitation. *Phys Sportsmed* 2005;33(5):32-43.
 47. Cook JL, Purdam C. Is compressive load a factor in the development of tendinopathy? *Br J Sports Med* 2012;46(3):163-8.
 48. Docking S, Samiric T, Scase E et al. Relationship between compressive loading and ECM changes in tendons. *Muscles Ligaments Tendons J* 2013;3(1):7-11.
 49. Young IJ, van Riet RP, Bell SN. Surgical release for proximal hamstring syndrome. *Am J Sports Med* 2008;36(12):2372-8.
 50. Cacchio A, Borra F, Severini G et al. Reliability and validity of three pain provocation tests used for the diagnosis of chronic proximal hamstring tendinopathy. *Br J Sports Med* 2012;46(12):883-7.
 51. Koulouris G, Connell D. Hamstring muscle complex: an imaging review. *Radiographics* 2005;25(3):571-86.
 52. White KE. High hamstring tendinopathy in 3 female long distance runners. *J Chiropr Med* 2011;10(2):93-9.
 53. Jayaseelan DJ, Moats N, Ricardo CR. Rehabilitation of proximal hamstring tendinopathy utilising eccentric training, lumbopelvic stabilisation and trigger point dry needling: 2 case reports. *J Orthop Sports Phys Ther* 2014;44(3):198-205.
 54. Threlkeld AJ. The effects of manual therapy on connective tissue. *Phys Ther* 1992;72(12):893-902.
 55. Malliaras P, Barton CJ, Reeves ND et al. Achilles and patellar tendinopathy loading programmes: a systematic review comparing clinical outcomes and identifying potential mechanisms for effectiveness. *Sports Med* 2013;43(4):267-86.
 56. Naugle KM, Naugle KE, Fillingim RB et al. Isometric exercise as a test of pain modulation: effects of experimental pain test, psychological variables and sex. *Pain Med* 2014;15(4):692-701.
 57. Misra G, Paris TA, Archer DB et al. Dose-response effect of isometric force production on the perception of pain. *PLoS One* 2014;9(2):e88105.
 58. Khan KM, Scott A. Mechanotherapy: how physical therapists' prescription of exercise promotes tissue repair. *Br J Sports Med* 2009;43(4):247-52.
 59. Thomee R. A comprehensive treatment approach for patellofemoral pain syndrome in young women. *Phys Ther* 1997;77(12):1690-703.
 60. Kongsgaard M, Qvortrup K, Larsen J et al. Fibril morphology and tendon mechanical properties in patellar tendinopathy: effects of heavy slow resistance training. *Am J Sports Med* 2010;38(4):749-56.
 61. Silbernagel KG, Thomee R, Eriksson BI et al. Continued sports activity, using a pain-monitoring model, during rehabilitation in patients with Achilles tendinopathy: a randomised controlled study. *Am J Sports Med* 2007;35(6):897-906.
 62. Cohen SP, Chen Y, Neufeld NJ. Sacroiliac joint pain: a comprehensive review of epidemiology, diagnosis and treatment. *Expert Rev Neurother* 2013;13(1):99-116.
 63. Timm KE. Sacroiliac joint dysfunction in elite rowers. *J Orthop Sports Phys Ther* 1999;29(5):288-93.
 64. Brolinson PG, Kozar AJ, Cibor G. Sacroiliac joint dysfunction in athletes. *Curr Sports Med Rep* 2003;2(1):47-56.
 65. Berthelot JM, Labat JJ, Le Goff B et al. Provocative sacroiliac joint manoeuvres and sacroiliac joint block are unreliable for diagnosing sacroiliac joint pain. *Joint Bone Spine* 2006;73(1):17-23.
 66. Nourbakhsh MR, Arab AM. Relationship between mechanical factors and incidence of low back pain. *J Orthop Sports Phys Ther* 2002;32(9):447-60.
 67. Richardson CA, Snijders CJ, Hides JA et al. The relation between the transversus abdominis muscles, sacroiliac joint mechanics and low back pain. *Spine (Phila Pa 1976)* 2002;27(4):399-405.
 68. van Wingerden JP, Vleeming A, Buyruk HM et al. Stabilisation of the sacroiliac joint in vivo: verification of muscular contribution to force closure of the pelvis. *Eur Spine J* 2004;13(3):199-205.
 69. Damen L, Buyruk HM, Guler-Uysal F et al. Pelvic pain during pregnancy is associated with asymmetric laxity of the sacroiliac joints. *Acta Obstet Gynecol Scand* 2001;80(11):1019-24.
 70. Vleeming A, Schuenke MD, Masi AT et al. The sacroiliac joint: an overview of its anatomy, function and potential clinical implications. *J Anat* 2012;221(6):537-67.
 71. Goode A, Hegedus EJ, Sizer P et al. Three-dimensional movements of the sacroiliac joint: a systematic review of the literature and assessment of clinical utility. *J Man Manip Ther* 2008;16(1):25-38.
 72. Garras DN, Carothers JT, Olson SA. Single-leg-stance (flamingo) radiographs to assess pelvic instability: how much motion is normal? *J Bone Joint Surg Am* 2008;90(10):2114-8.
 73. Lavignolle B, Vital JM, Senegas J et al. An approach to the functional anatomy of the sacroiliac joints in vivo. *Anat Clin* 1983;5(3):169-76.
 74. Vleeming A, Snijders CJ, Stoeckart R et al. The role of the sacroiliac joints in coupling between spine, pelvis, legs and arms. In: Vleeming A, Mooney V, Dorman T et al., eds. *Movement, Stability and Low Back Pain: The Essential Role of the Pelvis*. New York: Churchill Livingstone, 1997:53-71.
 75. van der Wurff P, Buijs EJ, Groen GJ. A multitest regimen of pain provocation tests as an aid to reduce unnecessary minimally invasive sacroiliac joint procedures. *Arch Phys Med Rehabil* 2006;87(1):10-4.

76. Freburger JK, Riddle DL. Using published evidence to guide the examination of the sacroiliac joint region. *Phys Ther* 2001;81(5):1135-43.
77. O'Sullivan PB, Beales DJ, Beetham JA et al. Altered motor control strategies in subjects with sacroiliac joint pain during the active straight-leg-raise test. *Spine (Phila Pa 1976)* 2002;27(1):E1-8.
78. Mens JM, Vleeming A, Snijders CJ et al. Validity of the active straight leg raise test for measuring disease severity in patients with posterior pelvic pain after pregnancy. *Spine (Phila Pa 1976)* 2002;27(2):196-200.
79. Mens JM, Vleeming A, Snijders CJ et al. The active straight leg raising test and mobility of the pelvic joints. *Eur Spine J* 1999;8(6):468-73.
80. Pålsson TS, Hirata RP, Graven-Nielsen T. Experimental pelvic pain impairs the performance during the active straight leg raise test and causes excessive muscle stabilisation. *Clin J Pain* 2015;31(7):642-51.
81. Shearar KA, Colloca CJ, White HL. A randomised clinical trial of manual versus mechanical force manipulation in the treatment of sacroiliac joint syndrome. *J Manipulative Physiol Ther* 2005;28(7):493-501.
82. Tullberg T, Blomberg S, Branth B et al. Manipulation does not alter the position of the sacroiliac joint. A roentgen stereophotogrammetric analysis. *Spine (Phila Pa 1976)* 1998;23(10):1124-8; discussion 1129.
83. Bialosky JE, Bishop MD, Price DD et al. The mechanisms of manual therapy in the treatment of musculoskeletal pain: a comprehensive model. *Man Ther* 2009;14(5):531-8.
84. Mens JM, Damen L, Snijders CJ et al. The mechanical effect of a pelvic belt in patients with pregnancy-related pelvic pain. *Clin Biomech (Bristol, Avon)* 2006;21(2):122-7.
85. Pel JJ, Spoor CW, Goossens RH et al. Biomechanical model study of pelvic belt influence on muscle and ligament forces. *J Biomech* 2008;41(9):1878-84.
86. Damen L, Spoor CW, Snijders CJ et al. Does a pelvic belt influence sacroiliac joint laxity? *Clin Biomech (Bristol, Avon)* 2002;17(7):495-8.
87. Kim WM, Lee HG, Jeong CW et al. A randomised controlled trial of intra-articular prolotherapy versus steroid injection for sacroiliac joint pain. *J Altern Complement Med* 2010;16(12):1285-90.
88. Cusi M, Saunders J, Hungerford B et al. The use of prolotherapy in the sacroiliac joint. *Br J Sports Med* 2010;44(2):100-4.
89. Robinson DR. Piriformis syndrome in relation to sciatic pain. *Am J Surg* 1947;73(3):355-8.
90. Smoll NR. Variations of the piriformis and sciatic nerve with clinical consequence: a review. *Clin Anat* 2010;23(1):8-17.
91. Papadopoulos EC, Khan SN. Piriformis syndrome and low back pain: a new classification and review of the literature. *Orthop Clin North Am* 2004;35(1):65-71.
92. Stewart JD. The piriformis syndrome is overdiagnosed. *Muscle Nerve* 2003;28(5):644-6.
93. Fishman LM, Schaefer MP. The piriformis syndrome is underdiagnosed. *Muscle Nerve* 2003;28(5):646-9.
94. Martin HD, Kivlan BR, Palmer JJ et al. Diagnostic accuracy of clinical tests for sciatic nerve entrapment in the gluteal region. *Knee Surg Sports Traumatol Arthrosc* 2014;22(4):882-8.
95. Miller TA, White KP, Ross DC. The diagnosis and management of piriformis syndrome: myths and facts. *Can J Neurol Sci* 2012;39(5):577-83.
96. Misirlioglu TO, Akgun K, Palamar D et al. Piriformis syndrome: comparison of the effectiveness of local anaesthetic and corticosteroid injections: a double-blinded, randomised controlled study. *Pain Physician* 2015;18(2):163-71.
97. Benson ER, Schutzer SF. Posttraumatic piriformis syndrome: diagnosis and results of operative treatment. *J Bone Joint Surg Am* 1999;81(7):941-9.
98. Dezawa A, Kusano S, Miki H. Arthroscopic release of the piriformis muscle under local anaesthesia for piriformis syndrome. *Arthroscopy* 2003;19(5):554-7.
99. Khodair SA, Ghieda UE, Elsayed AS. Ischiofemoral impingement syndrome: spectrum of MRI findings in comparison to normal subjects. *Egypt J Radiol Nuc Med* 2014;45(3):819-24.
100. Tosun O, Algin O, Yalcin N et al. Ischiofemoral impingement: evaluation with new MRI parameters and assessment of their reliability. *Skeletal Radiol* 2012;41(5):575-87.
101. Stafford GH, Villar RN. Ischiofemoral impingement. *J Bone Joint Surg Br* 2011;93(10):1300-2.
102. Ali AM, Whitwell D, Ostlere SJ. Case report: imaging and surgical treatment of a snapping hip due to ischiofemoral impingement. *Skeletal Radiol* 2011;40(5):653-6.
103. Torriani M, Souto SC, Thomas BJ et al. Ischiofemoral impingement syndrome: an entity with hip pain and abnormalities of the quadratus femoris muscle. *AJR Am J Roentgenol* 2009;193(1):186-90.
104. Ali AM, Teh J, Whitwell D et al. Ischiofemoral impingement: a retrospective analysis of cases in a specialist orthopaedic centre over a four-year period. *Hip Int* 2013;23(3):263-8.
105. Matsen FA, 3rd. Compartmental syndrome. A unified concept. *Clin Orthop Relat Res* 1975;113:8-14.
106. Kwong Y, Patel J, Ramanathan EB. Spontaneous complete hamstring avulsion causing posterior thigh compartment syndrome. *Br J Sports Med* 2006;40(8):723-4; discussion 724.
107. Mallo GC, Stanat SJ, Al-Humadi M et al. Posterior thigh compartment syndrome as a result of a basketball injury. *Orthopaedics* 2009;32(12):923.
108. Hynes JE, Jackson A. Atraumatic gluteal compartment syndrome. *Postgrad Med J* 1994;70(821):210-2.
109. Whitesides TE, Haney TC, Morimoto K et al. Tissue pressure measurements as a determinant for the need of fasciotomy. *Clin Orthop Relat Res* 1975;113:43-51.
110. Raether PM, Lutter LD. Recurrent compartment syndrome in the posterior thigh. Report of a case. *Am J Sports Med* 1982;10(1):40-3.
111. Abebe ES, Moorman CT, Garrett WE. Proximal hamstring avulsion injuries: injury mechanism, diagnosis and disease course. *Op Tech Sports Med* 2009;17(4):205-9.

112. Sallay PI. Diagnosis, classification and management of acute proximal hamstring avulsion injuries. *Op Tech Sports Med* 2009;17(4):196-204.
113. Cohen S, Bradley J. Acute proximal hamstring rupture. *J Am Acad Orthop Surg* 2007;15(6):350-5.
114. Carmichael J, Packham I, Trikha SP et al. Avulsion of the proximal hamstring origin. Surgical technique. *J Bone Joint Surg Am* 2009;91 Suppl 2:249-56.
115. Cross M, Vandersluis R, Wood D et al. Surgical repair of chronic complete hamstring tendon rupture in the adult patient. *Am J Sports Med* 1998; 26(6):785-8.
116. Harris JD, Griersser MJ, Best TM et al. Treatment of proximal hamstring ruptures: a systematic review. *Int J Sports Med* 2011;32(7):490-5.
117. Fernbach SK, Wilkinson RH. Avulsion injuries of the pelvis and proximal femur. *AJR Am J Roentgenol* 1981;137(3):581-4.
118. Rossi F, Dragoni S. Acute avulsion fractures of the pelvis in adolescent competitive athletes: prevalence, location and sports distribution of 203 cases collected. *Skeletal Radiol* 2001;30(3):127-31.
119. Wootton JR, Cross MJ, Holt KW. Avulsion of the ischial apophysis. The case for open reduction and internal fixation. *J Bone Joint Surg Br* 1990;72(4):625-7.
120. Lawrence RC, Helmick CG, Arnett FC et al. Estimates of the prevalence of arthritis and selected musculoskeletal disorders in the United States. *Arthritis Rheum* 1998;41(5):778-99.
121. Dougados M, van der Heijde D. Ankylosing spondylitis: how should the disease be assessed? *Best Pract Res Clin Rheumatol* 2002;16(4):605-18.
122. van der Linden S, Valkenburg HA, Cats A. Evaluation of diagnostic criteria for ankylosing spondylitis. A proposal for modification of the New York criteria. *Arthritis Rheum* 1984;27(4):361-8.
123. Dagfinrud H, Kvien TK, Hagen KB. The Cochrane review of physiotherapy interventions for ankylosing spondylitis. *J Rheumatol* 2005;32(10):1899-906.
124. Hosey RG, Fernandez MM, Johnson DL. Evaluation and management of stress fractures of the pelvis and sacrum. *Orthopaedics* 2008;31(4):383-5.
125. Aretxabala I, Fraiz E, Perez-Ruiz F et al. Sacral insufficiency fractures. High association with pubic rami fractures. *Clin Rheumatol* 2000;19(5):399-401.
126. Tsiridis E, Upadhyay N, Giannoudis PV. Sacral insufficiency fractures: current concepts of management. *Osteoporos Int* 2006;17(12):1716-25.
127. Longhino V, Bonora C, Sansone V. The management of sacral stress fractures: current concepts. *Clin Cases Miner Bone Metab* 2011;8(3):19-23.
128. Lever M, Lever E, Lever EG. Rethinking osteoporotic sacral fractures. *Injury* 2009;40(4):466.
129. Dimmen S, Nordsletten L, Engebretsen L et al. Negative effect of parecoxib on bone mineral during fracture healing in rats. *Acta Orthop* 2008;79(3):438-44.

130. Knobloch K, Schreibleueller L, Jagodzinski M et al. Rapid rehabilitation programme following sacral stress fracture in a long-distance running female athlete. *Arch Orthop Trauma Surg* 2007;127(9):809-13.

CHAPTER 31

1. <http://www.afl.com.au/injury%20report/tabid/13706/default.aspx>. 2008 AFL Injury Report.
2. Singh PJ, O'Donnell JM. The outcome of hip arthroscopy in Australian football league players: a review of 27 hips. *Arthroscopy* 2010;26(6):743-9.
3. Khanna V, Caragianis A, Diprimio G et al. Incidence of hip pain in a prospective cohort of asymptomatic volunteers: is the cam deformity a risk factor for hip pain? *Am J Sports Med* 2014;42(4):793-7.
4. Burnett RS, Della Rocca GJ, Prather H et al. Clinical presentation of patients with tears of the acetabular labrum. *J Bone Joint Surg Am* 2006;88(7):1448-57.
5. Philippon MJ, Weiss DR, Kuppersmith DA et al. Arthroscopic labral repair and treatment of femoroacetabular impingement in professional hockey players. *Am J Sports Med* 2010;38(1):99-104.
6. Reichenbach S, Leunig M, Werlen S et al. Association between cam-type deformities and magnetic resonance imaging-detected structural hip damage: a cross-sectional study in young men. *Arthritis Rheum* 2011;63(12):4023-30.
7. Crawford MJ, Dy CJ, Alexander JW et al. The 2007 Frank Stinchfield Award. The biomechanics of the hip labrum and the stability of the hip. *Clin Orthop Relat Res* 2007;465:16-22.
8. Beck M, Kalhor M, Leunig M et al. Hip morphology influences the pattern of damage to the acetabular cartilage—femoroacetabular impingement as a cause of early osteoarthritis of the hip. *J Bone Joint Surg Br* 2005;87b(7):1012-18.
9. Domayer SE, Mamisch TC, Kress I et al. Radial dGEMRIC in developmental dysplasia of the hip and in femoroacetabular impingement: preliminary results. *Osteoarthritis Cartilage* 2010;18(11):1421-8.
10. Hapa O, Yuksel HY, Muratli HH et al. Axial plane coverage and torsion measurements in primary osteoarthritis of the hip with good frontal plane coverage and spherical femoral head. *Arch Orthop Trauma Surg* 2010;130(10):1305-10.
11. Ito K, Minka MA, 2nd, Leunig M et al. Femoroacetabular impingement and the cam-effect. A MRI-based quantitative anatomical study of the femoral head-neck offset. *J Bone Joint Surg Br* 2001;83(2):171-6.
12. McCarthy JMD, Noble PP, Aluisio FVMD et al. Anatomy, pathologic features and treatment of acetabular labral tears. [Report]. *Clin Orthop Relat Res* 2003;406(1):38-47.
13. Tanzer M, Noiseux N. Osseous abnormalities and early osteoarthritis: the role of hip impingement. *Clin Orthop Relat Res* 2004;429:170-7.
14. Kemp JL, Makdissi M, Pritchard MG et al. Chondropathy of the hip at arthroscopy: prevalence and relationship to labral

- pathology, femoro-acetabular impingement and patient-reported outcomes. *Br J Sports Med* 2014;48(11):1102-7.
15. Diamond LE, Dobson FL, Bennell KL et al. Physical impairments and activity limitations in people with femoroacetabular impingement: a systematic review. *Br J Sports Med* 2015;49(4):230-42.
 16. Kemp JL, Schache AG, Makdissi M et al. The relationship between patient-reported outcomes and physical impairments in people with early hip osteoarthritis following hip arthroscopy. *Osteoarthritis Cartilage* 2015 Jul 3. [Epub ahead of print]
 17. Stem ES, O'Connor MI, Kransdorf MJ et al. Computed tomography analysis of acetabular anteversion and abduction. *Skeletal Radiol* 2006;35(6):385-9.
 18. Lewis CL, Sahrman SA. Acetabular labral tears. *Phys Ther* 2006;86(1):110-21.
 19. Petersen W, Petersen F, Tillmann B. Structure and vascularization of the acetabular labrum with regard to the pathogenesis and healing of labral lesions. *Arch Orthop Trauma Surg* 2003;123(6):283-8.
 20. Alzahrani A, Bali K, Gudena R et al. The innervation of the human acetabular labrum and hip joint: an anatomic study. *BMC Musculoskelet Disord* 2014;15(1):41.
 21. Kim SJ, Choi NH, Kim HJ. Operative hip arthroscopy. *Clin Orthop Relat Res* 1998;353:156-65.
 22. Narvani AA, Tsiroidis E, Tai CC et al. Acetabular labrum and its tears. *Br J Sports Med* 2003;37(3):207-11.
 23. Dholander AA, Lambrecht S, Verdonk PC et al. First insights into human acetabular labrum cell metabolism. *Osteoarthritis Cartilage* 2012;20(7):670-7.
 24. Ferguson SJ, Bryant JT, Ganz R et al. The acetabular labrum seal: a poroelastic finite element model. *Clin Biomech (Bristol, Avon)* 2000;15(6):463-8.
 25. Konrath GA, Hamel AJ, Olson SA et al. The role of the acetabular labrum and the transverse acetabular ligament in load transmission in the hip. *J Bone Joint Surg Am* 1998;80(12):1781-8.
 26. Dwyer MK, Jones HL, Hogan MG et al. The acetabular labrum regulates fluid circulation of the hip joint during functional activities. *Am J Sports Med* 2014;42(4):812-9.
 27. Dwyer MK, Jones HL, Field RE et al. Femoroacetabular impingement negates the acetabular labral seal during pivoting manoeuvres but not gait. *Clin Orthop Relat Res* 2015;473(2):602-7.
 28. Bardakos NV, Villar RN. The ligamentum teres of the adult hip. *J Bone Joint Surg Br* 2009;91(1):8-15.
 29. Pollard TC, Gwilym SE, Carr AJ. The assessment of early osteoarthritis. *J Bone Joint Surg Br* 2008;90(4):411-21.
 30. Pritzker KP, Gay S, Jimenez SA et al. Osteoarthritis cartilage histopathology: grading and staging. *Osteoarthritis Cartilage* 2006;14(1):13-29.
 31. Norkin C, LeVange P. *Joint Structure and Function*. Philadelphia: F.A. Davis Company, 1983.
 32. Retchford T, Crossley KM, Grimaldi A et al. Can local muscles augment stability in the hip? A narrative literature review. *J Musculoskelet Neuronal Interact* 2013;13(1):1-12.
 33. Neumann DA. Kinesiology of the hip: a focus on muscular actions. *J Orthop Sports Phys Ther* 2010;40(2):82-94.
 34. Ward SR, Winters TM, Blemker SS. The architectural design of the gluteal muscle group: implications for movement and rehabilitation. *J Orthop Sports Phys Ther* 2010;40(2):95-102.
 35. Correa TA, Crossley KM, Kim HJ et al. Contributions of individual muscles to hip joint contact force in normal walking. *J Biomech* 2010;43(8):1618-22.
 36. Kemp JL, Schache AG, Makdissi M et al. Greater understanding of normal hip physical function may guide clinicians in providing targeted rehabilitation programmes. *J Sci Med Sport* 2013;16(4):292-6.
 37. Thorborg K, Bandholm T, Schick M et al. Hip strength assessment using handheld dynamometry is subject to intertester bias when testers are of different sex and strength. *Scand J Med Sci Sports* 2013;23(4):487-93.
 38. Kemp JL, Makdissi M, Schache AG et al. Hip chondropathy at arthroscopy: prevalence and relationship to labral pathology, femoroacetabular impingement and patient-reported outcomes. *Br J Sports Med* 2014;48(14):1102-7.
 39. Bradshaw CJ, Bundy M, Falvey E. The diagnosis of longstanding groin pain: a prospective clinical cohort study. *Br J Sports Med* 2008;42(10):851-4.
 40. Reiman MP, Goode AP, Hegedus EJ et al. Diagnostic accuracy of clinical tests of the hip: a systematic review with meta-analysis. *Br J Sports Med* 2013;47(14):893-902.
 41. Clohisy JC, Baca G, Beaulé PE et al. Descriptive epidemiology of femoroacetabular impingement: a North American cohort of patients undergoing surgery. *Am J Sports Med* 2013;41(6):1348-56.
 42. Weir A, de Vos RJ, Moen M et al. Prevalence of radiological signs of femoroacetabular impingement in patients presenting with long-standing adductor-related groin pain. *Br J Sports Med* 2011;45(1):6-9.
 43. Byrd JW, Jones KS. Hip arthroscopy in athletes. *Clin Sports Med* 2001;20(4):749-61.
 44. Narvani AA, Tsiroidis E, Kendall S et al. A preliminary report on prevalence of acetabular labrum tears in sports patients with groin pain. *Knee Surg Sports Traumatol Arthrosc* 2003;11(6):403-8.
 45. Maslowski E, Sullivan W, Forster Harwood J et al. The diagnostic validity of hip provocation manoeuvres to detect intra-articular hip pathology. *PM&R* 2010;2(3):174-81.
 46. McCarthy JC, Busconi B. The role of hip arthroscopy in the diagnosis and treatment of hip disease. *Can J Surg* 1995;38 Suppl 1:S13-7.
 47. Bird PA, Oakley SP, Shnier R et al. Prospective evaluation of magnetic resonance imaging and physical examination findings in patients with greater trochanteric pain syndrome. *Arthritis Rheum* 2001;44(9):2138-45.
 48. Lequesne M, Mathieu P, Vuillemin-Bodaghi V et al. Gluteal tendinopathy in refractory greater trochanter pain syndrome: diagnostic value of two clinical tests. *Arthritis Rheum* 2008;59(2):241-6.

49. Sutlive TG, Lopez HP, Schnitker DE et al. Development of a clinical prediction rule for diagnosing hip osteoarthritis in individuals with unilateral hip pain. *J Orthop Sports Phys Ther* 2008;38(9):542-50.
50. Pollard TC, Batra RN, Judge A et al. Genetic predisposition to the presence and 5-year clinical progression of hip osteoarthritis. *Osteoarthritis Cartilage* 2012;20(5):368-75.
51. Mitchell B, McCrory P, Brukner P et al. Hip joint pathology: clinical presentation and correlation between magnetic resonance arthrography, ultrasound and arthroscopic findings in 25 consecutive cases. *Clin J Sport Med* 2003;13(3):152-6.
52. Thorborg K, Holmich P, Christensen R et al. The Copenhagen Hip and Groin Outcome Score (HAGOS): development and validation according to the COSMIN checklist. *Br J Sports Med* 2011;45(6):478-91.
53. Nilsson AK, Lohmander LS, Klassbo M et al. Hip disability and osteoarthritis outcome score (HOOS)—validity and responsiveness in total hip replacement. *BMC Musculoskelet Disord* 2003;4:10.
54. Mohtadi NG, Gryphon DR, Pedersen ME et al. The development and validation of a self-administered quality-of-life outcome measure for young, active patients with symptomatic hip disease: the International Hip Outcome Tool (iHOT-33). *Arthroscopy* 2012;28(5):595-605; quiz 606-10 e1.
55. Kemp JL, Collins NJ, Roos EM et al. Patient-reported outcome scores for hip arthroscopy demonstrate adequate psychometric properties. *Osteoarthritis Cartilage* 2013;21:S277-8.
56. Westacott DJ, Minns JL, Foguet P. The diagnostic accuracy of magnetic resonance imaging and ultrasonography in gluteal tendon tears—a systematic review. *Hip Int* 2011;21(6):637-45.
57. Jin W, Kim KI, Rhyu KH et al. Sonographic evaluation of anterosuperior hip labral tears with magnetic resonance arthrographic and surgical correlation. *J Ultrasound Med* 2012;31(3):439-47.
58. Troelsen A, Mechlenburg I, Gelinek J et al. What is the role of clinical tests and ultrasound in acetabular labral tear diagnostics? *Acta Orthop* 2009;80(3):314-8.
59. Nishii T, Tanaka H, Sugano N et al. Disorders of acetabular labrum and articular cartilage in hip dysplasia: evaluation using isotropic high-resolution CT arthrography with sequential radial reformation. *Osteoarthritis Cartilage* 2007;15(3):251-7.
60. Yamamoto Y, Tonotsuka H, Ueda T et al. Usefulness of radial contrast-enhanced computed tomography for the diagnosis of acetabular labrum injury. *Arthroscopy* 2007;23(12):1290-4.
61. Christie-Large M, Tapp MJ, Theivendran K et al. The role of multidetector CT arthrography in the investigation of suspected intra-articular hip pathology. *Br J Radiol* 2010;83(994):861-7.
62. Ha YC, Choi JA, Lee YK et al. The diagnostic value of direct CT arthrography using MDCT in the evaluation of acetabular labral tear: with arthroscopic correlation. *Skeletal Radiol* 2013;42(5):681-8.
63. Jung JY, Kim GU, Lee HJ et al. Diagnostic value of ultrasound and computed tomographic arthrography in diagnosing anterosuperior acetabular labral tears. *Arthroscopy* 2013;29(11):1769-76.
64. Sahin M, Calisir C, Omeroglu H et al. Evaluation of labral pathology and hip articular cartilage in patients with femoroacetabular impingement (FAI): comparison of multidetector CT arthrography and MR arthrography. *Pol J Radiol* 2014;79:374-80.
65. Smith TO, Hilton G, Toms AP et al. The diagnostic accuracy of acetabular labral tears using magnetic resonance imaging and magnetic resonance arthrography: a meta-analysis. *Eur Radiol* 2011;21(4):863-74.
66. Gold SL, Burge AJ, Potter HG. MRI of hip cartilage: joint morphology, structure and composition. *Clin Orthop Relat Res* 2012;470(12):3321-31.
67. Nishii T, Tanaka H, Nakanishi K et al. Fat-suppressed 3D spoiled gradient-echo MRI and MDCT arthrography of articular cartilage in patients with hip dysplasia. *AJR Am J Roentgenol* 2005;185(2):379-85.
68. Perdikakis E, Karachalios T, Katonis P et al. Comparison of MR-arthrography and MDCT-arthrography for detection of labral and articular cartilage hip pathology. *Skeletal Radiol* 2011;40(11):1441-7.
69. Knuesel PR, Pfirrmann CW, Noetzi HP et al. MR arthrography of the hip: diagnostic performance of a dedicated water-excitation 3D double-echo steady-state sequence to detect cartilage lesions. *AJR Am J Roentgenol* 2004;183(6):1729-35.
70. Schmid MR, Notzli HP, Zanetti M et al. Cartilage lesions in the hip: diagnostic effectiveness of MR arthrography. *Radiology* 2003;226(2):382-6.
71. Zaragoza E, Lattanzio PJ, Beaulieu PE. Magnetic resonance imaging with gadolinium arthrography to assess acetabular cartilage delamination. *Hip Int* 2009;19(1):18-23.
72. Fearon AM, Scarvell JM, Cook JL et al. Does ultrasound correlate with surgical or histologic findings in greater trochanteric pain syndrome? A pilot study. *Clin Orthop Relat Res* 2010;468(7):1838-44.
73. Ross JR, Nepple JJ, Philippon MJ et al. Effect of changes in pelvic tilt on range of motion to impingement and radiographic parameters of acetabular morphologic characteristics. *Am J Sports Med* 2014;42(10):2402-9.
74. Sankar WN, Matheney TH, Zaltz I. Femoroacetabular impingement: current concepts and controversies. *Orthop Clin North Am* 2013;44(4):575-89.
75. Ganz R, Parvizi J, Beck M et al. Femoroacetabular impingement: a cause for osteoarthritis of the hip. *Clin Orthop Relat Res* 2003;417:112-20.
76. Agricola R, Waarsing JH, Arden NK et al. Cam impingement of the hip: a risk factor for hip osteoarthritis. *Nat Rev Rheumatol* 2013;9(10):630-4.
77. Hack K, Di Primio G, Rakhra K et al. Prevalence of cam-type femoroacetabular impingement morphology in asymptomatic volunteers. *J Bone Joint Surg Am* 2010;92(14):2436-44.

78. Siebenrock KA, Ferner F, Noble PC et al. The cam-type deformity of the proximal femur arises in childhood in response to vigorous sporting activity. *Clin Orthop Relat Res* 2011;469(11):3229-40.
79. Gosvig KK, Jacobsen S, Sonne-Holm S et al. The prevalence of cam-type deformity of the hip joint: a survey of 4151 subjects of the Copenhagen Osteoarthritis Study. *Acta Radiol* 2008;49(4):436-41.
80. Kang C, Hwang DS, Cha SM. Acetabular labral tears in patients with sports injury. *Clin Orthop Surg* 2009;1(4):230-5.
81. Kapron AL, Peters CL, Aoki SK et al. The prevalence of radiographic findings of structural hip deformities in female collegiate athletes. *Am J Sports Med* 2015;43(6):1324-30.
82. Frank JM, Harris JD, Erickson BJ et al. Prevalence of femoroacetabular impingement imaging findings in asymptomatic volunteers: a systematic review. *Arthroscopy* 2015;31(6):1199-204.
83. Murray RO, Duncan C. Athletic activity in adolescence as an etiological factor in degenerative hip disease. *J Bone Joint Surg Br* 1971;53(3):406-19.
84. Agricola R, Bessems JH, Ginai AZ et al. The development of cam-type deformity in adolescent and young male soccer players. *Am J Sports Med* 2012;40(5):1099-106.
85. Agricola R, Heijboer MP, Ginai AZ et al. A cam deformity is gradually acquired during skeletal maturation in adolescent and young male soccer players: a prospective study with minimum 2-year follow-up. *Am J Sports Med* 2014;42(4):798-806.
86. Philippon MJ, Ho CP, Briggs KK et al. Prevalence of increased alpha angles as a measure of cam-type femoroacetabular impingement in youth ice hockey players. *Am J Sports Med* 2013;41(6):1357-62.
87. Samora JB, Ng VY, Ellis TJ. Femoroacetabular impingement: a common cause of hip pain in young adults. *Clin J Sport Med* 2011;21(1):51-6.
88. Allen D, Beaulé PE, Ramadan O et al. Prevalence of associated deformities and hip pain in patients with cam-type femoroacetabular impingement. *J Bone Joint Surg Br* 2009;91(5):589-94.
89. Larson CM, Sikka RS, Sardelli MC et al. Increasing alpha angle is predictive of athletic-related 'hip' and 'groin' pain in collegiate National Football League prospects. *Arthroscopy* 2013;29(3):405-10.
90. Siebenrock KA, Kaschka I, Frauchiger L et al. Prevalence of cam-type deformity and hip pain in elite ice hockey players before and after the end of growth. *Am J Sports Med* 2013;41(10):2308-13.
91. Pollard TC, McNally EG, Wilson DC et al. Localised cartilage assessment with three-dimensional dGEMRIC in asymptomatic hips with normal morphology and cam deformity. *J Bone Joint Surg Am* 2010;92(15):2557-69.
92. Glyn-Jones S, Palmer AJ, Agricola R et al. Osteoarthritis. *Lancet* 2015;386(9991):376-87.
93. Agricola R, Heijboer MP, Bierma-Zeinstra SM et al. Cam impingement causes osteoarthritis of the hip: a nationwide prospective cohort study (CHECK). *Ann Rheum Dis* 2013;72(6):918-23.
94. Nicholls AS, Kiran A, Pollard TCB et al. The association between hip morphology parameters and nineteen-year risk of end-stage osteoarthritis of the hip: a nested case-control study. *Arthritis Rheum* 2011;63(11):3392-400.
95. Thomas GE, Palmer AJ, Batra RN et al. Subclinical deformities of the hip are significant predictors of radiographic osteoarthritis and joint replacement in women. A 20 year longitudinal cohort study. *Osteoarthritis Cartilage* 2014;22(10):1504-10.
96. Agricola R, Heijboer MP, Roze RH et al. Pincer deformity does not lead to osteoarthritis of the hip whereas acetabular dysplasia does: acetabular coverage and development of osteoarthritis in a nationwide prospective cohort study (CHECK). *Osteoarthritis Cartilage* 2013;21(10):1514-21.
97. McCarthy JC, Noble PC, Schuck MR et al. The watershed labral lesion: its relationship to early arthritis of the hip. *J Arthroplasty* 2001;16(8 Suppl 1):81-7.
98. Martin RL, Ensey KR, Draovitch P et al. Acetabular labral tears of the hip: examination and diagnostic challenges. *J Orthop Sports Phys Ther* 2006;36(7):503-15.
99. Siebenrock KA, Schoeniger R, Ganz R. Anterior femoro-acetabular impingement due to acetabular retroversion—treatment with periacetabular osteotomy. *J Bone Joint Surg Am* 2003;85a(2):278-86.
100. Dorrell JH, Catterall A. The torn acetabular labrum. *J Bone Joint Surg Br* 1986;68(3):400-3.
101. McCarthy JC, Lee JA. Acetabular dysplasia: a paradigm of arthroscopic examination of chondral injuries. *Clin Orthop Relat Res* 2002;405:122-8.
102. Farjo LA, Glick JM, Sampson TG. Hip arthroscopy for acetabular labral tears. *Arthroscopy* 1999;15(2):132-7.
103. Lewis CL, Sahrman SA, Moran DW. Anterior hip joint force increases with hip extension, decreased gluteal force or decreased iliopsoas force. *J Biomech* 2007;40(16):3725-31.
104. Seldes RM, Tan V, Hunt J et al. Anatomy, histologic features and vascularity of the adult acetabular labrum. *Clin Orthop Relat Res* 2001;382:232-40.
105. Lewis CL, Sahrman SA, Moran DW. Effect of hip angle on anterior hip joint force during gait. *Gait Posture* 2010;32(4):603-7.
106. Grey AJ, Villar RN. The ligamentum teres of the hip: an arthroscopic classification of its pathology. *Arthroscopy* 1997;13(5):575-8.
107. Bardakos NV, Vasconcelos JC, Villar RN. Early outcome of hip arthroscopy for femoroacetabular impingement: the role of femoral osteoplasty in symptomatic improvement. *J Bone Joint Surg Br* 2008;90(12):1570-5.
108. McCarthy JC, Lee JA. Hip arthroscopy: indications, outcomes and complications. *J Bone Joint Surg Am* 2005;87a(5):1138-45.
109. Kemp JL, MacDonald D, Collins NJ et al. Hip arthroscopy in the setting of hip osteoarthritis: systematic review of outcomes

- and progression to hip arthroplasty. *Clin Orthop Relat Res* 2015;473(3):1055-73.
110. Grimaldi A, Richardson C, Durbridge G et al. The association between degenerative hip joint pathology and size of the gluteus maximus and tensor fascia lata muscles. *Man Ther* 2009;14(6):611-7.
 111. Grimaldi A, Richardson C, Stanton W et al. The association between degenerative hip joint pathology and size of the gluteus medius, gluteus minimus and piriformis muscles. *Man Ther* 2009;14(6):605-10.
 112. Baldwin KD, Harrison RA, Namdari S et al. Outcomes of hip arthroscopy for treatment of femoroacetabular impingement: a systematic review. *Curr Orth Practice* 2009;20(6):669-73.
 113. Bedi A, Chen N, Robertson W et al. The management of labral tears and femoroacetabular impingement of the hip in the young, active patient. *Arthroscopy* 2008;24(10):1135-45.
 114. Byrd JW, Jones KS. Prospective analysis of hip arthroscopy with 10-year followup. *Clin Orthop Relat Res* 2010;468(3):741-6.
 115. Enseki KR, Martin R, Kelly BT. Rehabilitation after arthroscopic decompression for femoroacetabular impingement. *Clin Sports Med* 2010;29(2):247-55, viii.
 116. Shindle MK, Domb BG, Kelly BT. Hip and pelvic problems in athletes. *Op Techn Sports Med* 2007;15(4):195-203.
 117. Stalzer S, Wahoff M, Scanlan M. Rehabilitation following hip arthroscopy. *Clin Sports Med* 2006;25(2):337-57, x.
 118. Tyler TF, Nicholas SJ, Campbell RJ et al. The association of hip strength and flexibility with the incidence of adductor muscle strains in professional ice hockey players. *Am J Sports Med* 2001;29(2):124-8.
 119. Kemp JL, Schache A, Makdissi M et al. Is hip range of motion and strength impaired in people with hip chondrolabral pathology? *J Musculoskelet Neuronal Interact* 2014;14(3):334-42.
 120. Nussbaumer S, Leunig M, Glatthorn JF et al. Validity and test-retest reliability of manual goniometers for measuring passive hip range of motion in femoroacetabular impingement patients. *BMC Musculoskelet Disord* 2010;11:194.
 121. Casartelli NC, Maffiuletti NA, Item-Glatthorn JF et al. Hip muscle weakness in patients with symptomatic femoroacetabular impingement. *Osteoarthritis Cartilage* 2011;19(7):816-21.
 122. Harris-Hayes M, Mueller MJ, Sahrman SA et al. Persons with chronic hip joint pain exhibit reduced hip muscle strength. *J Orthop Sports Phys Ther* 2014;44(11):890-8.
 123. Hatton AL, Kemp JL, Brauer SG et al. Dynamic single-leg balance performance is impaired in individuals with hip chondropathy. *Arthritis Care Res* 2014;66(5):709-16.
 124. Brisson N, Lamontagne M, Kennedy MJ et al. The effects of cam femoroacetabular impingement corrective surgery on lower-extremity gait biomechanics. *Gait Posture* 2013;37(2):258-63.
 125. Hunt MA, Guenther JR, Gilbert MK. Kinematic and kinetic differences during walking in patients with and without symptomatic femoroacetabular impingement. *Clin Biomech (Bristol, Avon)* 2013;28(5):519-23.
 126. Mendis MD, Wilson SJ, Hayes DA et al. Hip flexor muscle size, strength and recruitment pattern in patients with acetabular labral tears compared to healthy controls. *Man Ther* 2014;19(5):405-10.
 127. Lewis CL, Sahrman SA. Muscle activation and movement patterns during prone hip extension exercise in women. *J Athl Train* 2009;44(3):238-48.
 128. Hodges PW, Mellor R, Crossley K et al. Pain induced by injection of hypertonic saline into the infrapatellar fat pad and effect on coordination of the quadriceps muscles. *Arthritis Rheum* 2009;61(1):70-7.
 129. Kemp JL, Collins NJ, Makdissi M et al. Hip arthroscopy for intra-articular pathology: a systematic review of outcomes with and without femoral osteoplasty. *Br J Sports Med* 2012;46(9):632-43.
 130. Philippon MJ, Christensen JC, Wahoff MS. Rehabilitation after arthroscopic repair of intra-articular disorders of the hip in a professional football athlete. *J Sport Rehabil* 2009;18(1):118-34.
 131. Cheatham SW, Enseki KR, Kolber MJ. Post-operative rehabilitation after hip arthroscopy: a search for the evidence. *J Sport Rehabil* 2015;24(4):413-8.
 132. Connell DA, Bass C, Sykes CA et al. Sonographic evaluation of gluteus medius and minimus tendinopathy. *Eur Radiol* 2003;13(6):1339-47.
 133. Long SS, Surrey DE, Nazarian LN. Sonography of greater trochanteric pain syndrome and the rarity of primary bursitis. *AJR Am J Roentgenol* 2013;201(5):1083-6.
 134. Huang BK, Campos JC, Michael Peschka PG et al. Injury of the gluteal aponeurotic fascia and proximal iliotibial band: anatomy, pathologic conditions and MR imaging. *Radiographics* 2013;33(5):1437-52.
 135. Fearon A, Stephens S, Cook J et al. The relationship of femoral neck shaft angle and adiposity to greater trochanteric pain syndrome in women. A case control morphology and anthropometric study. *Br J Sports Med* 2012;46(12):888-92.
 136. Shimatsu K, Subramaniam S, Sim H et al. Ciprofloxacin-induced tendinopathy of the gluteal tendons. *J Gen Intern Med* 2014;29(11):1559-62.
 137. Kirchgessner T, Larbi A, Omoumi P et al. Drug-induced tendinopathy: from physiology to clinical applications. *Joint Bone Spine* 2014;81(6):485-92.
 138. Khan KM, Scott A. Mechanotherapy: how physical therapists' prescription of exercise promotes tissue repair. *Br J Sports Med* 2009;43(4):247-52.
 139. Magnusson SP, Langberg H, Kjaer M. The pathogenesis of tendinopathy: balancing the response to loading. *Nat Rev Rheumatol* 2010;6(5):262-8.
 140. Cook JL, Purdam C. Is compressive load a factor in the development of tendinopathy? *Br J Sports Med* 2012;46(3):163-8.

141. Almekinders LC, Weinhold PS, Maffulli N. Compression aetiology in tendinopathy. *Clin Sports Med* 2003;22(4):703-10.
142. Vogel KG, Ordog A, Pogany G et al. Proteoglycans in the compressed region of human tibialis posterior tendon and in ligaments. *J Orthop Res* 1993;11(1):68-77.
143. Thornton GM, Shao X, Chung M et al. Changes in mechanical loading lead to tendonspecific alterations in MMP and TIMP expression: influence of stress deprivation and intermittent cyclic hydrostatic compression on rat supraspinatus and Achilles tendons. *Br J Sports Med* 2010;44(10):698-703.
144. Birnbaum K, Siebert CH, Pandorf T et al. Anatomical and biomechanical investigations of the iliotibial tract. *Surg Radiol Anat* 2004;26(6):433-46.
145. Clancy WG. Runners' injuries. Part two. Evaluation and treatment of specific injuries. *Am J Sports Med* 1980;8(4):287-9.
146. Anderson K, Strickland SM, Warren R. Hip and groin injuries in athletes. *Am J Sports Med* 2001;29(4):521-33.
147. Bass CJ, Connell DA. Sonographic findings of tensor fascia lata tendinopathy: another cause of anterior groin pain. *Skeletal Radiol* 2002;31(3):143-8.
148. Sher I, Umans H, Downie SA et al. Proximal iliotibial band syndrome: what is it and where is it? *Skeletal Radiol* 2011;40(12):1553-6.
149. Fearon AM, Scarvell JM, Neeman T et al. Greater trochanteric pain syndrome: defining the clinical syndrome. *Br J Sports Med* 2013;47(10):649-53.
150. Woodley SJ, Nicholson HD, Livingstone V et al. Lateral hip pain: findings from magnetic resonance imaging and clinical examination. *J Orthop Sports Phys Ther* 2008;38(6):313-28.
151. Kingzett-Taylor A, Tirman PF, Feller J et al. Tendinosis and tears of gluteus medius and minimus muscles as a cause of hip pain: MR imaging findings. *AJR Am J Roentgenol* 1999;173(4):1123-6.
152. Blankenbaker DG, Ullrick SR, Davis KW et al. Correlation of MRI findings with clinical findings of trochanteric pain syndrome. *Skeletal Radiol* 2008;37(10):903-9.
153. Labrosse JM, Cardinal E, Leduc BE et al. Effectiveness of ultrasound-guided corticosteroid injection for the treatment of gluteus medius tendinopathy. *AJR Am J Roentgenol* 2010;194(1):202-6.
154. Rompe JD, Segal NA, Cacchio A et al. Home training, local corticosteroid injection or radial shock wave therapy for greater trochanter pain syndrome. *Am J Sports Med* 2009;37(10):1981-90.
155. Coombes BK, Bisset L, Brooks P et al. Effect of corticosteroid injection, physiotherapy or both on clinical outcomes in patients with unilateral lateral epicondylalgia: a randomised controlled trial. *JAMA* 2013;309(5):461-9.
156. Paavola M, Kannus P, Jarvinen TA et al. Treatment of tendon disorders. Is there a role for corticosteroid injection? *Foot Ankle Clin* 2002;7(3):501-13.
157. Furia JP, Rompe JD, Maffulli N. Low-energy extracorporeal shock wave therapy as a treatment for greater trochanteric pain syndrome. *Am J Sports Med* 2009;37(9):1806-13.
158. Mani-Babu S, Morrissey D, Waugh C et al. The effectiveness of extracorporeal shock wave therapy in lower limb tendinopathy: a systematic review. *Am J Sports Med* 2015;43(3):752-61.
159. van der Worp H, van den Akker-Scheek I, van Schie H et al. ESWT for tendinopathy: technology and clinical implications. *Knee Surg Sports Traumatol Arthrosc* 2013;21(6):1451-8.
160. Kosek E, Lundberg L. Segmental and plurisegmental modulation of pressure pain thresholds during static muscle contractions in healthy individuals. *Eur J Pain* 2003;7(3):251-8.
161. Naugle KM, Fillingim RB, Riley JL, 3rd. A meta-analytic review of the hypoalgesic effects of exercise. *J Pain* 2012;13(12):1139-50.
162. Cook JL, Purdam CR. The challenge of managing tendinopathy in competing athletes. *Br J Sports Med* 2014;48(7):506-9.
163. Rudavsky A, Cook J. Physiotherapy management of patellar tendinopathy (jumper's knee). *J Physiother* 2014;60(3):122-9.
164. Noehren B, Scholz J, Davis I. The effect of real-time gait retraining on hip kinematics, pain and function in subjects with patellofemoral pain syndrome. *Br J Sports Med* 2011;45(9):691-6.
165. Chumanov ES, Wille CM, Michalski MP et al. Changes in muscle activation patterns when running step rate is increased. *Gait Posture* 2012;36(2):231-5.
166. Davies JF, Stiehl JB, Davies JA et al. Surgical treatment of hip abductor tendon tears. *J Bone Joint Surg Am* 2013;95(15):1420-5.
167. Domb BG, Botser I, Giordano BD. Outcomes of endoscopic gluteus medius repair with minimum 2-year follow-up. *Am J Sports Med* 2013;41(5):988-97.
168. Dominguez A, Seijas R, Ares O et al. Clinical outcomes of trochanteric syndrome endoscopically treated. *Arch Orthop Trauma Surg* 2015;135(1):89-94.
169. Govaert LH, van Dijk CN, Zeegers AV et al. Endoscopic bursectomy and iliotibial tract release as a treatment for refractory greater trochanteric pain syndrome: a new endoscopic approach with early results. *Arthrosc Tech* 2012;1(2):e161-4.

CHAPTER 32

1. Bizzini M. The groin area: the Bermuda triangle of sports medicine? *Br J Sports Med* 2011;45:1.
2. Robinson P, Salehi F, Grainger A et al. Cadaveric and MRI study of the musculotendinous contributions to the capsule of the symphysis pubis. *Am J Roent* 2007;188:W440-5.
3. Werner J, Hägglund M, Waldén M et al. UEFA injury study: a prospective study of hip and groin injuries in professional football over seven consecutive seasons. *Br J Sports Med* 2009;43:1036-40.
4. Hölmich P, Thorborg K, Dehlendorff C et al. Incidence and clinical presentation of groin injuries in sub-elite male soccer. *Br J Sports Med* 2014;48(16):1245-50.

5. Serner A, Tol JL, Jomaah N et al. Diagnosis of acute groin injuries: a prospective study of 110 athletes. *Am J Sports Med* 2015;43(8):1857-64.
6. Robertson BA, Barker PJ, Fahrer M et al. The anatomy of the pubic region revisited: implications for the pathogenesis and clinical management of chronic groin pain in athletes. *Sports Med* 2009;39:225-34.
7. Charnock BL, Lewis CL, Garrett WE et al. Adductor longus mechanics during the maximal effort soccer kick. *Sports Biomech* 2009;8:223-34.
8. Polster JM, Elgabaly M, Lee H et al. MRI and gross anatomy of the iliopsoas tendon complex. *Skeletal Radiol* 2007;37:55-8.
9. Condon RE. Reassessment of groin anatomy during the evolution of preperitoneal hernia repair. *Am J Surg* 1996;172:5-8.
10. Emery CA, Meeuwisse WH, Powell JW. Groin and abdominal strain injuries in the National Hockey League. *Clin J Sport Med* 1999;9(3):151-6.
11. Orchard JW. Men at higher risk of groin injuries in elite team sports: a systematic review. *Br J Sports Med* 2015;49:798-802.
12. Weir A, Hölmich P, Schache AG et al. Terminology and definitions on groin pain in athletes: building agreement using a short Delphi method. *Br J Sports Med* 2015;49(12):825-7.
13. Serner A, van Eijck CH, Beumer BR et al. Study quality on groin injury management remains low: a systematic review on treatment of groin pain in athletes. *Br J Sports Med* 2015;49(12):813.
14. Waldén M, Häggglund M, Ekstrand J. The epidemiology of groin injury in senior football: a systematic review of prospective studies. *Br J Sports Med* 2015;49(12):792-7.
15. Thorborg K, Rathleff MS, Petersen P et al. Prevalence and severity of hip and groin pain in sub-elite male football: a cross-sectional cohort study of 695 players. *Scand J Med Sci Sports*. Published online 8 December 2015. doi:10.1111/sms.12623.
16. Hölmich P, Renstrom PA. Long-standing groin pain in sportspeople falls into three primary patterns, a 'clinical entity' approach: a prospective study of 207 patients. *Br J Sports Med* 2007;41:247-52.
17. Lovell G. The diagnosis of chronic groin pain in athletes: a review of 189 cases. *Aust J Sci Med Sport* 1995;27:76-9.
18. Bradshaw CJ, Bundy M, Falvey E. The diagnosis of longstanding groin pain: a prospective clinical cohort study. *Br J Sports Med* 2008;42:551-4.
19. Maffey L, Emery C. What are the risk factors for groin strain injury in sport? A systematic review of the literature. *Sports Med* 2007;37:881-94.
20. Whittaker JL, Small C, Maffey L et al. Risk factors for groin injury in sport: an updated systematic review. *Br J Sports Med* 2015;49(12):803-9.
21. Mosler AB, Agricola R, Weir A et al. *Br J Sports Med* 2015;49(12):810.
22. Weir A, Brukner P, Delahunt E et al. Doha agreement meeting on terminology and definitions in groin pain in athletes. *Br J Sports Med* 2015;49:768-74.
23. Thorborg K, Petersen J, Magnusson SP et al. Clinical assessment of hip strength using a hand-held dynamometer is reliable. *Scand J Med Sci Sports* 2010;20:493-501.
24. Tyler TF, Nicholas SJ, Campbell RJ et al. The association of hip strength and flexibility with the incidence of adductor muscle strains in professional ice hockey players. *Am J Sports Med* 2001;29:124-8.
25. Thorborg K, Serner A, Petersen J et al. Hip adduction and abduction strength profiles in elite soccer players: implications for clinical evaluation of hip adductor muscle recovery after injury. *Am J Sports Med* 2011;39:121-6.
26. Nevin F, Delahunt E. Adductor squeeze test values and hip joint range of motion in Gaelic football athletes with longstanding groin pain. *J Sci Med Sport* 2014;17:155-9.
27. Thorborg K, Branci S, Nielsen MP et al. Eccentric and isometric hip adduction strength in male soccer players with and without adductor-related groin pain: an assessor-blinded comparison. *Orthop J Sports Med* 2014;2(2):2325967114521778.
28. Malliaras P, Hogan A, Nawrocki A et al. Hip flexibility and strength measures: reliability and association with athletic groin pain. *Br J Sports Med* 2009;43:739-44.
29. Light N, Thorborg K. The precision and torque production of common hip adductor squeeze tests used in elite football. *J Sci Med Sport*. Published online: 12 December 2015. doi:10.1016/j.jsams.2015.12.009.
30. Thorborg K, Tijssen M, Habets B et al. Patient-Reported Outcome (PRO) questionnaires for young-aged to middle-aged adults with hip and groin disability: a systematic review of the clinimetric evidence. *Br J Sports Med* 2015;49(12):812.
31. Thorborg K, Hölmich P, Christensen R et al. The Copenhagen Hip and Groin Outcome Score (HAGOS): development and validation according to the COSMIN checklist. *Br J Sports Med* 2011;45:478-91.
32. Thorborg K, Branci S, Stensbirk F et al. Copenhagen hip and groin outcome score (HAGOS) in male soccer: reference values for hip and groin injury-free players. *Br J Sports Med* 2014;48(7):557-9.
33. Branci S, Thorborg K, Nielsen MB et al. Radiological findings in symphyseal and adductor-related groin pain in athletes: a critical review of the literature. *Br J Sports Med* 2013;47:611-9.
34. Branci S, Thorborg K, Bech BH et al. MRI findings in soccer players with long-standing adductor-related groin pain and asymptomatic controls. *Br J Sports Med* 2015;49(10):681-91.
35. Robinson P, Grainger AJ, Hensor EMA et al. Do MRI and ultrasound of the anterior pelvis correlate with or predict, young football players' clinical findings? A 4-year prospective study of elite academy soccer players. *Br J Sports Med* 2015;49:176-82.
36. Weir A, de Vos RJ, Moen M et al. Prevalence of radiological signs of femoroacetabular impingement in patients presenting with long-standing adductor-related groin pain. *Br J Sports Med* 2009;45:6-9.
37. Hölmich P, Thorborg K, Nyvold P et al. Does bony hip morphology affect the outcome of treatment for patients with

- adductor-related groin pain? Outcome 10 years after baseline assessment. *Br J Sports Med* 2014;48(16):1240-4.
38. Harris NH, Murray RO. Lesions of the symphysis in athletes. *Br Med J* 1974;4:211-4.
 39. Verrall GM, Henry L, Fazzalari NL et al. Bone biopsy of the parasymphyseal pubic bone region in athletes with chronic groin injury demonstrates new woven bone formation consistent with a diagnosis of pubic bone stress injury. *Am J Sports Med* 2008;36:2425-31.
 40. Slavotinek JP. Groin pain in footballers: the association between preseason clinical and pubic bone magnetic resonance imaging findings and athlete outcome. *Am J Sports Med* 2005;33:894-9.
 41. Lovell G, Galloway H, Hopkins W et al. Osteitis pubis and assessment of bone marrow oedema at the pubic symphysis with MRI in an elite junior male soccer squad. *Clin J Sport Med* 2006;16:117-22.
 42. Robinson P, Barron DA, Parsons W et al. Adductor-related groin pain in athletes: correlation of MR imaging with clinical findings. *Skeletal Radiol* 2004;33:451-7.
 43. Falvey E, King E, Kinsella S et al. Athletic groin pain (part 1): a prospective anatomical diagnosis of 382 patients—clinical findings, MRI findings and patient-reported outcome measures at baseline. *Br J Sports Med* 2016;50(7):423-30.
 44. Branci S, Thorborg K, Bech BH et al. The Copenhagen Standardised MRI protocol to assess the pubic symphysis and adductor regions of athletes: outline and intratester and intertester reliability. *Br J Sports Med* 2015;49:692-9.
 45. de Jonge S, Tol JL, Weir A et al. The tendon structure returns to asymptomatic values in nonoperatively treated Achilles tendinopathy but is not associated with symptoms: a prospective study. *Am J Sports Med* 2015;43:2950-8.
 46. Brennan D, O'Connell MJ, Ryan M et al. Secondary cleft sign as a marker of injury in athletes with groin pain: MR image appearance and interpretation. *Radiology* 2005;235:162-7.
 47. van Schie HTM, de Vos RJ, de Jonge S et al. Ultrasonographic tissue characterisation of human Achilles tendons: quantification of tendon structure through a novel non-invasive approach. *Br J Sports Med* 2010;44:1153-9.
 48. Kulig K, Landel R, Chang Y-J et al. Patellar tendon morphology in volleyball athletes with and without patellar tendinopathy. *Scand J Med Sci Sports* 2013;23:e81-8.
 49. Saily M, Whiteley R, Read JW et al. Pubic apophysitis: a previously undescribed clinical entity of groin pain in athletes. *Br J Sports Med* 2015;49:828-34.
 50. Hölmich P, Uhrskou P, Ulnits L et al. Effectiveness of active physical training as treatment for long-standing adductor-related groin pain in athletes: randomised trial. *Lancet* 1999;353:439-43.
 51. Rio E, Kidgell D, Purdam C et al. Isometric exercise induces analgesia and reduces inhibition in patellar tendinopathy. *Br J Sports Med* 2015;49:1277-83.
 52. Chaudhari AMW, Jamison ST, McNally MP et al. Hip adductor activations during run-to-cut manoeuvres in compression shorts: implications for return to sport after groin injury. *J Sports Sci* 2014;32:1333-40.
 53. Weir A, Veger SAS, Van de Sande HBA et al. A manual therapy technique for chronic adductor-related groin pain in athletes: a case series. *Scand J Med Sci Sports* 2009;19:616-20.
 54. Weir A, Jansen JACG, van de Port IGL et al. Manual or exercise therapy for long-standing adductor-related groin pain: a randomised controlled clinical trial. *Man Ther* 2011;16:148-54.
 55. Thorborg K, Bandholm T, Zebis M et al. Large strengthening effect of a hip-flexor training programme: a randomised controlled trial. *Knee Surg Sports Traumatol Arthrosc* 2016;24(7):2346-52.
 56. Kroger EW, Griesser MJ, Kolovich GP et al. Efficacy of surgery for internal snapping hip. *Int J Sports Med* 2013;34:851-5.
 57. Sheen AJ, Stephenson BM, Lloyd DM et al. 'Treatment of the sportsman's groin': British Hernia Society's 2014 position statement based on the Manchester Consensus Conference. *Br J Sports Med* 2014;48:1079-87.
 58. Garvey JFW. Sportsman's groin: reaching a consensus. *Hernia* 2014;18:811-3.
 59. Hölmich P. Long-standing groin pain in sportspeople falls into three primary patterns, a 'clinical entity' approach: a prospective study of 207 patients. *Br J Sports Med* 2007;41:247-52; discussion 252.
 60. Yuill EA, Pajaczkowski JA, Howitt SD. Conservative care of sports hernias within soccer players: a case series. *J Bodywork Mov Ther* 2012;16:540-8.
 61. Woodward JS, Parker A, MacDonald RM. Non-surgical treatment of a professional hockey player with the signs and symptoms of sports hernia: a case report. *Int J Sports Phys Ther* 2012;7:85.
 62. Schoenfeld BJ, Contreras B, Tiriyaki-Sonmez G et al. An electromyographic comparison of a modified version of the plank with a long lever and posterior tilt versus the traditional plank exercise. *Sports Biomech* 2014;13:296-306.
 63. Sundstrup E, Jakobsen MD, Andersen CH et al. Swiss ball abdominal crunch with added elastic resistance is an effective alternative to training machines. *Int J Sports Phys Ther* 2012;7:372-80.
 64. Sheen AJ, Paaanen H. The next step towards rational treatment for 'the sportsman's groin'. *Br J Sports Med* 2015;49:764-5.
 65. Paaanen H, Brinck T, Hermunen H et al. Laparoscopic surgery for chronic groin pain in athletes is more effective than nonoperative treatment: a randomised clinical trial with magnetic resonance imaging of 60 patients with sportsman's hernia (athletic pubalgia). *Surgery* 2011;150:99-107.
 66. Fitzgibbons RJ, Forse RA. Clinical practice. Groin hernias in adults. *N Engl J Med* 2015;372:756-63.
 67. Verrall GM, Slavotinek JP, Barnes PG et al. Description of pain provocation tests used for the diagnosis of sports-related chronic groin pain: relationship of tests to defined clinical (pain and tenderness) and MRI (pubic bone marrow oedema) criteria. *Scand J Med Sci Sports* 2005;15:36-42.
 68. Drew MK, Osmotherly PG, Chiarelli PE. Imaging and clinical tests for the diagnosis of long-standing groin pain in athletes. A systematic review. *Phys Ther Sport* 2014;15(2):124-9.

69. Verrall GM, Slavotinek JP, Fon GT et al. Outcome of conservative management of athletic chronic groin injury diagnosed as pubic bone stress injury. *Am J Sports Med* 2007;35:467-74.
70. Schlegel TF, Bushnell BD, Godfrey J et al. Success of nonoperative management of adductor longus tendon ruptures in National Football League athletes. *Am J Sports Med* 2009;37:1394-9.
71. Thorborg K, Petersen J, Nielsen MB et al. Clinical recovery of two hip adductor longus ruptures: a case-report of a soccer player. *BMC Res Notes* 2013;6:205.
72. Bradshaw C, McCrory P, Bell S et al. Obturator nerve entrapment. A cause of groin pain in athletes. *Am J Sports Med* 1997;25:402-8.
73. Brukner P, Bradshaw C, McCrory P. Obturator neuropathy: a cause of exercise-related groin pain. *Phys Sportsmed* 1999;27:62-73.
74. Labat J-J, Riant T, Robert R et al. Diagnostic criteria for pudendal neuralgia by pudendal nerve entrapment (Nantes criteria). *Neurourol Urodyn* 2008;27:306-10.
75. Lappe J, Cullen D, Haynatzki G et al. Calcium and vitamin D supplementation decreases incidence of stress fractures in female navy recruits. *J Bone Miner Res* 2008;23:741-9.
76. Shaffer RA, Rauh MJ, Brodine SK et al. Predictors of stress fracture susceptibility in young female recruits. *Am J Sports Med* 2006;34:108-15.
77. Esteve E, Rathleff MS, Bagur-Calafat C et al. Prevention of groin injuries in sports: a systematic review with meta-analysis of randomised controlled trials. *Br J Sports Med* 2015;49(12):785-91.
78. Thorborg K, Rathleff MS, Petersen P et al. Prevalence and severity of hip and groin pain in sub-elite male football: a cross-sectional cohort study of 695 players. *Scand J Med Sci Sports*. Published online: 8 Dec 2015. doi: 10.1111/sms.12623.
79. Emery CA, Meeuwisse WH. Risk factors for groin injuries in hockey. *Med Sci Sports Exerc* 2001;33:1423-33.
80. Crow JF, Pearce AJ, Veale JP et al. Hip adductor muscle strength is reduced preceding and during the onset of groin pain in elite junior Australian football players. *J Sci Med Sport* 2010;13:202-4.
5. Gyftopoulos S, Rosenberg ZS, Schweitzer ME et al. Normal anatomy and strains of the deep musculotendinous junction of the proximal rectus femoris: MRI features. *Am J Roentgenol* 2008;190(3):W182-6.
6. Kassarian A, Rodrigo RM, Santisteban JM. Current concepts in MRI of rectus femoris musculotendinous (myotendinous) and myofascial injuries in elite athletes. *Eur J Radiol* 2012;81(12):3763-71.
7. Mendiguchia J, Alentorn-Geli E, Idoate F et al. Rectus femoris muscle injuries in football: a clinically relevant review of mechanisms of injury, risk factors and preventive strategies. *Br J Sports Med* 2013;47(6):359-66.
8. Kassarian A, Rodrigo RM, Santisteban JM. Intramuscular degloving injuries to the rectus femoris: findings at MRI. *Am J Roentgenol* 2014;202(5):W475-80.
9. Chen YT, Tenforde AS, Fredericson M. Update on stress fractures in female athletes: epidemiology, treatment, and prevention. *Curr Rev Musculoskel Med* 2013;6(2):173-81.
10. Veal C, Auyong DB, Hanson NA et al. Delayed quadriceps weakness after continuous adductor canal block for total knee arthroplasty: a case report. *Acta Anaesthesiol Scand* 2014;58(3):362-4.
11. Mens JM, Vleeming A, Snijders CJ et al. Reliability and validity of the active straight leg raise test in posterior pelvic pain since pregnancy. *Spine* 2001;26(10):1167-71.
12. Ross MD, Langford B, Whelan PJ. Test-retest reliability of 4 single-leg horizontal hop tests. *J Strength Cond Res* 2002;16(4):617-22.
13. Augustsson J, Thomee R, Linden C et al. Single-leg hop testing following fatiguing exercise: reliability and biomechanical analysis. *Scand J Med Sci Sports* 2006;16(2):111-20.
14. Kang L, Belcher D, Hulstyn MJ. Stress fractures of the femoral shaft in women's college lacrosse: a report of seven cases and a review of the literature. *Br J Sports Med* 2005;39(12):902-6.
15. Johnson AW, Weiss CB, Wheeler DL. Stress fractures of the femoral shaft in athletes—more common than expected: a new clinical test. *Am J Sports Med* 1994;22(2):248-56.
16. Trainor K, Pinnington MA. Reliability and diagnostic validity of the slump knee bend neurodynamic test for upper/mid lumbar nerve root compression: a pilot study. *Physiotherapy* 2011;97(1):59-64.
17. Butler DS, Jones MA. *Mobilisation of the nervous system*. Melbourne: Churchill Livingstone, 1991.
18. Binkley JM, Stratford PW, Lott SA et al. The Lower Extremity Functional Scale (LEFS): scale development, measurement properties, and clinical application. North American Orthopaedic Rehabilitation Research Network. *Phys Ther* 1999;79(4):371-83.
19. Pua Y-H, Cowan SM, Wrigley TV et al. The Lower Extremity Functional Scale could be an alternative to the Western Ontario and McMaster Universities Osteoarthritis Index physical function scale. *J Clin Epidemiol* 2009;62(10):1103-11.

CHAPTER 33

1. Alonso J-M, Tscholl PM, Engebretsen L et al. Occurrence of injuries and illnesses during the 2009 IAAF World Athletics Championships. *Br J Sports Med* 2014;44(15):1100-5.
2. Ekstrand J, Häggglund M, Walden M. Injury incidence and injury patterns in professional football: the UEFA injury study. *Br J Sports Med* 2011;45(7):553-8.
3. Ekstrand J, Häggglund M, Walden M. Epidemiology of muscle injuries in professional football (soccer). *Am J Sports Med* 2011;39(6):1226-32.
4. Costa PB, Ryan ED, Herda TJ et al. Acute effects of static stretching on peak torque and the hamstrings-to-quadriceps conventional and functional ratios. *Scand J Med Sci Sports* 2013;23(1):38-45.

20. Whiteley R, Jacobsen P, Prior S et al. Correlation of isokinetic and novel hand-held dynamometry measures of knee flexion and extension strength testing. *J Sci Med Sport* 2012;15(5):444-50.
21. Laprade RF, Surowiec RK, Sochanska AN et al. Epidemiology, identification, treatment and return to play of musculoskeletal-based ice hockey injuries. *Br J Sports Med* 2014;48(1):4-10.
22. Ryan JB, Wheeler JH, Hopkinson WJ et al. Quadriceps contusions. West Point update. *Am J Sports Med* 1991;19(3):299-304.
23. Kary JM. Diagnosis and management of quadriceps strains and contusions. *Curr Rev Musculoskel Med* 2010;3(1-4):26-31.
24. Orchard JW, Best TM, Mueller-Wohlfahrt HW et al. The early management of muscle strains in the elite athlete: best practice in a world with a limited evidence basis. *Br J Sports Med* 2008;42(3):158-9.
25. Aspelin P, Ekberg O, Thorsson O et al. Ultrasound examination of soft tissue injury of the lower limb in athletes. *Am J Sports Med* 1992;20(5):601-3.
26. Hayashi D, Hamilton B, Guermazi A et al. Traumatic injuries of thigh and calf muscles in athletes: role and clinical relevance of MR imaging and ultrasound. *Insights Imaging* 2012;3(6):591-601.
27. Alonso A, Hekeik P, Adams R. Predicting a recovery time from the initial assessment of a quadriceps contusion injury. *Aust J Physiother* 2000;46(3):167-77.
28. McBrier NM, Lekan JM, Druhan LJ et al. Therapeutic ultrasound decreases mechano-growth factor messenger ribonucleic acid expression after muscle contusion injury. *Arch Phys Med Rehabil* 2007;88(7):936-40.
29. Shu B, Yang Z, Li X et al. Effect of different intensity pulsed ultrasound on the restoration of rat skeletal muscle contusion. *Cell Biochem Biophysics* 2012;62(2):329-36.
30. Aronen JG, Garrick JG, Chronister RD et al. Quadriceps contusions: clinical results of immediate immobilization in 120 degrees of knee flexion. *Clin J Sport Med* 2006;16(5):383-7.
31. Gerrard DF. The use of padding in rugby union. An overview. *Sports Med* 1998;25:329-32.
32. Riede U, Schmid MR, Romero J. Conservative treatment of an acute compartment syndrome of the thigh. *Arch Orthop Trauma Surg* 2007;127(4):269-75.
33. Ojike NI, Roberts CS, Giannoudis PV. Compartment syndrome of the thigh: a systematic review. *Injury* 2010;41(2):133-6.
34. Diaz JA, Fischer DA, Rettig AC et al. Severe quadriceps muscle contusions in athletes. A report of three cases. *Am J Sports Med* 2003;31(2):289-93.
35. King JB. Post-traumatic ectopic calcification in the muscles of athletes: a review. *Br J Sports Med* 1998;32(4):287-90.
36. Lacout A, Jarraya M, Marcy PY et al. Myositis ossificans imaging: keys to successful diagnosis. *Indian J Radiol Imaging* 2012;22(1):35-9.
37. Miller AE, Davis BA, Beckley OA. Bilateral and recurrent myositis ossificans in an athlete: a case report and review of treatment options. *Arch Phys Med Rehabil* 2006;87(2):286-90.
38. Tsuno MM, Shu GJ. Myositis ossificans. *J Manipulative Physiol Ther* 1990;13(6):340-2.
39. Danchilk JJ, Yochum TR, Aspergren DD. Myositis ossificans traumatica. *J Manip Physiol Ther* 1993;16:605-14.
40. Buselli P, Coco V, Notarnicola A et al. Shock waves in the treatment of post-traumatic myositis ossificans. *Ultrasound Med Biol* 2010;36(3):397-409.
41. Torrance DA, Degraauw C. Treatment of post-traumatic myositis ossificans of the anterior thigh with extracorporeal shock wave therapy. *J Can Chiropr Assoc* 2011;55(4):240-6.
42. Gard K, Ebaugh D. The use of acetic acid iontophoresis in the management of soft tissue injury. *Nth Am J Phys Ther* 2010;5(4):220-6.
43. Wieder DL. Treatment of traumatic myositis ossificans with acetic acid iontophoresis. *Phys Ther* 1992;72(2):133-7.
44. Mani-Babu S, Wolman R, Keen R. Quadriceps traumatic myositis ossificans in a football player: management with intravenous pamidronate. *Clin J Sport Med* 2014;24(5):e56-8.
45. Beatty K, McIntosh A, Savage T et al. Biomechanics of the running drop punt kick with respect to the development of quadriceps strains. *Sport Health* 2007;25(4):18-23.
46. Figved W, Grindem H, Aaberg M et al. Muscle strength measurements and functional outcome of an untreated complete distal rectus femoris muscle tear. *BMJ Case Rep* 2014;2014:bcr2013203191.
47. Orchard JW. Intrinsic and extrinsic risk factors for muscle strains in Australian football. *Am J Sports Med* 2001;29(3):300-3.
48. Hasselman CT, Best TM, Hughes IV C et al. An explanation for various rectus femoris strain injuries using previously undescribed muscle architecture. *Am J Sports Med* 1995;23(4):493-9.
49. Hughes IV C, Hasselman CT, Best TM et al. Incomplete, intrasubstance strain injuries of the rectus femoris muscle. *Am J Sports Med* 1995;23(4):500-6.
50. Garcia VV, Duhrkop DC, Seijas R et al. Surgical treatment of proximal ruptures of the rectus femoris in professional soccer players. *Arch Orthop Trauma Surg* 2012;132(3):329-33.
51. Cross TM, Gibbs N, Houang MT et al. Acute quadriceps muscle strains: magnetic resonance imaging features and prognosis. *Am J Sports Med* 2004;32(3):710-9.
52. Hosalkar HS, Pennock AT, Zaps D et al. The hip antero-superior labral tear with avulsion of rectus femoris (HALTAR) lesion: does the SLAP equivalent in the hip exist? *Hip Int* 2012;22(4):391-6.
53. Small K, Mc Naughton L, Matthews M. A systematic review into the efficacy of static stretching as part of a warm-up for the prevention of exercise-related injury. *Res Sports Med* 2008;16(3):213-31.
54. Yamaguchi T, Ishii K. Effects of static stretching for 30 seconds and dynamic stretching on leg extension power. *J Strength Cond Res* 2005;19(3):677-83.
55. Meroni R, Cerri CG, Lanzarini C et al. Comparison of active stretching technique and static stretching technique on hamstring flexibility. *Clin J Sport Med* 2010;20(1):8-14.

56. Simic L, Sarabon N, Markovic G. Does pre-exercise static stretching inhibit maximal muscular performance? A meta-analytical review. *Scand J Med Sci Sports* 2013;23(2):131-48.
57. Aguilar AJ, DiStefano LJ, Brown CN et al. A dynamic warm-up model increases quadriceps strength and hamstring flexibility. *J Strength Cond Res* 2012;26(4):1130-41.
58. Costa PB, Ryan ED, Herda TJ et al. Effects of static stretching on the hamstrings-to-quadriceps ratio and electromyographic amplitude in men. *J Sports Med Phys Fitness* 2009;49(4):401-9.
59. Sekir U, Arabaci R, Akova B et al. Acute effects of static and dynamic stretching on leg flexor and extensor isokinetic strength in elite women athletes. *Scand J Med Sci Sports* 2010;20(2):268-81.
60. Behm DG, Chaouachi A. A review of the acute effects of static and dynamic stretching on performance. *Eur J Appl Physiol* 2011;111(11):2633-51.
61. Chaouachi A, Castagna C, Chtara M et al. Effect of warm-ups involving static or dynamic stretching on agility, sprinting, and jumping performance in trained individuals. *J Strength Cond Res* 2010;24(8):2001-11.
62. Breugem SJM, Hulscher JBF, Steller P. Stress fracture of the femoral neck in a young female athlete. *Eur J Trauma Emerg Surg* 2009;35:192-5.
63. Arriaza R, Senaris J, Couceiro G et al. Stress fractures of the femur after ACL reconstruction with transfemoral fixation. *Knee Surg Sports Traumatol Arthrosc* 2006;14(11):1148-50.
64. Heng CH, Wang Bde H, Chang PC. Distal femoral fracture after double-bundle anterior cruciate ligament reconstruction surgery. *Am J Sports Med* 2015;43(4):953-6.
65. Behrens SB, Deren ME, Matson A et al. Stress fractures of the pelvis and legs in athletes: a review. *Sports Health* 2013;5(2):165-74.
66. Lee CH, Huang GS, Chao KH et al. Surgical treatment of displaced stress fractures of the femoral neck in military recruits: a report of 42 cases. *Arch Orthop Trauma Surg* 2003;123(10):527-33.
67. Fredericson M, Jennings F, Beaulieu C et al. Stress fractures in athletes. *Top Magn Reson Imaging* 2006;17(5):309-25.
68. Warren MP, Chua AT. Exercise-induced amenorrhea and bone health in the adolescent athlete. *Ann N Y Acad Sci* 2008;1135(1):244-52.
69. Wentz L, Liu P-Y, Haymes E et al. Females have a greater incidence of stress fractures than males in both military and athletic populations: a systemic review. *Mil Med* 2011;176(4):420-30.
70. Tenforde AS, Watanabe LM, Moreno TJ et al. Use of an antigravity treadmill for rehabilitation of a pelvic stress injury. *PM R* 2012;4(8):629-31.
71. Cheatham SW, Kolber MJ, Salameh PA. Meralgia paresthetica: a review of the literature. *Int J Sports Phys Ther* 2013;8(6):883-93.
72. Lorei MP, Hershman EB. Peripheral nerve injuries in athletes. Treatment and prevention. *Sports Med* 1993;16(2):130-47.
73. Khalil N, Nicotra A, Rakowicz W. Treatment for meralgia paraesthetica. *Cochrane Database Syst Rev* 2012;12:CD004159.
74. Herring SA, Akuthota V, Kaufman MS et al. Peripheral nerve injuries of the proximal lower limb in athletes. In Akuthota V, Herring S (eds). *Nerve and Vascular Injuries in Sports Medicine*. New York: Springer, 2009:161-70.
75. Toussaint CP, Perry EC, 3rd, Pisansky MT et al. What's new in the diagnosis and treatment of peripheral nerve entrapment neuropathies. *Neurol Clin* 2010;28(4):979-1004.
76. Maak TG, Osei D, Delos D et al. Peripheral nerve injuries in sports-related surgery: presentation, evaluation, and management: AAOS exhibit selection. *J Bone Joint Surg Am* 2012;94(16):e1211-10.
77. Ulkar B, Yildiz Y, Kunduracioglu B. Meralgia paresthetica: a long-standing performance-limiting cause of anterior thigh pain in a soccer player. *Am J Sports Med* 2003;31(5):787-9.
78. Kalichman L, Vered E, Volchek L. Relieving symptoms of meralgia paresthetica using Kinesio taping: a pilot study. *Arch Phys Med Rehabil* 2010;91(7):1137-9.
79. Lai WH, Shih YF, Lin PL et al. Normal neurodynamic responses of the femoral slump test. *Man Ther* 2012;17(2):126-32.

CHAPTER 34

1. Hallen A, Ekstrand J. Return to play following muscle injuries in professional footballers. *J Sports Sci* 2014;32(13):1229-36.
2. Orchard J, Best TM, Verrall GM. Return to play following muscle strains. *Clin J Sport Med* 2005;15(6):436-41.
3. Battermann N, Appell HJ, Dargel J et al. An anatomical study of the proximal hamstring muscle complex to elucidate muscle strains in this region. *Int J Sports Med* 2011;32(3):211-5.
4. Tubbs RS, Caycedo FJ, Oakes WJ et al. Descriptive anatomy of the insertion of the biceps femoris muscle. *Clin Anat* 2006;19(6):517-21.
5. Kellis E, Galanis N, Natsis K et al. Muscle architecture variations along the human semitendinosus and biceps femoris (long head) length. *J Electromyogr Kinesiol* 2010;20(6):1237-43.
6. Connell DA, Schneider-Kolsky ME, Hoving JL et al. Longitudinal study comparing sonographic and MRI assessments of acute and healing hamstring injuries. *Am J Roentgenol* 2004;183(4):975-84.
7. Gibbs NJ, Cross TM, Cameron M et al. The accuracy of MRI in predicting recovery and recurrence of acute grade one hamstring muscle strains within the same season in Australian Rules football players. *J Sci Med Sport* 2004;7(2):248-58.
8. Verrall GM, Slavotinek JP, Barnes PG et al. Diagnostic and prognostic value of clinical findings in 83 athletes with posterior thigh injury: comparison of clinical findings with magnetic resonance imaging documentation of hamstring muscle strain. *Am J Sports Med* 2003;31(6):969-73.
9. Ekstrand J, Healy JC, Walden M et al. Hamstring muscle injuries in professional football: the correlation of MRI findings with return to play. *Br J Sports Med* 2012;46:112-117.
10. Guillodo Y, Here-Dorignac C, Thoribe B et al. Clinical predictors of time to return to competition following hamstring injuries. *Muscles Ligaments Tendons J* 2014;4(3):386-90.

11. Wangenstein A, Almusa E, Boukarroum S et al. MRI does not add value over and above patient history and clinical examination in predicting time to return to sport after acute hamstring injuries: a prospective cohort of 180 male athletes. *Br J Sports Med* 2015;49(24):1579-87.
12. Warren P, Gabbe BJ, Schneider-Kolsky M et al. Clinical predictors of time to return to competition and of recurrence following hamstring strain in elite Australian footballers. *Br J Sports Med* 2010;44(6):415-9.
13. Moen MH, Reurink G, Weir A et al. Predicting return to play after hamstring injuries. *Br J Sports Med* 2014;48(18):1358-63.
14. De Vos RJ, Reurink G, Goudswaard GJ et al. Clinical findings just after return to play predict hamstring re-injury, but baseline MRI findings do not. *Br J Sports Med* 2014;48(18):1377-84.
15. Reurink G, Brilman EG, de Vos RJ et al. Magnetic resonance imaging in acute hamstring injury: can we provide a return to play prognosis? *Sports Med* 2015;45:133-46.
16. Reurink G, Almusa E, Goudswaard GJ et al. No association between fibrosis on magnetic resonance imaging at return to play and hamstring reinjury risk. *Am J Sports Med* 2015;43(5):1228-34.
17. Orchard J, Seward H. Epidemiology of injuries in the Australian Football League, seasons 1997-2000. *Br J Sports Med* 2002;36(1):39-44.
18. Hawkins RD, Fuller CW. A prospective epidemiological study of injuries in four English professional football clubs. *Br J Sports Med* 1999;33(3):196-203.
19. Hawkins RD, Hulse MA, Wilkinson C et al. The association football medical research programme: an audit of injuries in professional football. *Br J Sports Med* 2001;35(1):43-7.
20. Elliott MC, Zarins B, Powell JW et al. Hamstring muscle strains in professional football players: a 10-year review. *Am J Sports Med* 2011;39(4):843-50.
21. Ekstrand J, Häggglund M, Walden M. Epidemiology of muscle injuries in professional football (soccer). *Am J Sports Med* 2011;39(6):1226-32.
22. Cross KM, Gurka KK, Saliba S et al. Comparison of hamstring strain injury rates between male and female intercollegiate soccer athletes. *Am J Sports Med* 2013;41(4):742-8.
23. Woods C, Hawkins RD, Maltby S et al. The Football Association Medical Research Programme: an audit of injuries in professional football—analysis of hamstring injuries. *Br J Sports Med* 2004;38(1):36-41.
24. Askling C, Lund H, Saartok T et al. Self-reported hamstring injuries in student-dancers. *Scand J Med Sci Sports* 2002;12(4):230-5.
25. Orchard JW, Seward H, Orchard JJ. Results of 2 decades of injury surveillance and public release of data in the Australian Football League. *Am J Sports Med* 2013;41(4):734-41.
26. Askling CM, Tengvar M, Tarassova O et al. Acute hamstring injuries in Swedish elite sprinters and jumpers: a prospective randomised controlled clinical trial comparing two rehabilitation protocols. *Br J Sports Med* 2014;48(7):532-9.
27. Askling CM, Tengvar M, Thorstensson A. Acute hamstring injuries in Swedish elite football: a prospective randomised controlled clinical trial comparing two rehabilitation protocols. *Br J Sports Med* 2013;47(15):953-9.
28. Askling C. Types of hamstring injuries in sports. *Br J Sports Med*. Published online: 2011;45:e2 doi:10.1136/bjsm.2010.081570.15.
29. Askling C, Tengvar M, Saartok T et al. Sports related hamstring strains—two cases with different etiologies and injury sites. *Scand J Med Sci Sports* 2000;10(5):304-7.
30. Askling CM, Malliaropoulos N, Karlsson J. High-speed running type or stretching-type of hamstring injuries makes a difference to treatment and prognosis. *Br J Sports Med* 2012;46(2):86-7.
31. Askling CM, Tengvar M, Saartok T et al. Acute first-time hamstring strains during high-speed running: a longitudinal study including clinical and magnetic resonance imaging findings. *Am J Sports Med* 2007;35(2):197-206.
32. Askling C, Thorstensson A. Hamstring muscle strain in sprinters. *New Stud Athlet* 2008;23(3):67-79.
33. Woodley SJ, Mercer SR. Hamstring muscles: architecture and innervation. *Cells Tissues Organs* 2005;179(3):125-41.
34. Askling C, Saartok T, Thorstensson A. Type of acute hamstring strain affects flexibility, strength and time to return to pre-injury level. *Br J Sports Med* 2006;40(1):40-4.
35. Askling CM, Tengvar M, Saartok T et al. Acute first-time hamstring strains during slow-speed stretching: clinical, magnetic resonance imaging and recovery characteristics. *Am J Sports Med* 2007;35(10):1716-24.
36. Brooks JH, Fuller CW, Kemp SP et al. Incidence, risk and prevention of hamstring muscle injuries in professional rugby union. *Am J Sports Med* 2006;34(8):1297-306.
37. Gabbe BJ, Finch CF, Bennell KL et al. Risk factors for hamstring injuries in community level Australian football. *Br J Sports Med* 2005;39(2):106-10.
38. Heiderscheit BC, Hoerth DM, Chumanov ES et al. Identifying the time of occurrence of a hamstring strain injury during treadmill running: a case study. *Clin Biomech (Bristol, Avon)* 2005;20(10):1072-8.
39. Schache AG, Wrigley TV, Baker R et al. Biomechanical response to hamstring muscle strain injury. *Gait Posture* 2009;29(2):332-8.
40. Chumanov ES, Heiderscheit BC, Thelen DG. Hamstring musculotendon dynamics during stance and swing phases of high-speed running. *Med Sci Sports Exerc* 2011;43(3):525-32.
41. Silder A, Sherry MA, Sanfilippo J et al. Clinical and morphological changes following 2 rehabilitation programs for acute hamstring strain injuries: a randomised clinical trial. *J Orthop Sports Phys Ther* 2013;43(5):284-99.
42. Askling CM, Tengvar M, Saartok T et al. Proximal hamstring strains of stretching type in different sports: injury situations, clinical and magnetic resonance imaging characteristics and return to sport. *Am J Sports Med* 2008;36(9):1799-804.
43. Heiderscheit BC, Sherry MA, Silder A et al. Hamstring strain injuries: recommendations for diagnosis, rehabilitation and injury prevention. *J Orthop Sports Phys Ther* 2010;40(2):67-81.

44. Comin J, Malliaras P, Baquie P et al. Return to competitive play after hamstring injuries involving disruption of the central tendon. *Am J Sports Med* 2013;41(1):111-5.
45. Schache AG, Koulouris G, Kofoed W et al. Rupture of the conjoint tendon at the proximal musculotendinous junction of the biceps femoris long head: a case report. *Knee Surg Sports Traumatol Arthrosc* 2008;16(8):797-802.
46. Brukner P, Connell D. 'Serious thigh muscle strains': beware the intramuscular tendon which plays an important role in difficult hamstring and quadriceps muscle strains. *Br J Sports Med* 2016;50(4):205-8.
47. Malliaropoulos N, Papacostas E, Kiritsi O et al. Posterior thigh muscle injuries in elite track and field athletes. *Am J Sports Med* 2010;38(9):1813-19.
48. Cibulka MT, Rose SJ, Delitto A et al. Hamstring muscle strain treated by mobilising the sacroiliac joint. *Phys Ther* 1986;66(8):1220-3.
49. Malliaropoulos N, Papalexandris S, Papalada A et al. The role of stretching in rehabilitation of hamstring injuries: 80 athletes follow-up. *Med Sci Sports Exerc* 2004;36(5):756-9.
50. Sherry MA, Best TM. A comparison of 2 rehabilitation programs in the treatment of acute hamstring strains. *J Orthop Sports Phys Ther* 2004;34(3):116-25.
51. Petersen J, Hölmich P. Evidence-based prevention of hamstring injuries in sport. *Br J Sports Med* 2005;39(6):319-23.
52. Quintero AJ, Wright VJ, Fu FH et al. Stem cells for the treatment of skeletal muscle injury. *Clin Sports Med* 2009;28(1):1-11.
53. Khan KM, Scott AR. Mechanotherapy: how physical therapists' prescription of exercise promotes tissue repair. *Br J Sports Med* 2009;43(4):247-52.
54. Robinson M, Hamilton B. Medical interventions in the management of hamstring muscle injury. *Eur J Sport Sci* 2014;14(7):743-51.
55. Reynolds JF, Noakes TD, Schwellnus MP et al. Non-steroidal anti-inflammatory drugs fail to enhance healing of acute hamstring injuries treated with physiotherapy. *S Afr Med J* 1995;85(6):517-22.
56. Levine WN, Bergfeld JA, Tessendorf W et al. Intramuscular corticosteroid injection for hamstring injuries. A 13-year experience in the National Football League. *Am J Sports Med* 2000;28(3):297-300.
57. A Hamid MS, Mohamed Ali MR, Yusof A et al. Platelet-rich plasma injections for the treatment of hamstring injuries: a randomised controlled trial. *Am J Sports Med* 2014;42(10):2410-8.
58. Reurink G, Goudswaard GJ, Moen MH et al. Platelet-rich plasma injections in acute muscle injury. *N Engl J Med* 2014;370(26):2546-7.
59. Hamilton B, Tol JL, Almusa E et al. Platelet-rich plasma does not enhance return to play in hamstring injuries: a randomised controlled trial. *Br J Sports Med* 2015;49:943-50.
60. Martinez-Zapata MJ, Orozco L, Balias R et al. Efficacy of autologous platelet-rich plasma for the treatment of muscle rupture with haematoma: a multicentre, randomised, double-blind, placebo-controlled clinical trial. *Blood Transfus* 2016;14(2):245-54.
61. Linklater JM, Hamilton B, Carmichael J et al. Hamstring injuries: anatomy, imaging and intervention. *Semin Musculoskelet Radiol* 2010;14(2):131-61.
62. Orchard JW, Best TM, Mueller-Wohlfahrt HW et al. The early management of muscle strains in the elite athlete: best practice in a world with a limited evidence basis. *Br J Sports Med* 2008;42(3):158-9.
63. Lee P, Rattenberry A, Connelly S et al. Our experience on Actovegin, is it cutting edge? *Int J Sports Med* 2011;32(4):237-41.
64. Chumanov ES, Heiderscheit BC, Thelen DG. The effect of speed and influence of individual muscles on hamstring mechanics during the swing phase of sprinting. *J Biomech* 2007;40(16):3555-62.
65. Gabbe BJ, Bennell KL, Finch CF. Why are older Australian football players at greater risk of hamstring injury? *J Sci Med Sport* 2006;9(4):327-33.
66. Mendiguchia J, Garrues MA, Cronin JB et al. Nonuniform changes in MRI measurements of the thigh muscles after two hamstring strengthening exercises. *J Strength Cond Res* 2013;27(3):574-81.
67. Zebis MK, Skotte J, Andersen CH et al. Kettlebell swing targets semitendinosus and supine leg curl targets biceps femoris: an EMG study with rehabilitation implications. *Br J Sports Med* 2013;47(18):1192-8.
68. McAllister MJ, Hammond KG, Schilling BK et al. Muscle activation during various hamstring exercises. *J Strength Cond Res* 2014;28(6):1573-80.
69. Schache AG, Dorn TW, Blanch PD et al. Mechanics of the human hamstring muscles during sprinting. *Med Sci Sports Exerc* 2012;44(4):647-58.
70. Brockett CL, Morgan DL, Proske U. Human hamstring muscles adapt to eccentric exercise by changing optimum length. *Med Sci Sports Exerc* 2001;33(5):783-90.
71. Proske U, Morgan DL, Brockett CL et al. Identifying athletes at risk of hamstring strains and how to protect them. *Clin Exp Pharmacol Physiol* 2004;31(8):546-50.
72. Aagaard P, Simonsen EB, Trolle M et al. Specificity of training velocity and training load on gains in isokinetic knee joint strength. *Acta Physiol Scand* 1996;156(2):123-9.
73. Seger JY, Thorstensson A. Effects of eccentric versus concentric training on thigh muscle strength and EMG. *Int J Sports Med* 2005;26(1):45-52.
74. Mjolsnes R, Arnason A, Osthaugen T et al. A 10-week randomised trial comparing eccentric vs. concentric hamstring strength training in well-trained soccer players. *Scand J Med Sci Sports* 2004;14(5):311-17.
75. Arnason A, Andersen TE, Holme I et al. Prevention of hamstring strains in elite soccer: an intervention study. *Scand J Med Sci Sports* 2008;18(1):40-8.
76. Petersen J, Thorborg K, Nielsen MB et al. Preventive effect of eccentric training on acute hamstring injuries in men's

- soccer: a cluster-randomised controlled trial. *Am J Sports Med* 2011;39(11):2296-303.
77. van der Horst N, Smits DW, Petersen J et al. The preventative effect of the Nordic hamstring exercise on hamstring injuries in amateur soccer players: a randomized controlled trial. *Am J Sports Med* 2015;43(6):1316-1323.
 78. Lieberman DE, Raichlen DA, Pontzer H et al. The human gluteus maximus and its role in running. *J Exp Biol* 2006;209(Pt 11): 2143-55.
 79. Mendiguchia J, Brughelli M. A return-to-sport algorithm for acute hamstring injuries. *Phys Ther Sport* 2011;12(1):2-14.
 80. Nemeth G, Ohlsén H. In vivo moment arm lengths for hip extensor muscles at different angles of hip flexion. *J Biomech* 1985;18(2):129-40.
 81. Contreras B, Vigotsky AD, Schoenfeld BJ et al. A comparison of gluteus maximus, biceps femoris, and vastus lateralis EMG activity in the back squat and barbell hip thrust exercises. *J App Biomech* 2015;31(6):452-8.
 82. Brughelli M, Cronin J. Preventing hamstring injuries in sport. *Strength Cond J* 2008;30(1):55-64.
 83. Hides JA, Stanton WR, Mendis MD et al. Effect of motor control training on muscle size and football games missed from injury. *Med Sci Sports Exerc* 2012;44(6):1141-9.
 84. Hides JA, Stanton WR. Can motor control training lower the risk of injury for professional football players? *Med Sci Sports Exerc* 2014;46(4):762-8.
 85. Cameron ML, Adams RD, Maher CG et al. Effect of the HamSprint Drills training programme on lower limb neuromuscular control in Australian football players. *J Sci Med Sport* 2009;12(1):24-30.
 86. Cameron M, Adams R, Maher C. Motor control and strength as predictors of hamstring injury in elite players of Australian football. *Phys Ther Sport* 2003;4(4):159-66.
 87. Verrall GM, Slavotinek JP, Barnes PG. The effect of sports specific training on reducing the incidence of hamstring injuries in professional Australian Rules football players. *Br J Sports Med* 2005;39(6):363-8.
 88. Opar DA, Piatkowski T, Williams MD et al. A novel device using the Nordic hamstring exercise to assess eccentric knee flexor strength: a reliability and retrospective injury study. *J Orthop Sports Phys Ther* 2013;43(9):636-40.
 89. Walden M, Häggglund M, Ekstrand J. UEFA Champions League study: a prospective study of injuries in professional football during the 2001-2002 season. *Br J Sports Med* 2005;39(8):542-6.
 90. Henderson G, Barnes CA, Portas MD. Factors associated with increased propensity for hamstring injury in English Premier League soccer players. *J Sci Med Sport* 2010;13(4):397-402.
 91. Witvrouw E, Danneels L, Asselman P et al. Muscle flexibility as a risk factor for developing muscle injuries in male professional soccer players. A prospective study. *Am J Sports Med* 2003;31(1):41-6.
 92. Whiteley R, Jacobsen P, Prior S et al. Correlation of isokinetic and novel hand-held dynamometry measures of knee flexion and extension strength testing. *J Sci Med Sport* 2012;15(5): 444-50.
 93. Croisier JL, Ganteaume S, Binet J et al. Strength imbalances and prevention of hamstring injury in professional soccer players: a prospective study. *Am J Sports Med* 2008;36(8):1469-75.
 94. Tol JL, Hamilton B, Eirale C et al. At return to play following hamstring injury the majority of professional football players have residual isokinetic deficits. *Br J Sports Med* 2014;48(18):1364-9.
 95. Freckleton G, Cook J, Pizzari T. The predictive validity of a single leg bridge test for hamstring injuries in Australian Rules Football Players. *Br J Sports Med* 2014;48(8):713-7.
 96. Askling CM, Nilsson J, Thorstensson A. A new hamstring test to complement the common clinical examination before return to sport after injury. *Knee Surg Sports Traumatol Arthrosc* 2010;18(12):1798-803.
 97. Sanfilippo JL, Silder A, Sherry MA et al. Hamstring strength and morphology progression after return to sport from injury. *Med Sci Sports Exerc* 2013;45(3):448-54.
 98. Reurink G, Goudswaard GJ, Tol JL et al. MRI observations at return to play of clinically recovered hamstring injuries. *Br J Sports Med* 2014;48(18):1370-6.
 99. Orchard J, Best TM. The management of muscle strain injuries: an early return versus the risk of recurrence. *Clin J Sport Med* 2002;12(1):3-5.
 100. Freckleton G, Pizzari T. Risk factors for hamstring muscle strain injury in sport: a systematic review and meta-analysis. *Br J Sports Med* 2013;47(6):351-8.
 101. van Beijsterveldt AM, van de Port IG, Vereijken AJ et al. Risk factors for hamstring injuries in male soccer players: a systematic review of prospective studies. *Scand J Med Sci Sports* 2013;23(3):253-62.
 102. Arnason A, Sigurdsson SB, Gudmundsson A et al. Risk factors for injuries in football. *Am J Sports Med* 2004;32(1 Suppl): 5S-16S.
 103. Engebretsen AH, Myklebust G, Holme I et al. Intrinsic risk factors for hamstring injuries among male soccer players: a prospective cohort study. *Am J Sports Med* 2010;38(6):1147-53.
 104. Orchard JW. Intrinsic and extrinsic risk factors for muscle strains in Australian football. *Am J Sports Med* 2001;29(3): 300-3.
 105. Verrall GM, Slavotinek JP, Barnes PG et al. Clinical risk factors for hamstring muscle strain injury: a prospective study with correlation of injury by magnetic resonance imaging. *Br J Sports Med* 2001;35(6):435-9; discussion 440.
 106. Orchard JW, Farhart P, Leopold C. Lumbar spine region pathology and hamstring and calf injuries in athletes: is there a connection? *Br J Sports Med* 2004;38(4):502-4; discussion 504.
 107. Lee MJ, Reid SL, Elliott BC et al. Running biomechanics and lower limb strength associated with prior hamstring injury. *Med Sci Sports Exerc* 2009;41(10):1942-51.