

Observation-conditioned bounds define experimental decision criteria for bifunctional enzyme systems

Yule Choi^{1*}

¹ Korea Science Academy of KAIST, Busan, Republic of Korea

* 25-117@ksa.hs.kr

Short title: Decision criteria for bifunctional enzyme systems

Abstract

Distinct biochemical mechanisms can reproduce the same measured response while disagreeing about unobserved states. We develop an observation-conditioned framework that derives sufficient conditions for experimental decisions under this ambiguity. Constructive models of bacterial nitrogen signaling preserve complete stationary target distributions while changing enzyme occupancy and kinetics. Protein conservation and a concurrently checked modification-count constraint then bound differences between compatible mechanisms. In a prospective GlnE assay, the count constraint reduces a total-target difference bound from 0.370 to 0.041, below a prescribed 0.05 tolerance. This conclusion requires the stated topology, stationarity, coefficient units and matched observation; it is not an experimentally validated concentration choice. A general graph construction extends the method to positive-polynomial stationary weights without requiring enzyme-concentration cancellation. In the independent AceK–isocitrate dehydrogenase system, it bounds a network with two-enzyme-bound target complexes and identifies another prospective observation gate. Six published kinetic summaries also admit distinct substrate-binding realizations with nearly identical initial rates but different protein occupancies; withheld ATPase measurements test additional biological implications. Finally, raw-assay simulations compare shared and condition-specific signal fits using 3,000 calibration trials per null setting and 2,000 independent trials per evaluation setting, including nonlinear halfpoint estimation. Condition-specific fits remove tested affine signal biases, whereas continuous affinity mixtures still alter mechanistic interpretation. The framework connects explicitly delimited mechanism classes to measurable certification conditions and complementary observations, rather than selecting an unrestricted native mechanism.

Author summary

27

Matching a biological response does not necessarily reveal how the underlying proteins work. We study
enzymes that both add and remove chemical groups from another protein. Different binding and reaction
mechanisms can produce the same response, even when every modification state is measured. We show how
additional observations and known protein amounts can place limits on the predictions shared by these
mechanisms. The resulting decision rules are conditional: the required measurements must be checked in the
proposed experiment. We develop a general calculation and apply it to bacterial nitrogen signaling and a
separate enzyme system controlling carbon metabolism. We also follow simulated binding experiments from
prepared concentrations to statistical interpretation. Allowing each experimental condition to have its own
background and signal scale corrects tested measurement distortions, but mixtures of hidden binding states
remain misleading. These results identify which assumptions an experiment must check before a good fit can
support a mechanistic conclusion.

38

Introduction

Ultrasensitivity describes a sharp input–output response, but the shape of that response does not uniquely identify its molecular source. Enzyme saturation, interactions among modification sites, reciprocal regulation, and downstream recognition can produce similar titrations [1,2]. A useful mechanistic analysis must therefore answer two questions: which response properties follow from the reaction architecture, and which observations distinguish competing explanations of the same measured response?

The *Escherichia coli* nitrogen-signaling cascade provides a concrete setting for both questions. Glutamine regulates the bifunctional enzyme GlnD, which adds and removes UMP groups on trimeric PII. PII modification states regulate GlnE, which adds and removes adenylyl groups on the twelve-subunit glutamine synthetase (GS) complex [3–5]. Both layers combine multisite targets with opposing activities of a shared enzyme. Fig 1 summarizes the regulatory directions and state classes.




Fig 1. The GlnD–PII–GlnE–GS nitrogen-signaling cascade. Low free glutamine favors the UTase activity of GlnD, increasing PII uridylylation; uridylylated PII favors the adenylyl-removing (AR) activity of GlnE, driving GS toward lower adenylylation. High free glutamine favors the UR activity of GlnD, increasing unmodified PII; unmodified PII favors GlnE ATase, driving GS toward higher adenylylation. PII trimers carry 0–3 UMP groups, and GS dodecamers carry 0–12 adenylyl groups. The schematic summarizes regulatory direction and state classes rather than a kinetic model.

Published measurements report a glutamine response coefficient near two for the GlnD–PII cycle and approximately 4.5–6.5 for the full cascade, depending on PII concentration [6,7]. Proposed explanations

include allosteric regulation and bifunctionality [8,9], zero-order and multistep effects [7], and selective GlnE activation by fully unmodified PII [6]. Bifunctionality can suppress sensitivity in some architectures but permit it through sequestration or additional enzyme–substrate complexes in others [10–13]. Reconstituted-cycle studies also show that concentration robustness depends on strong glutamine inhibition of UTase and on the enzyme construct [14]. These alternatives make the observation rule and assay conditions central to any quantitative attribution.

We ask which experimentally checkable conditions permit a decision despite mechanistic ambiguity. Three results connect that question to practice: observation-conditioned bounds for a declared mechanism class, a general graph construction applicable beyond enzyme-concentration cancellation, and a calibrated binding benchmark that distinguishes correctable signal errors from hidden-state ambiguity. The nitrogen cascade supplies a detailed concentration example; AceK–isocitrate dehydrogenase supplies an independent biochemical topology and published kinetic summaries. Stationary elimination, scaling non-identifiability, prediction profile likelihood and model-discrimination design are established foundations [15–22]. Our contribution is a conservation-aware construction that turns bounded observations into sufficient class-wide decision criteria, together with explicit tests of the observation process and the biochemical restrictions needed to use them.

Results

Complete target distributions can leave enzyme mechanisms unresolved

The mathematical foundation is a stationary modification ladder. For a distributive target with n modification sites, a shared bifunctional enzyme, and regulatory exchange restricted to free enzyme, elimination of catalytic complexes gives

$$p_i = \frac{c_i u^i}{\sum_{j=0}^n c_j u^j}, \quad \theta = \frac{1}{n} \sum_i i p_i.$$

Here u is the ratio of opposing catalytic drives and $c_i > 0$ describes the effective ladder. This reduction uses established stationary elimination and rational parameterization [15–18]. Differentiation gives $d\theta/d\log u = \text{Var}(i)/n$. Consequently, local sensitivity separates into a ladder factor and switch elasticity. Independent identical sites give $c_i = \binom{n}{i}$ and $\theta = u/(1+u)$ for every n . Neither site count nor enzyme saturation alone steepens that free-target response. S1 Appendix derives the result and its restrictions; S1–S2 Figs retain the concentration and estimator comparisons.

This distinction matters for the twelve digitized GlnD–PII measurements [6]. The bounded fit has exponent 2.018 but a physical-occupancy midpoint local coefficient of 1.725. Those are different summaries of the same curve, not interchangeable measurements of cooperativity. Independent and interacting ladders fitted with the same four-parameter switch give RMSEs of 0.0352–0.0392 UMP groups per trimer. Exact inverse constructions can also preserve the mean while changing state fractions or introducing a refractory population. Downstream gain calibration can preserve some of those ambiguities. The titration, plateau, shared-parameter and downstream comparisons are retained in S3–S4 Figs, S2 Appendix and the repository extended-analysis archive; they compare stated models, not experimental probabilities of their native correctness.

A stronger ambiguity survives measurement of every target state. GlnD domain experiments motivate separate catalytic and regulatory functions [23–25], without establishing a unique assignment of model states to ACT domains. Let three free enzyme states have weights $w = (1, G/K_1, G^2/(K_1K_2))$ and UT/UR catalytic specificities a_j, b_j . The stationary target drive depends on A/B , where $A = \sum_j a_j w_j$ and $B = \sum_j b_j w_j$. The transformation

$$K_1 \mapsto \lambda K_1, \quad K_2 \mapsto K_2/\lambda, \quad (a_1, b_1) \mapsto \lambda(a_1, b_1)$$

preserves both numerators: the singly liganded weight and its specificity compensate exactly. It therefore preserves every free PII modification-state fraction at every free glutamine concentration, not merely a fitted mean (Fig 2).

A reference independent-ligand model fitted to the twelve coordinates has RMSE 0.0366 UMP groups per trimer and binding scale $K = 0.0520$ mM. Its transported UR specificity-fold scenario is stated in A5–A6 and D4; these coefficients are not independently measured microscopic rates. Transformations with $\lambda = 1, 2, 4$ preserve monotonically decreasing UT and increasing UR capacities but predict singly liganded fractions $1/2, 1/3, 1/5$ at $G = K$. Under rapid binding and unsaturated conversion, their relative relaxation rates are $1, 4/3, 8/5$. Average ligand occupancy at that input remains one for all three: measuring an average at a symmetry point can miss the very difference revealed by a resolved enzyme state.

Good titration agreement is not sufficient biochemical validation. The fixed- C construction transports two endpoint specificity ratios and a 0.080 mM UR half-range, giving RMSE 0.540. Relaxing its fixed- C , intermediate-state and half-range constraints, while retaining both endpoint ratios and monotone state activities, gives RMSE 0.0409. Its UR half-range is 0.2566 mM and a binding scale reaches the 148 mM fit boundary. The improvement cannot be attributed to a single relaxation. A fixed- K_1 profile shows the




Fig 2. GlnD mechanisms with identical PII states predict different enzyme occupancies and kinetics. (A) Three finite binding-and-catalysis constructions preserve the same fitted mean and every binomial PII state fraction. Points are the digitized coordinates. (B) Singly liganded enzyme fractions differ, reaching 0.5, 0.333, and 0.2 at the common binding scale. (C) PII relaxation-rate ratios in the rapid-binding, unsaturated conversion reduction; both capacities share the same multiplier. (D) An illustrative glutamine decrease from 0.5 to 0.05 mM gives different PII time courses with the same endpoint. Time is in reference-model units, not experimentally estimated minutes. The baseline is fitted once; the other models are obtained by the exact transformation in A6.

consequence for design: from $K_1 = 0.3$ to 10 mM, RMSE improves from 0.0493 to 0.0411 while the maximum
singly liganded occupancy contrast within the monotone transformed family falls from 0.242 to 0.0108 (S8
Fig B). These are cross-assay stress hypotheses, not confidence intervals. The 1998 UT and UR assays use 25
and 50 mM MgCl_2 , respectively, compared with 10 mM in the 2011 cascade assay [6,24]. D4 documents the
transfer conditions. Equivalence persists in this profile, but the original occupancy gap is not a robust
biochemical prediction.

The published six-state GlnE topology supplies a downstream counterpart [26]. With $g = G/K_G$,
 $p = P/K_P$ and $u = U/K_U$, its binding polynomial is $Z_E = 1 + g + p + u + pg/\alpha_1 + pu/\alpha_2$. The stationary
ATase/AR drive is

$$v = \frac{k_P}{k_R} \frac{p + \beta g + \beta' pg/\alpha_1}{u}.$$

The common binding denominator cancels. Thus α_2 is silent in this ratio, while common scaling of α_1 and β'
trades binding synergy against catalysis without changing it. GlnE variants coupled to the GlnD family can
preserve the GS response without condition-specific recalibration. They predict different PII-binding
halfpoints and reduced relaxation rates (S5 Fig). The direct glutamine term βg is retained: the cascade
response is not attributable to PII alone.

These are constructive applications of binding-polynomial and scaling ideas [20,21,27], not a claim that
stationary invariants or non-identifiability are new. Their specific contribution is an explicit biochemical
family that defeats increasingly resolved target measurements. A state-weight change $w_j \mapsto s_j w_j$,
compensated by $k_j \mapsto k_j/s_j$, preserves the corresponding catalytic numerator under the stipulated topology.
A15 of S1 Appendix compares each result with prior theory. The construction neither exhausts equivalent
mechanisms nor proves uniqueness of a native interpretation.

A concurrent observation defines a sufficient concentration criterion

Experiments often count enzyme-bound target, whereas the invariant concerns free target. Finite
mass-action binding need not destroy the stationary identity: complex balances recover the same free ratio
when regulatory exchange is restricted to free enzyme. Total fractions can nevertheless differ. With one
target per enzyme complex, two mechanisms sharing the free fraction and conserved totals satisfy

$$|\Delta\theta_{\text{total}}| \leq \min(1, E_T/T_T).$$

This is a conservation bound, not a low-occupancy approximation or a Hill-slope bound. Finite binding
changes transients; regulatory exchange on catalytic complexes can change the stationary identity itself. S6

Fig separates these effects.

We then retain four PII states, thirteen GS states, three GlnD regulatory states and a refinement of the GlnE classes in a 120-species mass-action cascade. Catalytic complexes conserve PII, GS and both enzymes. Partly modified PII recognition is specified and varied as a model assumption; glutamine and nucleotide pools are buffered. Let P_T, S_T, D_T, E_T denote total PII trimers, GS dodecamers, GlnD and GlnE. Define

$$\epsilon_P = (D_T + 2E_T)/P_T < 1, \quad \epsilon_S = \min(1, E_T/S_T).$$

One GlnD can bind at most one PII and one GlnE at most two. The total-variation distance between total PII distributions in the equivalent pair is therefore at most ϵ_P , while their free PII distributions coincide.

Downstream, the specified GlnE drive has the form $r(T) = a + b/T$ for free PII pool size T , with $a, b \geq 0$. The direct glutamine route supplies b . Combining the permitted free-pool range with a ladder sensitivity bound gives

$$|\Delta\theta_{GS,\text{total}}| \leq \min \left\{ 1, \tanh \left[\frac{\nu_*}{4} \log \frac{1}{1 - \epsilon_P} \right] + \epsilon_S \right\}.$$

Independent GS sites permit $\nu_* = 1$; $\nu_* = 12$ covers any positive twelve-site ladder shared by the pair.

Without the direct route, $b = 0$, free GS fractions coincide and only ϵ_S remains. These conclusions require the stated recognition and reaction structure (A10–A11).

The general bound can be very conservative: a small difference in one fitted realization does not establish a uniform interpretation guarantee. A concurrent observation can tighten it. Suppose the total PII mean UMP count has a justified upper bound M . The free zero-UMP fraction satisfies

$$\pi_0 \geq p_{\min} = \max\{0, 1 - M/(1 - \epsilon_P)\}.$$

This follows from nonnegative modification in the bound pool and requires no independent PII-site assumption. At fixed glutamine the source GlnE drive can be written $r(T) = k(a + b/T)$. A common positive factor k , including the AR gain, cancels from the drive contrast. With $g = G/K_G$, define

$$\ell = \frac{b}{a} \leq \frac{\beta g K_P}{p_{\min}(1 + \beta' g/\alpha_1)}, \quad d = \log \frac{1 + \ell/[P_T(1 - \epsilon_P)]}{1 + \ell/P_T}.$$

Then $|\Delta\theta_{GS,\text{total}}| \leq \min\{1, \epsilon_S + \tanh(\nu_* d/4)\}$. If $p_{\min} = 0$, the generic bound is recovered as $\ell \rightarrow \infty$. A11 proves the result and specifies its reaction assumptions.

Consider a prospective assay at $P_T = 5$, $S_T = 0.5$, $D_T = 0.5$, $E_T = 0.02 \mu\text{M}$ and $G = 2 \text{ mM}$, with a

concurrently verified count bound $M \leq 0.67$. The source GlnE fit gives $K_G = 15.6$ mM, $K_P = 0.18$ μ M, $\alpha_1 = 0.17$, $\beta = 175/7.5$ and $\beta' = 1315/7.5$ [26]. For every positive twelve-site GS ladder shared by the pair, the generic bound is 0.370, while the count-conditioned bound is 0.0412. If the count gate is confirmed in that assay, the resulting bound crosses the prescribed 0.05 tolerance (Fig 3A). At $E_T = 0.10$ μ M, the GS sequestration allowance alone is 0.20; a fivefold reduction in GlnE concentration, without also reducing GlnD, is sufficient under the stated coefficient and observation conditions. This is a pairwise difference guarantee within the stated model class, not an error bound relative to biological truth or a globally optimal assay.

The numerical gate is motivated by the historical digitized mean 0.65 at 2 mM, but is not a confidence limit inferred from it. It must be measured in the proposed assay. Likewise, the source coefficients are a fitted example rather than independent microscopic measurements. At the proposed totals, an independently justified $\ell \leq 0.1417$ μ M would suffice (rounded down); the nominal value is 0.0162. Using an apparent-affinity sensitivity scale $K_P = 1$ μ M gives a bound of 0.0464. The source supplementary legend instead prints $K_P = 0.18$ mM, inconsistent with the main text and its stated direct-scale comparison; literal use gives 0.300 and loses the guarantee. Fig 3B shows this dependence explicitly. If the observation or coefficient bounds cannot be justified, the general bound must be used.

Five thousand arbitrary positive ladders and 5,000 free/bound PII mixtures verify the component inequalities. An additional 120 paired full-reaction realizations, using the source GlnE rate ratios and varied unknown gains, lie below the bound (Fig 3C); maximum stationary vector-field residual is 1.23×10^{-15} . These are finite-model checks, not validation of native kinetics or equilibration times. The earlier 8,200-condition sweep and topology checks remain in A10–A11 and S1 Data.

The boundary depends on topology and input control. Allowing the previously silent PII–PII-UMP GlnE class to catalyze adenylation makes its interaction parameter observable; the selected extension reaches a GS fraction separation of 0.154 (Fig 3D). Yet the compensated family retains a drive $a + b/T + cT$ with log-elasticity magnitude at most one, so the general fraction bound survives. Fixing total instead of free glutamine is another intervention: different enzyme occupancies can produce different free inputs, with $0 \leq G_T - G \leq 2D_T + E_T$ in this network. These examples identify assumptions whose alteration changes the inference; they do not establish an additional native catalytic route.




Fig 3. A prospective PII count gate defines a GlnE concentration criterion. (A) General and count-conditioned upper bounds on pairwise total GS fraction difference versus GlnE total. At the fivefold-reduced $E_T = 0.02 \mu\text{M}$, the count-conditioned bound is 0.0412 and crosses the prescribed 0.05 tolerance; the general bound is 0.370. Other fixed conditions are $P_T = 5$, $S_T = 0.5$, $D_T = 0.5 \mu\text{M}$, $G = 2$ mM, $M \leq 0.67$, and the source-fit GlnE coefficients. (B) Bound at $E_T = 0.02 \mu\text{M}$ versus the count upper limit and K_P ; the contour marks the 0.05 decision boundary. Markers identify the main-text source unit, an apparent-scale sensitivity scenario and the conflicting literal supplementary unit. (C) Actual differences and their general/count-conditioned bounds in 120 paired 120-species realizations; GlnD $\lambda = 1$ versus 4, GlnE $\sigma = 1$, three glutamine doses, two K_P values, five gains and four GlnE totals. Each bound uses the larger total PII count of the pair. These selected checks need not saturate a uniform bound. (D) Topology control: allowing EPU to catalyze adenylation makes α_2 observable. Curves compare $\alpha_2 = 2.17$ and 8.85, maximizing over 41 specified doses at each added productive coefficient. All panels are conditional model calculations.

End-to-end observation models define when complementary measurements discriminate mechanisms

The next measurement should target what remains unresolved. State-resolved PII distinguishes the stipulated ladder and population alternatives. The adjacent-state contrast

$$\frac{p_{i-1}p_{i+1}}{p_i^2} = \frac{c_{i-1}c_{i+1}}{c_i^2}$$

is the established distributive-modification invariant [28]; here its input dependence tests a fixed homogeneous ladder. Rare-state ratios are error-sensitive. For four mean-matched candidates, the minimum pairwise state-distribution distance is largest at 0.747 mM on the disclosed grid. A 10 mM measurement separates the independent and refractory scenarios' predicted fully modified fractions (S4 Fig). These are candidate-dependent designs, not experimentally established optima.

Such state measurements cannot separate the GlnD family. In the illustrative independent-binding family, a target-free measurement resolving singly liganded GlnD gives a minimum gap of 0.244 under a factor-of-two binding-scale variation. The kinetic-constraint profile above shows why this gap cannot be transported to every acceptable fit. Moreover, the assay must distinguish zero, one and two bound glutamines, rather than their mean. Intact native MS could be evaluated in a pilot, but successful GlnD state resolution and precision are not established here. The central observation benchmark therefore uses GlnE binding curves. Table 1 distinguishes the evidence and observations needed for these decisions.

With PII-UMP absent, normalized GlnE PII-binding halfpoints obey

$$\frac{H(G)}{H(0)} = \frac{1 + G/K_G}{1 + G/(\alpha_1 K_G)}.$$

The unknown PII affinity cancels. At appreciable enzyme concentration, a total-PII halfpoint equals $H(G) + E_T/2$ in the single-binding-class model. Two nonzero glutamine conditions plus a shared zero-glutamine curve identify K_G and α_1 ideally, except at $\alpha_1 = 1$ (D5). A solution assay proportional to bound GlnE fraction would require free and saturating-ligand controls, concentration standards and checks for glutamine-dependent signal changes.

We test the complete observation process at $E_T = 0.04 \mu\text{M}$: eleven total-PII doses, three glutamine concentrations and two technical readings per dose (S8 Fig A). Each trial contains eight independently generated complete titration sets. Finite-depletion fits estimate three halfpoints with either shared background and gain or separate background and gain for each glutamine condition. The average

Table 1. Evidence, observables and conditions for the central decisions.

Evidence class	Quantity supplied	What the inference additionally requires
Published observations	Twelve mean-UMP coordinates; source assay compositions and fitted GlnE coefficients	Cross-assay transfer is not automatic; rounded coordinates supply no biological confidence interval.
Exact conditional result	Free-state equivalence; count-conditioned total-GS bound	Stated reaction topology, common free input, conserved protein units and an upper coefficient bound.
Prospective observation	Total PII count M in the proposed cascade assay	Concurrent calibration; $M \leq 0.67$ is a gate to check, not an observed confidence limit.
Synthetic performance	Raw GlnE binding curves through halfpoint estimation and testing	Proportional binding readout, concentration controls and a tested binding-state model.
Second-system calibration	Six AceK kinetic summaries and a published dimeric-target network	Matched assay conditions; reconstructed curves are not raw observations.
Independent-activity check	Three ATPase halfpoints excluded from kinase calibration	Additional biochemical assumptions; exact errors and covariance are unavailable.
Prospective extension	Paired-paralog calibration and sequestration intervals (S7 Fig)	Matched calibration transfer and independent direct binding validation.

log-halfpoint-ratio score retains the shared zero-glutamine measurement within each experiment. The reference is $\alpha_1 = 0.17$, with $K_G \in [7.8, 15.6]$ mM. This source-based sensitivity range is narrower than the earlier illustrative $[7.8, 31.2]$ mM range (D5); neither is a confidence region.

For each fitting strategy, fixed-null parametric calibration uses 3,000 raw-simulation and nonlinear-refitting trials at each of 27 nuisance settings. A binomial-tolerance order statistic, adjusted across the grid, sets the largest null cutoff. Independent evaluation uses 2,000 trials at each of 210 settings spanning K_G , concentration and signal errors, regulatory effects, dose biases and eleven mixture strengths (E5 in S3 Appendix). The 8,016,000 nonlinear fits include both strategies, which use paired synthetic observations. The calibrated cutoffs are 2.0625 and 2.1036; the largest independent null rejection estimates are 0.0415 and 0.0385 (Fig 4A). The calibration statement applies to the finite declared grid, not every intermediate nuisance value or a native assay. Small-effect power remains sensitive to the unknown K_G (Fig 4B); S9 Fig shows the complete null and small-effect nuisance grid.

Condition-specific signal fitting changes the experimental interpretation. At $K_G = 15.6$ mM, a nonzero-glutamine background shift of -0.02 produces reference rejection 0.924 with a shared signal fit and

0.027 with condition-specific fits (Fig 4C). A log-gain shift of -0.05 gives 0.992 versus 0.0295. Thus
 matching the signal model to condition-specific controls can repair these tested distortions. Optimizer
 failures and active-bound fits are counted separately: no optimizer failures occurred in evaluation, while 17
 shared-fit and 22 condition-specific-fit solutions reached a parameter bound. Their trials remain in the
 denominator and do not reject.

An affinity mixture remains a distinct limitation. Along a continuous path from one binding class to two,
 a mixture strength of 0.2 gives rejection 0.567 even with condition-specific signal fits, and a strength of 0.5
 gives 1.0 at $K_G = 15.6$ mM (Fig 4D). Residuals remain close to the single-class noise scale. Such rejection
 challenges the joint binding and calibration hypothesis; it does not identify a change in α_1 . Concentration
 controls and condition-specific signal fits therefore address some errors, while heterogeneous binding requires
 a different observation or model.

A general graph enclosure extends the workflow beyond enzyme cancellation

The cancellation of a shared enzyme is a property to test, not a consequence of bifunctionality alone. Fix free
 ligands and a free enzyme concentration e . Suppose each complex contains one target molecule, possibly an
 oligomer, and target-state transitions form an irreducible linear generator $Q(e)$ with nonnegative polynomial
 edge rates. The established matrix-tree theorem gives positive stationary weights $\tau_i(e)$ [29]. For the unbound
 target states, write $\tau_F(e) = C(1, e, \dots, e^n)^T$. Their normalized distribution is independent of e exactly when
 C has rank one. This criterion follows from polynomial proportionality and provides an explicit check before
 applying a cancellation-based interpretation.

When cancellation fails, the same weights yield an enclosure. Express them in the Bernstein basis on a
 justified interval $e \in [L, U]$, and normalize each nonzero coefficient column to form a vertex of V . The
 free-state distribution lies in the convex hull of these vertices (E1). With free-pool weights $\lambda \geq 0$ and
 bound-state masses $z \geq 0$ aggregated into the same target-state categories, conservation gives $\mathbf{1}^T \lambda + \mathbf{1}^T z = 1$
 and $\mathbf{1}^T z \leq \epsilon$. A matched observation imposes

$$a \leq A(V\lambda + z) \leq b. \quad (1)$$

Minimizing and maximizing a reporter $h^T(V\lambda + z)$ are linear programs. Their interval contains every
 compatible stationary mechanism in the declared class; its width bounds pairwise disagreement. An empty
 feasible set indicates incompatibility, not a successful certificate. The enclosure may be conservative, and
 biological observation intervals or kinetic coefficients must be justified separately. E1–E2 supply the proof,




Fig 4. Raw-assay calibration separates correctable signal errors from mechanistic ambiguity.

(A) Independent null rejection at 27 settings per fitting strategy; each point summarizes 2,000 trials, with nine error settings at each K_G . Cutoffs use separate parametric null simulations including nonlinear halfpoint fitting. (B) Condition-specific-fit rejection versus α_1 ; lines are means and shading is the range across nine error settings at each K_G , not a confidence band. (C) A glutamine-dependent background shift causes strong rejection under a shared signal model; separate background and gain parameters correct the tested affine distortion. (D) A continuous hidden-affinity-mixture path still alters the single-class interpretation. C–D fix $K_G = 15.6$ mM; shading shows 95% Wilson Monte Carlo intervals. Dashed lines mark 0.05. Each trial contains eight complete titrations. Error magnitudes are specified scenarios, and all panels are simulations.

reusable graph algorithm and independent checks.

AceK, which phosphorylates and dephosphorylates isocitrate dehydrogenase (IDH), provides an independent application. The published dimeric-target network allows two AceK molecules to bind one IDH dimer [30]. Its free-state weights reduce to three positive quadratic polynomials with coefficient rank three: enzyme cancellation fails, but the enclosure remains available. At prospective totals of $1\ \mu\text{M}$ IDH dimers and $0.05\ \mu\text{M}$ AceK, its unphosphorylated-dimer fraction has class interval width 0.056995. A concurrently verified mean phosphate-count interval $[1.435, 1.475]$ would reduce that width to 0.036257, below 0.05 (Fig 5D). These are proposed gate limits under source coefficients and a constructed bound-state dwell-time class. They are not existing experimental measurements. Full conservation checks and independent integration retain singly and doubly bound complexes (E2).

Published AceK kinetics also test the choice of complementary observation. Six parameter summaries describe wild type, AceK3 and AceK4 in kinase and phosphatase assays [31]. An ordered substrate-binding realization reproduces their common sequential initial-rate law exactly. A finite random-binding realization approaches the same law as binding becomes fast relative to catalysis. At a binding/catalysis ratio of 1,000, the largest relative difference is 0.00098–0.00114 over 625 dose pairs per summary (Fig 5A–B). Nevertheless, matched low-ATP protein-binding predictions differ by 0.495–0.497 in bound fraction (Fig 5C). The construction is calibrated to published kinetic summaries; it is not an independent fit to raw replicates or a validated complete AceK cycle.

Independent-activity data restrict the interpretation further. Adding ATPase only from the ATP-bound state preserves the ordered kinase rate law, but predicts ATPase halfpoints of 170, 580 and $5,300\ \mu\text{M}$ for wild type, AceK3 and AceK4. The withheld measured summaries are 230, 770 and $2,900\ \mu\text{M}$ [31]. This exposes an additional biochemical assumption that kinase agreement alone does not establish. Missing covariance prevents assigning significance to these discrepancies (E4). The general workflow therefore separates algebraic compatibility, kinetic calibration, and independent biological checks before proposing a discriminating assay.

The paired GlnB/GlnK reporter extension is retained in S7 Fig, A13–A14 and D6. Published peptide data provide conditional sequestration ceilings, but a matched no-binding calibration and direct bound-fraction endpoint are still required for calibrated reconstruction [5, 32–35]. This prospective extension is separate from the two network demonstrations above.




Fig 5. AceK extends the workflow to a second biochemical system. (A) Wild-type phosphatase rate law reconstructed from published parameter summaries, with ordered and finite random-binding realizations; the reference line is not raw experimental data. (B) Largest relative random-binding deviation over 625 ATP–protein input pairs for each of six source summaries. The dotted 1% line is a computational tolerance. Colors denote protein variants; solid and dashed lines denote kinase and phosphatase. (C) Protein-binding contrast at the source-derived apparent binding scale versus $\text{ATP}/K_{m, \text{ATP}}$, with binding/catalysis ratio 1,000. These are prospective predictions under the specified binding models. (D) General and observation-conditioned interval widths versus AceK total at $1 \mu\text{M}$ IDH dimers in the full dimeric-target topology. The count gate is $[1.435, 1.475]$ phosphates per dimer. The 0.05 tolerance is crossed at $0.05 \mu\text{M}$ AceK only after conditioning. The reporter is total unphosphorylated-dimer fraction, not measured physiological activity. Coefficients, dwell-time class and observation limits remain conditional.

Discussion

The main result is a method for deriving decision conditions despite mechanistic ambiguity. Complete stationary target distributions can leave enzyme-state mechanisms unresolved. Conservation and additional observations can nevertheless bound the predictions of a specified class. The nitrogen example requires a count gate, coefficient envelope, source-unit interpretation and stationarity; the AceK example shows how a graph enclosure can replace exact enzyme cancellation when additional enzyme-bound states are present.

These claims build on existing theory rather than replacing it. Prediction profile likelihood propagates a specified likelihood into prediction intervals even with non-identifiable parameters [22]. Model-discrimination design compares prospective observations under candidate models [19]. Our enclosure supplies a deterministic sufficient condition from a declared stationary graph, kinetic coefficients, protein conservation and bounded observations, without requiring a likelihood or choosing one microscopic member. It does not inherit frequentist coverage unless its input intervals have that coverage, and may be wider than a likelihood-based interval using richer information. The methods are complementary: a certificate can justify a decision shared by a class, while profile or discrimination methods can prioritize observations when the certificate is uninformative.

The expanded binding benchmark makes measurement controls part of this distinction. Condition-specific background and gain fits correct the tested affine signal distortions, whereas hidden affinity mixtures remain misleading under a single-class interpretation. The numerical calibration accounts for nonlinear fitting on a finite nuisance grid; it does not establish uniform performance outside that grid. GlnD profiles and the AceK ATPase check additionally show why fit quality within one assay does not establish the biochemical assumptions of another.

The biological scope is explicit. The graph construction requires one target per complex, fixed free ligands, irreducibility and positive-polynomial transitions; multiple targets in one complex or changing external pools require a different state description. The 120-species cascade verifies specified kinetics, not the full physiological network. ATP/ADP, metabolism, expression feedback and spatial modification arrangements can matter [36,37]. Published kinetic summaries and cellular activity measurements obtained under different conditions cannot supply a matched count gate. The framework consequently defines which measurements would certify a class-wide decision and which observations would challenge a proposed mechanism; it does not report an already validated native concentration choice.

Materials and methods

309

Reaction models and verification

310

Core models use positive mass-action rates, distributive catalytic release and explicitly stated regulatory exchange. Complex balances and zero flux across ladder cuts give the stationary distributions; derivatives and conservation yield the response and observation bounds. S1 Appendix supplies complete reaction schemes, proofs, parameter origins and boundary constructions. S2 Appendix supplies data provenance, fitting sensitivity and prospective assay precision; earlier exploratory analyses are preserved in repository documentation. Shared rate-rescaling constructions are not fitted independently to the same data and are not assigned model-selection probabilities.

311
312
313
314
315
316
317

The closed cascade enumerates 120 species and 300 directed reactions; an additional productive route gives a 156-species model. Stationary solutions reduce to conserved free-pool balances and are checked in the independently enumerated mass-action vector field. Conserved-input calculations add a scalar glutamine balance. Randomized comparisons, alternative recognition weights and independent BDF integrations test specified realizations. Dimensions, tolerances, seeds and parameter ranges are recorded in A10–A12 and S1 Data.

318
319
320
321
322
323

The earlier source regeneration included the original-workbook extraction and exploratory 8,700-trial benchmark. The present extension adds independently calibrated raw-assay simulations, exact graph-cofactor checks, observation linear programs and a second biochemical network. All 13 stages of the current integrated check passed, including 22 baseline checks, matched-model verification, three extension test groups, the expanded bootstrap audit and four generalization checks. The integrated entry point additionally checks 2,000 independent paired-reporter generators and 10,000 admissible interval configurations. The generalized calibration extension is checked against 3,000 independent mixture linear programs, 500 reporter-interval linear programs and 1,000 positive ladder pairs. Machine-readable reports distinguish clean regeneration from subsequent checks. They validate computational statements under the model assumptions, not experimental uncertainty or physiological correctness.

324
325
326
327
328
329
330
331
332
333

Published data and comparison rules

334

The first-layer analysis uses twelve author-supplied digitized coordinate pairs, retained with provenance. Bounded Hill and specified ladder fits use identical observations. Corrected AIC includes the residual variance parameter under a working independent Gaussian model; profile calculations follow established practice [38]. Coordinate perturbations use prespecified input/output scales, not estimated biological errors.

335
336
337
338

Two thousand refits assess coordinate sensitivity and 300 shared perturbations propagate state predictions. 339
 Historical downstream summaries receive shared-parameter comparisons without inventing unreported 340
 titration coordinates or likelihoods. Numerical and source details are in S2 Appendix and the repository 341
 extended-analysis archive. 342

The AceK analysis transcribes six kinetic summaries and three withheld ATPase halfpoints from Miller et 343
 al. [31]. Each initial-rate realization is checked against the published sequential law; no reconstructed curve is 344
 labeled raw data. The full dimeric-target topology and reference rates follow Dexter and Gunawardena [30]. 345
 Separately transcribed in vivo IDH activity data [39] are not converted from units/mg to micromolar 346
 concentrations or treated as matched reconstitution measurements. E3–E4 document the assay and 347
 calibration limits. 348

The independent Gosztolai et al. workbook and model are retained unchanged from 349
 doi:10.6084/m9.figshare.4880003. Extraction records source cells, means, SE and sample counts. 350
 Paired-peptide fractions divide modified signal by modified plus unmodified signal; they differ from 351
 normalization by a separately measured total. First-order SE uses the conservative sum of absolute 352
 contributions because covariance is unavailable. It does not establish joint confidence coverage. Raw 353
 copy-per-cell values are preserved; the source model corresponds to $0.00161 \mu\text{M}$ per copy per cell, and the 354
 workbook’s inconsistent printed unit is documented. 355

Prospective observation design 356

State-distribution designs maximize a disclosed candidate-dependent distance. The new GlnE benchmark 357
 generates finite-depletion observations, fits three halfpoints and either shared or condition-specific signal 358
 parameters, and tests the average log-ratio score using variance across eight complete experiments. The 359
 shared zero-glutamine measurement is retained within each score. Fixed-null parametric calibration repeats 360
 the raw nonlinear fit at three K_G values, three concentration-error scales and three signal-error scales. 361
 Per-cell order statistics provide simultaneous 95% Monte Carlo confidence for the nominal 5% tail over the 362
 declared 27-point grid; independent evaluation reports Wilson intervals. This is finite-grid calibration, not a 363
 data-dependent plug-in bootstrap or a continuous-nuisance theorem. Trials with failed or active-bound fits 364
 do not reject and remain in the denominator. Full doses, error magnitudes, mixture path, seeds and cutoff 365
 construction are in E5 and S1 Data. 366

The count-conditioned bound is tested with arbitrary positive ladders, independently formed protein 367
 mixtures and full conserved reaction realizations. Paired-reporter intervals require their calibration and 368
 uncertainty sets to hold jointly. D4–D6 and A11–A14 separate reported values, source-fit examples, chosen 369

scenarios and measurements required for application.

370

Data and code availability

371

No new experimental measurements were generated. Source observations, digitized coordinates, parameter records, prediction grids and verification outputs are supplied as S1 Data and in GitHub release v1.4.0. The version-independent Zenodo concept DOI, doi:10.5281/zenodo.22731768, indexes the versioned source archive; use version 1.4.0 for this manuscript. The submission manifest records the verified archive record and release hashes. The earlier v1.3.4 archive does not contain the new analyses. The independent source dataset is available at doi:10.6084/m9.figshare.4880003. S1 Data retains source licenses and attribution.

372

373

374

375

376

377

Acknowledgments

378

I would like to express my sincere gratitude to Dr Seonhwa Park for introducing me to the practice of research and for her guidance and encouragement.

379

380

Author contributions

381

Yule Choi: conceptualization, methodology, formal analysis, software, investigation, validation, visualization, writing – original draft, writing – review and editing.

382

383

Competing interests

384

The author declares no competing interests.

385

References

386

1. Goldbeter A, Koshland DE. An amplified sensitivity arising from covalent modification in biological systems. *Proc Natl Acad Sci USA*. 1981;78:6840–6844. doi:10.1073/pnas.78.11.6840.
2. Ferrell JE, Ha SH. Ultrasensitivity part III: cascades, bistable switches, and oscillators. *Trends Biochem Sci*. 2014;39(12):612-618. doi:10.1016/j.tibs.2014.10.002.

387

388

389

390

3. van Heeswijk WC, Westerhoff HV, Boogerd FC. Nitrogen Assimilation in *Escherichia coli*: Putting
Molecular Data into a Systems Perspective. *Microbiol Mol Biol Rev.* 2013;77(4):628-695.
doi:10.1128/mmmbr.00025-13.
4. Ninfa AJ, Atkinson MR. PII signal transduction proteins. *Trends Microbiol.* 2000;8:172–179.
doi:10.1016/s0966-842x(00)01709-1.
5. Gosztolai A, Schumacher J, Behrends V, Bundy JG, Heydenreich F, Bennett MH, et al. GlnK
Facilitates the Dynamic Regulation of Bacterial Nitrogen Assimilation. *Biophys J.* 2017;112:2219–2230.
doi:10.1016/j.bpj.2017.04.012.
6. Jiang P, Ninfa AJ. A Source of Ultrasensitivity in the Glutamine Response of the Bicyclic Cascade
System Controlling Glutamine Synthetase Adenylylation State and Activity in *Escherichia coli*.
Biochemistry. 2011;50(50):10929-10940. doi:10.1021/bi201410x.
7. Ventura AC, Jiang P, Van Wassenhove L, Del Vecchio D, Merajver SD, Ninfa AJ. Signaling properties
of a covalent modification cycle are altered by a downstream target. *Proc Natl Acad Sci U S A.*
2010;107(22):10032-10037. doi:10.1073/pnas.0913815107.
8. Mutalik VK, Shah P, Venkatesh KV. Allosteric Interactions and Bifunctionality Make the Response of
Glutamine Synthetase Cascade System of *Escherichia coli* Robust and Ultrasensitive. *J Biol Chem.*
2003;278(29):26327-26332. doi:10.1074/jbc.m300129200.
9. Bruggeman FJ, Boogerd FC, Westerhoff HV. The multifarious short-term regulation of ammonium
assimilation of *Escherichia coli*: dissection using an in silico replica. *FEBS J.* 2005;272(8):1965-1985.
doi:10.1111/j.1742-4658.2005.04626.x.
10. Ortega F, Ehrenberg M, Acerenza L, Westerhoff HV, Mas F, Cascante M. Sensitivity Analysis of
Metabolic Cascades Catalyzed by Bifunctional Enzymes. *Mol Biol Rep.* 2002;29(1-2):211-215.
doi:10.1023/a:1020334014497.
11. Ortega F, Acerenza L, Westerhoff HV, Mas F, Cascante M. Product dependence and bifunctionality
compromise the ultrasensitivity of signal transduction cascades. *Proc Natl Acad Sci USA.*
2002;99:1170–1175. doi:10.1073/pnas.022267399.
12. Straube R. Sensitivity and Robustness in Covalent Modification Cycles with a Bifunctional Converter
Enzyme. *Biophys J.* 2013;105(8):1925-1933. doi:10.1016/j.bpj.2013.09.010.

13. Straube R. Reciprocal Regulation as a Source of Ultrasensitivity in Two-Component Systems with a Bifunctional Sensor Kinase. *PLoS Comput Biol.* 2014;10(5):e1003614. doi:10.1371/journal.pcbi.1003614.
14. Jiang P, Zhang Y, Atkinson MR, Ninfa AJ. The Robustness of the Escherichia coli Signal-Transducing UTase/UR-PII Covalent Modification Cycle to Variation in the PII Concentration Requires Very Strong Inhibition of the UTase Activity of UTase/UR by Glutamine. *Biochemistry.* 2012;51(45):9032-9044. doi:10.1021/bi3005736.
15. Thomson M, Gunawardena J. The rational parameterisation theorem for multisite post-translational modification systems. *J Theor Biol.* 2009;261:626–636. doi:10.1016/j.jtbi.2009.09.003.
16. Feliu E, Wiuf C. Variable elimination in post-translational modification reaction networks with mass-action kinetics. *J Math Biol.* 2013;66(1-2):281-310. doi:10.1007/s00285-012-0510-4.
17. Feliu E, Wiuf C. Variable Elimination in Chemical Reaction Networks with Mass-Action Kinetics. *SIAM J Appl Math.* 2012;72:959–981. doi:10.1137/110847305.
18. Xu Y, Gunawardena J. Realistic enzymology for post-translational modification: Zero-order ultrasensitivity revisited. *J Theor Biol.* 2012;311:139–152. doi:10.1016/j.jtbi.2012.07.012.
19. Silk D, Kirk PDW, Barnes CP, Toni T, Stumpf MPH. Model Selection in Systems Biology Depends on Experimental Design. *PLoS Comput Biol.* 2014;10(6):e1003650. doi:10.1371/journal.pcbi.1003650.
20. Castro M, de Boer RJ. Testing structural identifiability by a simple scaling method. *PLoS Comput Biol.* 2020;16(11):e1008248. doi:10.1371/journal.pcbi.1008248.
21. Villaverde AF, Massonis G. On testing structural identifiability by a simple scaling method: Relying on scaling symmetries can be misleading. *PLoS Comput Biol.* 2021;17(10):e1009032. doi:10.1371/journal.pcbi.1009032.
22. Kreutz C, Raue A, Timmer J. Likelihood based observability analysis and confidence intervals for predictions of dynamic models. *BMC Syst Biol.* 2012;6:120. doi:10.1186/1752-0509-6-120.
23. Zhang Y, Pohlmann EL, Serate J, Conrad MC, Roberts GP. Mutagenesis and Functional Characterization of the Four Domains of GlnD, a Bifunctional Nitrogen Sensor Protein. *J Bacteriol.* 2010;192(11):2711-2721. doi:10.1128/jb.01674-09.

24. Jiang P, Peliska JA, Ninfa AJ. Enzymological Characterization of the Signal-Transducing Uridylyltransferase/Uridylyl-Removing Enzyme (EC 2.7.7.59) of *Escherichia coli* and Its Interaction with the PII Protein. *Biochemistry*. 1998;37(37):12782-12794. doi:10.1021/bi980667m.
25. Grant GA. The ACT Domain: A Small Molecule Binding Domain and Its Role as a Common Regulatory Element. *J Biol Chem*. 2006;281:33825-33829. doi:10.1074/jbc.r600024200.
26. Jiang P, Mayo AE, Ninfa AJ. *Escherichia coli* glutamine synthetase adenylyltransferase (ATase, EC 2.7.7.49): kinetic characterization of regulation by PII, PII-UMP, glutamine, and alpha-ketoglutarate. *Biochemistry*. 2007;46:4133-4146. doi:10.1021/bi0620510. Supporting information: doi:10.1021/bi0620510.s001.
27. Adair GS, Bock AV, Field H Jr. The hemoglobin system: VI. The oxygen dissociation curve of hemoglobin. *J Biol Chem*. 1925;63(2):529-545. doi:10.1016/S0021-9258(18)85018-9.
28. Gunawardena J. Distributivity and processivity in multisite phosphorylation can be distinguished through steady-state invariants. *Biophys J*. 2007;93(11):3828-3834. doi:10.1529/biophysj.107.110866.
29. Gunawardena J. A linear framework for time-scale separation in nonlinear biochemical systems. *PLoS One*. 2012;7(5):e36321. doi:10.1371/journal.pone.0036321.
30. Dexter JP, Gunawardena J. Dimerization and bifunctionality confer robustness to the isocitrate dehydrogenase regulatory system in *Escherichia coli*. *J Biol Chem*. 2013;288(8):5770-5778. doi:10.1074/jbc.M112.339226.
31. Miller SP, Karschnia EJ, Ikeda TP, LaPorte DC. Isocitrate dehydrogenase kinase/phosphatase: kinetic characteristics of the wild-type and two mutant proteins. *J Biol Chem*. 1996;271(32):19124-19128. doi:10.1074/jbc.271.32.19124.
32. Durand A, Merrick M. In vitro analysis of the *Escherichia coli* AmtB-GlnK complex reveals a stoichiometric interaction and sensitivity to ATP and 2-oxoglutarate. *J Biol Chem*. 2006;281(40):29558-29567. doi:10.1074/jbc.M602477200.
33. van Heeswijk WC, Wen D, Clancy P, Jaggi R, Ollis DL, Westerhoff HV, et al. The *Escherichia coli* signal transducers PII (GlnB) and GlnK form heterotrimers in vivo: fine tuning the nitrogen signal cascade. *Proc Natl Acad Sci USA*. 2000;97(8):3942-3947. doi:10.1073/pnas.97.8.3942.
34. Cong X, Liu Y, Liu W, Liang X, Laganowsky A. Allosteric modulation of protein-protein interactions by individual lipid binding events. *Nat Commun*. 2017;8:2203. doi:10.1038/s41467-017-02397-0.

35. Radchenko MV, Thornton J, Merrick M. Association and dissociation of the GlnK–AmtB complex in response to cellular nitrogen status can occur in the absence of GlnK post-translational modification. *Front Microbiol.* 2014;5:731. doi:10.3389/fmicb.2014.00731.
36. Callenbach A, Dorešić D, Düster R, Nakonecnij V, Dudkin E, Geyer M, et al. Quantitative modelling of P-TEFb mediated CTD phosphorylation identifies local cooperativity. *PLoS Comput Biol.* 2026;22(7):e1014531. doi:10.1371/journal.pcbi.1014531.
37. Emori MT, Tomazini LF, Souza EM, Pedrosa FO, Chubatsu LS, Oliveira MAS. The deuridylylation activity of *Herbaspirillum seropedicae* GlnD protein is regulated by the glutamine:2-oxoglutarate ratio. *Biochim Biophys Acta Proteins Proteom.* 2018;1866(12):1216–1223. doi:10.1016/j.bbapap.2018.09.009.
38. Raue A, Kreutz C, Maiwald T, Bachmann J, Schilling M, Klingmüller U, et al. Structural and practical identifiability analysis of partially observed dynamical models by exploiting the profile likelihood. *Bioinformatics.* 2009;25(15):1923–1929. doi:10.1093/bioinformatics/btp358.
39. LaPorte DC, Thorsness PE, Koshland DE Jr. Compensatory phosphorylation of isocitrate dehydrogenase: a mechanism for adaptation to the intracellular environment. *J Biol Chem.* 1985;260(19):10563–10568. doi:10.1016/S0021-9258(19)85122-0.

Supporting information

S1 Appendix. Reaction mechanisms, conservation bounds and proofs (A1–A15).

S2 Appendix. Data provenance, biochemical constraints and sensitivity analyses (D1–D7).

S3 Appendix. General algorithm, AceK evidence, expanded calibration and paired-paralog extension (E1–E7).

S1–S9 Figs. Supplementary analyses, raw-assay diagnostics and nuisance grids; individual captions accompany the files.

S1 Data. Source measurements, parameter records, code, numerical results and verification reports.