SHEET AND AND	tipsychotics	Brand name	Details	Side effects	
	1st gen			lower seizure threshold, prolong QT, NMS	
W J F	laloperidol	Haldal	anti D2		
	uphenazine	Haldol Prolixin	high potency. Available in depot	more EPS, less (HAM), galactorrea	
of Chi	lorpromazine	Thorazine/ Largactil	high potency. Available in depot	to con and that social is alderly a falls	
SEX SERVICE	2nd gen	The care of the ca	low potency	less EPS, more HAM. Avoid in elderly -> falls	
nut (Olanzapine	Zyprexa	anti D2 and 5HT2	metabolic syndrome most wt gain and sedation	
	Clozapine	Scany lupe Clozaril	Dec suicide, effective. Good for pts w/TD	agranulocytosis, sz, cardiomyopathy, hypoTN. Highest risk of met syd. Makes pts drool. Level will inc with smoking cessation	
	Risperidone	Risperdal 5Ht-	most similar to typicals. Depot	orthostasis Similar SE profile to Haldol prolaction took	
	Quetiapine	solon Seroquel	Better for dementia pts b/c not as much dopa blockade so less motor SE	orthostasis and sedation (HI effects). Cataracts sudden death in elderly w/ Jamenta related psychosis	of odn.
	Ziprasidone	meets and (C Geodon	Slow titration, must take w/ food	[MI.GT (EKG) least likely to cause met. syd.	Lemans!
V .	Aripiprazole	Abilify	partial DA agonist. Augmentation in MDD.	Akathesias. May be activating.	J. O
- A Description	THE RESERVE THE PROPERTY OF THE PARTY OF THE	TV ads		Et H. Sture	
(MAOIs		MADA = SHT/NE (antidept) MADB = DA (antiPD)	avoid tyramine (aged/cured foods/beans) -> will cause HTN emergency from NE; serotonin synd w/ SSRIs {2 wk washout} CANNOT GIVE W/ MEPERIDINE, TRAMADOL, or DEX Need to wait 5 weeks w/ fluoxetine Plugher dozes Plug	- desmay
	Phenilizine	Nardil	Anxious depression	CANNOT GIVE W/ MEPERIDINE, TRAMADUL, OF DEX - Supp.	, 0
T	ranylcypromine	Parnate /	Melancholic depression	Need to wait 5 weeks w/ Huoxetine no cheese w	me, favo
-	Selegiline	Emsam/Eldepryl	Selective for MAOb nonsellachic	(eluquer doses fatal in OD (torsade de pointe) stous cardiac conduction	
5	TCAs		incr SHT, NE	Serotonin Synd: Supportive care, benzos as needed	
3	Amitriptyline	Elavil/ Endep	used for pain syndromes		
	Clomipramine	Anafranil	used for OCD	Sent + Nepi	
	Imipramine	Tofranil	used for urinary urgency/enuresis		
	Nortriptyline	Pamelor/Aventyl/ Noritren	used for elderly depression	diastolic HTN (due to NE), SSRI side effects	
	SNRI		use for comorbid pain syd	5179. rate of HTN] India for	pain.
3	Venlafaxine	Effexor			
	Duloxetine	Cymblata	MANAGEM CONTRACTOR STREET	Ann Their Brook	= gr
CALCULATE AND ADDRESS OF THE PARTY OF THE PA	SSRI		2nd Shortest acting Short 2 life-	Worst discontinuation syndrome WAYTSEW	
D	Paroxetine	Paxil	2110 3110 110	@ inten of warfann is so now mounted 2 and was	ant
	Citalopram	Celexa	LAL I	PTATES 2 WK Was	A PENUL
	Escitalopram	Lexapro	longest half life	Howass seizure threshold in eating disorders. Can potentially make	T WOULD
	Fluoxetine	Prozac	Shortest acting spector + MC) Ind	sucola
	Fluvoxamine	Luvox	4 800	4	
	Sertraline	Zoloft		Sfx:	sexual
GEOGRAPH .	Other		NDRI. Fewer studies of efficacy than SSRI. Good for	Lowers seizure times and an additional and an additional and an additional and additional additional and additional and additional and additional	+91
Contract of	Washington State of the A	Wellbutrin/Zyban	smoking, less sexual dysfx. with los	2 M. anxious/finery	
	Bupropion		Thrane Sincr NE and St	HT COURT SUITE 1/2 offects Very selective for 5HT1	
	Mirtazapine	Remeron/Avanza/Zispin	secretion. Promotes sleep/appetite by checks shall be secretion.	& Rxisudated	
	Trazodone		agonist 5HT1A) kup Joe Partial 5HT1A agonist, more used in GAD	o hot be for deposition praprism 7 this so wast monitor FT	sur.
	Buspirone	Buspar Ionhanto anarolytic			

or puts brakes on serotomin,

7 B metabolism NSAIDS inc Lith Levels els. 15-2 >2 72.5 tremor nauxo atania diarrho -> 32 -> ARF -> coma + death Mood stabilizers Lithium neurotoxicity, thyroid dysfn, DI,Ebstein's abnormality (fetal cardio use in pregnancy. Decreases suicide. (.8-1.2) augmentation for depression effects) SA nodesick sinces. antronvulsants autoinducer of metabolism at liver (decr concen of Carbamazepine Tegretol SIS rash, CBC abnl, fetal craniofacial defects INEY blood skin CYP3A4 self and VPA) (8-12) John hepatitis, pancreatitis, CBC abnl, sedation, wt gain, alopecia, fetal Valproic acid Depakote V Can be loaded, available IV. (80-120) Thumbout agains neural tube defects Lamotrigine Lamictal OF T SJS rash (so titrate SLOW) Worse than corb Not indicated for acute mania Benzos GAU as topion this cause) teletice periodical thin avenit Alprazolam Xanax Highly abused Quick onset, short duration Oprazepam Equipotent IM/PO/IV Ativan use in liver dz. Long onset, avg duration. Clonazepam Klonipin Long onset/duration, low risk of abuse xazepam Serax use in liver dz Temazepam Restoril use in liver dz, for sleep Dementia the set one restorming the set of the set one restorment on the set on the se Gi upset, bradycardia Donepezil Aricept Central ACH inhibitor, used in Alz Headache, dizziness, insomnia Memantine Namenda NMDA-type glutamate antagonist in Alz Used in Parkinsons Levadopa sleep attacks Dopa agonist, used in parkinsons/RLS Requip Roperinole Dopa agonist, used in parkinsons/RLS Mirapex Pramipexole Etoh Cannot use in renal pts NMDA antagonist/GABA agonist Campral Acamprosate Liver damage, GI upset, will cause withdraw Opiod antagonist. Also used in opiod dep Vivitrol Naltrexone HA, can cause neurotoxicity/EPS inhibits acetaldehyde dehydrogenase Antabuse Disulfiram Opiate Similar to taking opiods Synthetic opiod Methadone Harder to abuse Suboxone Bupren.+ Naloxone Similar to taking opiods, liver damage Semi-synthetic opiod Buprenorphine Acute withdrawal Opiod antagonist. Use in OD. Naloxone Rebound HTN, causes hypoTN Central alpha2 agonist Clonidine Stimulants Do not give w/ TCA, or pts w/ HTN/Sz/Arrythmias Inhibit NE/D reuptake. Use in ADHD Ritalin Methylphenidate SJS, Induces P450 Used in narcolopsy/fatigue Provigil Modafinil Not stimulant, NRI used in ADHD Strattera Atomoxetine prodrug to dextroamphetamine Vyvanse Lisdexamfetamine enantiomer of the amphetamine Adderall Dextroamphetamine Autionivitants excapateprie hyponatrem 2.5%. toleptal

GARA activity nonberge Blyiden Zaliphlon

Ambien

abuse potential relators